

An observational, prospective, two-cohort comparison of a fixed versus variable dosing strategy of prothrombin complex concentrate to counteract vitamin K antagonists in 240 bleeding emergencies

Nakisa Khorsand,¹ Nic J.G.M. Veeger,² Reinier M. van Hest,^{1,3} Paula F. Ypma,⁴ Jeroen Heidt,⁵ and Karina Meijer⁶

¹Department of Hospital Pharmacy, Haga Teaching Hospital, The Hague; ²Department of Epidemiology, University Medical Centre, Groningen; ³Department of Clinical Pharmacy, Academic Medical Center, Amsterdam; ⁴Department of Haematology, Haga Teaching Hospital, The Hague; ⁵Department of Internal Medicine, Medical Centre Haaglanden, The Hague; and ⁶Division of Haemostasis and Thrombosis, Department of Haematology, University Medical Centre Groningen, The Netherlands

Citation: Khorsand N, Veeger NJGM, van Hest RM, Ypma PF, Heidt J, and Meijer K. An observational, prospective, two-cohort comparison of a fixed versus variable dosing strategy of prothrombin complex concentrate to counteract vitamin K antagonists in 240 bleeding emergencies. *Haematologica* 2012;97(10):1501-1506. doi:10.3324/haematol.2012.063701

Online Supplementary Table S1. PCC variable dosing regimen

Initial INR →	7.5	5.9	4.8	4.2	3.6	3.3	3.0	2.8	2.6	2.5	2.3	2.2
↓ Body weight (kg)												
Target 50	1040	1040	1040	780	780	780	520	520	X	X	X	X
INR 60	1300	1300	1040	1040	780	780	780	520	X	X	X	X
≤2.1	70	1560	1300	1300	1300	1040	1040	780	780	X	X	X
	80	1560	1560	1560	1300	1300	1040	1040	780	X	X	X
	90	1560	1560	1560	1560	1300	1300	1040	780	X	X	X
100	1560	1560	1560	1560	1560	1300	1040	1040	X	X	X	X
Target 50	1560	1560	1560	1300	1300	1300	1040	1040	780	780	780	780
INR 60	2080	1820	1820	1560	1560	1560	1300	1300	1040	1040	1040	780
≤1.5	70	2340	2080	2080	1820	1820	1820	1560	1560	1300	1040	1040
	80	2600	2600	2340	2340	2340	2080	2080	1820	1560	1300	1300
	90	2600	2600	2600	2340	2340	2340	2080	2080	1820	1560	1300
100	2600	2600	2600	2600	2600	2340	2340	2080	1820	1820	1560	1300

The dosage is shown as International Units of Factor IX, based on a Cofact® batch with 26 IU of F IX per mL as used for this study. This table is based on the manufacturer's algorithm (Sanquin, Amsterdam, The Netherlands).