

Manuscript no. HAEMATOL/2012/066217 entitled "Increased plasma thrombopoietin levels in patients with myelodysplastic syndrome: a reliable marker for a benign subset of bone marrow failure"

Authors: Yu Seiki, Yumi Sasaki, Kohei Hosokawa, Chizuru Saito, Naomi Sugimori, Hirohito Yamazaki, Akiyoshi Takami, and Shinji Nakao

Information about the contributions of each person named as having participated in the study

1) Guarantor(s), i.e., person(s) who is (are) responsible for the integrity of the work as a whole: • Shinji Nakao, Cellular Transplantation Biology, Kanazawa University Graduate School of Medical Science, Kanazawa, Japan, snakao8205@staff.kanazawa-u.ac.jp

According to the International Committee of Medical Journal Editors (ICMJE) (http://www.icmje.org/ethical_1author.html): "Authorship credit should be based on: 1) substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; 2) drafting the article or revising it critically for important intellectual content; and 3) final approval of the version to be published. Authors should meet conditions 1, 2, and 3 Acquisition of funding, collection of data, or general supervision of the research group alone does not constitute authorship".

The guarantors of this manuscript confirm that all persons designated as authors qualify for authorship, and that each author has participated sufficiently in the work to take public responsibility for appropriate portions of the content.

2) Authors who participated in the conception of the study: Shinji Nakao

3) Design & Methods. The following authors were responsible for specific investigations:

- Yu Seiki was responsible for measuring plasma TPO levels of the patients
- Yumi Sasaki was responsible for clonality analysis
- Yu Seiki, Kohei Hosokawa, Chizuru Saito, Naomi Sugimori, and Hirohito Yamazaki were responsible for detecting GPI-AP⁺ cells in patients with MDS and collecting clinical data of the patients

4) Results. The following authors were responsible for specific portions of the results, including figures and tables:

- Yu Seiki was responsible for all tables and figures including supplementary figures
- Yumi Sasaki was responsible for supplementary figure S1
- Akiyoshi Takami was responsible for statistical analyses

5) Writing the manuscript. The following authors were responsible for writing the manuscript:

- Shinji Nakao and Yu Seiki were responsible for writing the manuscript.

6) Contributors Listed in Acknowledgments:

We are deeply indebted to Ms. Rie Ooumi for technical assistance and to the following physicians for providing patients' data; A. Urabe of NTT Kanto Hospital; T. Endo and T. Kondo of the Hokkaido University Hospital; Y. Kataoka of the Ikeda City Hospital; M. Kofune of the Sapporo Medical University Hospital; T. Ueki, D. Akabane and H. Kobayashi of Nagano Red Cross Hospital; A. Sawazaki of NTT Kanazawa Hospital; A. Matsuda of Saitama Medical School Hospital; H. Fuse of Matsudo City Hospital; M. Ueda of Ishikawa Prefectural Hospital; S. Yoshida of Miyazaki Prefectural Miyazaki Hospital; K. Wakasa of Aiiku Hospital; R. Matsuoka and K. Kawakami of Kagawa Prefectural Central Hospital; T. Masunari and T. Kiguchi of Chugoku Central Hospital; R. Imamura of Kurume University Hospital; S. Matano of Tonami General Hospital; T. Sekine of Matsuzaka Central Hospital; A. Nishiyama of Ichinomiya Municipal Hospital; M. Iino of Yamanashi Prefectural Central Hospital; M. Tai of Municipal Tsuruga Hospital; K. Yoshinaga of Tokyo

Women's Medical University Hospital; N. Seno of Shinshu University Hospital; K. Ito of Ogawa Red Cross Hospital; M Masuya of Mie University Hospital; N. Aotsuka of Narita Red Cross Hospital; Y. Terasaki of Toyama City Hospital; T.Yoshida and T. Kurokawa of Toyama Prefectural Central Hospital.