

Online Supplementary Table S1. Age-related normal values of immunoglobulin levels¹

Age group	IgM (g/L)	IgG (g/L)	IgA (g/L)	
newborn	0,06–0,3	6,1–15,4	0,01–0,04	
3-6 months	0,3–1,0	2,5–5,6	0,05–0,5	
6 months-1 year	0,3–1,0	2,5–6,7	0,08–0,7	
1-2 year	0,4–1,7	3,3–11,6	0,1–1,0	
2–6 year	0,5–1,8	4,0–11,0	0,1–1,6	
7–12 year	0,5–2,0	6,0–12,3	0,3–2,0	
Adults	0,4–2,3	7,0–16,0	0,7–4,0	
	IgG1	IgG2	IgG3	IgG4
0-1 month	2,4-10,6	0,8-4,1	0,14-0,55	0,04-0,6
1-6 month	1,8-7,0	0,4-2,1	0,14-0,70	<0,03-0,4
6 month-2 years	2,2-8,2	0,4-2,4	0,15-1,0	<0,03-0,6
2-7 years	3,5-10,0	0,6-3,5	0,14-1,3	<0,03-1,2
7-18 years	3,8-10,0	0,9-5,0	0,15-1,5	<0,03-2,1

Reference

1. De Vries E, Kuijpers TW, Van Tol MJ, Van der Meer JW, Weemaes CM, Van Dongen JJ. Immunology in medical practice. XXXV. Screening of suspected immunodeficiency: diagnostic protocols for patients with opportunistic or recurrent severe infections, wasting and failure to thrive. *Ned Tijdschr Geneesk* 2000;144:2197-203.