

## APRIL levels are associated with disease activity in human chronic graft-versus-host disease

François Chasset,<sup>1,2,3</sup> Adèle de Masson,<sup>1,2,3</sup> Hélène Le Buane,<sup>1,2</sup> Aliénor Xhaard,<sup>2,4</sup> Flore Sicre de Fontbrune,<sup>2,4</sup> Marie Robin,<sup>2,4</sup> Michel Rybajad,<sup>2,3</sup> Nathalie Parquet,<sup>2,4,5</sup> Anne C. Brignier,<sup>2,4,5</sup> Tereza Coman,<sup>2,4</sup> Djaouida Bengoufa,<sup>2,6</sup> Anne Bergeron,<sup>2,7</sup> Régis Peffault de Latour,<sup>2,4</sup> Martine Bagot,<sup>1,2,3</sup> Armand Bensussan,<sup>1,2\*</sup> Gérard Socié,<sup>2,4,8\*</sup> and Jean-David Bouaziz<sup>1,2,3\*</sup>

\*Have equally contributed to that work and share last author seniorship

<sup>1</sup>INSERM and UMRS 976, Laboratory of Onco dermatology, Immunology, and Cutaneous Stem cells, F-75739 Paris; <sup>2</sup>University Paris Diderot, Sorbonne Paris Cité; <sup>3</sup>Dermatology Department, Saint-Louis Hospital, Paris; <sup>4</sup>Hematology and Transplantation, Saint-Louis Hospital, Paris; <sup>5</sup>Therapeutic Apheresis Unit, Saint-Louis Hospital, Paris; <sup>6</sup>Immunobiology Department, Saint-Louis Hospital, Paris; and <sup>7</sup>Pneumology Department, Saint-Louis Hospital, Paris; <sup>8</sup>INSERM and UMRS1160, France

The online version of this letter has a Supplementary Appendix.

Correspondence: gerard.socie@aphp.fr  
doi:10.3324/haematol.2016.145409

**Supplemental table 1: Antibodies used for cell surface and intracellular flow cytometry staining**

Specificity	Fluorophore	Clone	Manufacturer
Fixable Viability	eFluor® 506	NA <sup>1</sup>	eBioscience®
CD19	PerCP-Cy5.5	Hib19	eBioscience®
CD27	APC-eFluor 780	LG7F9	eBioscience®
CD24	FITC	ML5	BD Pharmigen®
CD38	APC	HIT2	BD Pharmigen®
CD267 TACI	PE	1A1	Miltenyi Biotec®
CD269 BCMA	APC	REA31	Miltenyi Biotec®
BAFF-R	FITC	7H22-E16	Miltenyi Biotec®


NA<sup>1</sup>: Not applicable

**Supplemental table 2. Clinical and biological parameters associated with High APRIL levels**

Variables	Low APRIL levels	High APRIL levels	p value (univariate analyze)	p value (multivariate analyze)
Severe cGVHD; n (%)	18 (40.9)	10 (90.9)	0.005*	0.03*
Lichenoid cGVHD; n (%)	21 (48.8)	6 (66.6)	0.46	-
Sclerodermiform cGVHD; n (%)	20 (46.5)	2 (22.2)	0.27	-
Ocular cGVHD; n (%)	12 (29.3)	3 (33.3)	1	-
Oral mucosa cGVHD; n (%)	20 (50)	2 (22.2)	0.26	-
Digestive tract cGVHD; n (%)	5 (11.3)	2 (18.1)	0.61	-
Hepatic cGVHD; n (%)	9 (20.4)	3 (27.27)	0.68	-
Pulmonary cGVHD; n (%)	8 (18)	6 (54.5)	0.02*	0.2
ANA +; n (%)	25 (60.9)	11 (100)	0.01*	0.9
Corticosteroids; n (%)	32 (78)	6 (54.5)	0.14	0.07

\* Statistically significant; ANA: Antinuclear autoantibody

**Supplemental Figure 1. Correlation between circulating BAFF and APRIL levels and B cell subsets and association with antinuclear antibodies.**



**Supplemental Figure 1. Correlation between BAFF and APRIL circulating levels and B cell subsets and association with antinuclear antibodies.**

**(A).** Representative flow cytometry dot plot gating on CD19<sup>+</sup> B cell populations according to CD24 and CD38 immunostaining. **(B).** Correlation between circulating BAFF levels and B-cell subsets among no cGVHD patients. **(C).** Correlation between circulating APRIL levels and B-cell subsets among cGVHD patients. **(D).** Median APRIL circulating levels between patients with and without antinuclear antibodies (ANA) and frequency of ANA positivity between patients with circulating APRIL levels > or < 83 ng/ml. Correlations were measured using the Spearman test: \* p<0.05, \*\* p<0.01, \*\*\* p<0.001, NS: Non-significant