

Haematologica
HAEMATOL/2018/203166
Version 4

Unraveling the effect of silent, intronic and missense mutations on VWF splicing: contribution of next generation sequencing in the study of mRNA.

Nina Borràs, Gerard Orriols, Javier Batlle, Almudena Pérez-Rodríguez, Teresa Fidalgo, Patricia Martinho, María Fernanda López-Fernández, Ángela Rodríguez-Trillo, Esther Lourés, Rafael Parra, Carme Altisent, Ana Rosa Cid, Santiago Bonanad, Noelia Cabrera, Andrés Moret, María Eva Mingot-Castellano, Nira Navarro, Rocío Pérez-Montes, Sally Marcellini, Ana Moreto, Sonia Herrero, Inmaculada Soto, Núria Fernández-Mosteirín, Víctor Jiménez-Yuste, Nieves Alonso, Aurora de Andrés-Jacob, Emilia Fontanes, Rosa Campos, María José Paloma, Nuria Bermejo, Ruben Berrueco, José Mateo, Karmele Arribalzaga, Pascual Marco, Ángeles Palomo, Nerea Castro Quismondo, Belén Iñigo, María del Mar Nieto, Rosa Vidal, María Paz Martínez, Reyes Aguinaco, Jesús María Tenorio, María Ferreiro, Javier García-Frade, Ana María Rodríguez-Huerta, Jorge Cuesta, Ramón Rodríguez-González, Faustino García-Candel, Manuela Dobón, Carlos Aguilar, Francisco Vidal, and Irene Corrales

Disclosures: None.

Contributions: Contribution: This study derived from PCM-EVW-ES registry. I.C. and F.V coordinated the study; N.B and G.O selected the patients included in the study; J.B, A.P.R, M.F.L.F, T.F, P.M, R.P and C.A obtained patient's sample; N.B, G.O, F.V. and I.C. analyzed and interpreted results; N.B, F.V. and I.C. wrote the paper; and all the authors that participated in the PCM-EVW-ES registry checked the final version of the manuscript.