



Journal of The Ferrata Storti Foundation

Donor cell leukemia: reappearance of gene mutations in donor cells - more than an incidental phenomenon?

by Tal Shahar Gabay, Noa Chapal-Ilani, Yoni Moskovitz, Tamir Biezuner, Barak Oron, Yardena Brilon, Anna Fridman-Dror, Rawan Sabah, Ran Balicer, Amos Tanay, Netta Mendelson-Cohen, Eldad J. Dann, Riva Fineman, Nathali Kaushansky, Shlomit Yehudai-Reshef, Tsila Zuckerman, and Liran I. Shlush

Haematologica 2020 [Epub ahead of print]

Citation: Tal Shahar Gabay, Noa Chapal-Ilani, Yoni Moskovitz, Tamir Biezuner, Barak Oron, Yardena Brilon, Anna Fridman-Dror, Rawan Sabah, Ran Balicer, Amos Tanay, Netta Mendelson-Cohen, Eldad J. Dann, Riva Fineman, Nathali Kaushansky, Shlomit Yehudai-Reshef, Tsila Zuckerman, and Liran I. Shlush. Donor cell leukemia: reappearance of gene mutations in donor cells - more than an incidental phenomenon?

*Haematologica. 2020; 105:xxx
doi:10.3324/haematol.2019.242347*

Publisher's Disclaimer.

E-publishing ahead of print is increasingly important for the rapid dissemination of science. Haematologica is, therefore, E-publishing PDF files of an early version of manuscripts that have completed a regular peer review and have been accepted for publication. E-publishing of this PDF file has been approved by the authors. After having E-published Ahead of Print, manuscripts will then undergo technical and English editing, typesetting, proof correction and be presented for the authors' final approval; the final version of the manuscript will then appear in print on a regular issue of the journal. All legal disclaimers that apply to the journal also pertain to this production process.

Letter to the Editor

Donor cell leukemia: reappearance of gene mutations in donor cells – more than an incidental phenomenon?

Tal Shahar Gabay^{1,2}, Noa Chapal-Ilani³, Yoni Moskovitz³, Tamir Biezuner³, Barak Oron³, Yardena Brilon³, Anna Fridman-Dror¹, Rawan Sabah^{1,2}, Ran Balicer⁴, Amos Tanay^{3,5}, Netta Mendelson-Cohen⁵, Eldad J. Dann⁶, Riva Fineman⁶, Nathali Kaushansky³, Shlomit Yehudai-Reshef^{1*}, Tsila Zuckerman^{2,6*}, Liran I Shlush^{3,6*}

1. Hematology Research Center, Rambam Health Care Campus, Haifa, Israel
2. Bruce Rappaport Faculty of Medicine, Technion, Haifa, Israel
3. Department of Immunology, Weizmann Institute of Science, Rehovot, Israel
4. Clalit Research Institute, Tel Aviv, Israel
5. Department of Computer Science and Applied Mathematics, Weizmann Institute of Science, Rehovot, Israel
6. Department of Hematology and BMT, Rambam Health Care Campus, Haifa, Israel

*Equal contribution

Running title: Donor Cell Leukemia and Recipient Microenvironment

Conflict of Interest Statement

The authors declare no potential conflicts of interest.

Corresponding author:

Tsila Zuckerman, MD

Department of Hematology and Bone Marrow Transplantation

Rambam Health Care Campus

8, Ha'Aliya Street

Haifa 3109601, Israel

Tel: +97247772541

Fax: +97247772343

e-mail: t_zuckerman@rambam.health.gov.il

Text word count: 1227

Number of figures: 1

Number of tables: 1

Number of references: 15

Keywords: Acute myeloid leukemia, donor cell leukemia, bone marrow microenvironment, allogeneic stem cell transplantation, red blood cell distribution width, relapse

Acute myeloid leukemia (AML) is one of the extreme outcomes of age-related clonal hematopoiesis (ARCH)¹. With aging, mutations accumulate in hematopoietic stem and progenitor cells (HSPCs)^{2,3}. Based on the estimated number of HSPCs (~50,000) in the human body and the number of somatic mutations in adult single cells (~1000)⁴, it is predicted that every ~100 nucleotides, a somatic mutation will occur at a low variant allele frequency (VAF). Indeed, recent studies have demonstrated that virtually all elderly healthy individuals carry protein damaging mutations in *DNMT3a* and *TET2* at a low VAF⁵. Although mutations in preleukemic genes are inevitable, only 10-30% of older adults develop ARCH (large clone VAF>0.02)⁶. Moreover, only 1 out of 1000 carriers of somatic mutations will develop a myeloid malignancy. Recent studies have revealed that individuals with ARCH who carry larger clones and/or more than one mutation and specific mutations, are at a higher risk of AML development. However, the underlying mechanisms resulting in clonal evolution that eventually lead to full-blown AML remain unclear. HSPCs reside in the bone marrow microenvironment (BMM), where they are supported by mesenchymal stromal cells (MSCs), that maintain their quiescence and/or proliferation and differentiation. BMM can determine whether HSPCs carrying mutations will outgrow their wild type counterparts. The role of the BMM in leukemia induction has been described in a rodent model⁷; however, studies in humans are scarce.

We herein present two rare cases of late-onset donor cell leukemia (DCL) that developed in allogeneic stem cell transplant (allo-SCT) recipients and not in their sex-mismatched donors (Table 1). These findings might imply BMM involvement in the initiation and evolution of myeloid malignancies. The study was approved by the Institutional Review Board of the Rambam Health Care Campus (Approval #0016-15). The molecular profile of the patients was assessed using deep error corrected sequencing (average coverage ~5000X) (Supplementary Methods section). The presence of mutations (somatic and germline) at diagnosis and at DCL occurrence was evaluated (variants with VAF>0.1) in patients' peripheral blood (PB) and saliva as well as in donors' PB. Karyotype and chimerism analyses were performed in DCL samples. To confirm the presence of donor sex chromosomes and full donor chimerism, polymerase chain reaction (PCR) for short tandem repeats (STR) and fluorescence in situ hybridization (FISH) analyses were done (Supplementary Table 1). To explore the possibility that DCL originated from ARCH mutations present in the donor we applied ARCH variant calling algorithms aimed to accurately detect recurrent AML mutations at VAF>0.005. Additionally, we longitudinally followed complete blood count (CBC) parameters in 71 long-term AML patients post-allo-SCT. These individuals were identified in the electronic database of a health maintenance organization (HMO)

encompassing 3.45 million people. CBCs of these patients were compared to those of 500,000 age/gender-matched controls.

The current report analyzed two MDS/AML patients who underwent allo-SCT from HLA-matched, sex-mismatched family donors and relapsed with DCL years after the procedure.

In the first case, a 54-year-old male, diagnosed with AML in 2006, exhibited a mutation in the *U2AF1* S34F gene (Supplementary Tables 1 and 2). The patient achieved complete remission and was transplanted from his HLA-matched sister (52 y.o.) who carried a mutation in *DNMT3a* F752Y gene at low VAF (2.96%) at the donation time point (Supplementary Table 3). Nine years later (2015) the patient relapsed, presenting with different mutations, i.e., *U2AF1* (S34Y instead of the original S34F), *IDH1* R132C and *DNMT3a* F752Y at higher VAF (43.3%). At the time of DCL onset he had full chimerism and 100% XX donor cells, as confirmed by PCR for STR and FISH. Notably, mutations in *U2AF1* and *IDH1* genes were not detected in donor's PB either at the time of cell donation or 10 years later (Figure 1, Case 1). To date, the donor remains in good general condition with normal CBC values. In the second case, a 26-year-old female was diagnosed with high-risk MDS in 2000 and transplanted from her haplo-matched father (73 y.o.) (Supplementary Tables 1 and 2). At the time of MDS diagnosis a mutation in the *U2AF1* S34F gene was found. Seventeen years later (2017) she relapsed with high-risk MDS, presenting with mutations in *TET2* S1059X and *ASXL1* W1411X genes, not observed at diagnosis. At the DCL onset the patient had full donor chimerism and 100% XY donor cells, as confirmed by PCR for STR and FISH. These mutations were not detected in donor's PB during donation. (Figure 1, Case 2; Supplementary Table 3). Notably, the donor died from a cerebrovascular accident (CVA).

In both analyzed DCL patients, the red blood cell distribution width (RDW) remained elevated even 9 and 17 years (respectively) after the transplant (Figure 1). As increased RDW is reported to be associated with dyserythropoiesis and a poor prognosis in patients with hematological malignancies, the present study has analyzed longitudinal CBC indexes of other 71 allo-SCT recipients, recorded in the Clalit HMO dataset (Supplementary methods). While most CBC parameters, including RDW, gradually normalized in the majority of patients in this cohort over the first 4 years following allo-SCT, ~25% of these long-term survivors still exhibited high RDW. This evidence could be suggestive of underlying MDS, or other hematological pathologies associated with increased RDW⁸, DCL being one of them.

Moreover, in an Italian study including 94 adults with acute leukemia, unfavorable cytogenetics was found in 26% of patients with high RDW compared to 8% of patients with RDW within the normal range ($p=0.10$)⁸. Additionally, increased RDW was reported among individuals with ARCH⁶ and specifically those at a greater risk of AML development. This change in erythropoiesis might be mediated, at least in part, by impaired BMM, although an abnormal crosstalk between BMM cells and clonal HSPCs (ARCH) could also be a possible contributor⁹. While such damaged microenvironment might be involved both in DCL evolution and RDW dynamics, the mechanisms underlying these effects may differ.

The long period between allo-SCT and DCL development observed in our patients could imply either that the evolution occurred in donor HSPCs, originating from mutations that were scanty (below the detection limit) in the donor and then expanded in the recipient or that new mutations were acquired and the mutated cells proliferated in the recipient only. In case 1, a pre-leukemic mutation (*DNMT3a*) was found in the donor at the time of donation¹⁰⁻¹². While inherited predisposition to MDS/AML is known, it is less likely to be the case in the current patient, as no other leukemias have been reported among the family members. Hence, the DCL initiation in the reported cases could be triggered by the BMM involvement in abnormal hematopoiesis¹³⁻¹⁵. This interpretation of ours is supported by the fact that in patient #1 the same gene has been affected twice in the same position, as revealed at diagnosis and in transplanted donor's cells 9 years post-SCT. Of note, the chance of the same patient to develop AML twice in the same position within the gene (*U2AF1* S34) is less than 1:500,000².

Overall, our findings suggest that at least in some MDS/AML cases, after allo-SCT, certain mutations in the donor cells might have selective advantage in specific BMM conditions. As a result, hematopoiesis is not fully normalized due to such abnormalities of recipient's BMM which trigger leukemogenesis in the donor cells, that may eventually lead to DCL. Better understanding of this interplay could shed light on the mechanisms of AML evolution and contribute to advances in prevention and treatment of this disease.

Acknowledgements

This work was supported by the Israel Cancer Association, grants # 20170179 and 20181136.

Author Contribution:

T.S.-G. designed and performed research, wrote the paper, approved the final version of the paper

N.C.-I. performed research, interpreted the data, approved the final version of the paper

Y.M. performed research, approved the final version of the paper

T.B. performed research, approved the final version of the paper

B.A. performed research, approved the final version of the paper

Y.B. performed research, approved the final version of the paper

A.F.-D. performed research, approved the final version of the paper

R.S. performed research, approved the final version of the paper

R.B. performed research, approved the final version of the paper

O. N. performed research, approved the final version of the paper

A.T. performed research, approved the final version of the paper

N.M.-C. performed research, approved the final version of the paper

E.J.D. collected the data, approved the final version of the paper

R.V. collected the data, approved the final version of the paper

N.K. designed research, interpreted the data, approved the final version of the paper

S.Y.-R. designed research, interpreted the data, wrote the paper, approved the final version of the paper

T.Z. designed research, interpreted the data, wrote the paper, approved the final version of the paper

L.S. designed research, interpreted the data, wrote the paper, approved the final version of the paper

References

1. Shlush LI. Age-related clonal hematopoiesis. *Blood*. 2018;131(5):496-504.
2. Shlush LI, Zandi S, Mitchell A, et al. Identification of pre-leukaemic haematopoietic stem cells in acute leukaemia. *Nature*. 2014;506(7488):328-333.
3. Welch JS, Ley TJ, Link DC, et al. The origin and evolution of mutations in acute myeloid leukemia. *Cell*. 2012;150(2):264-278.
4. Lee-Six H, Obro NF, Shepherd MS, et al. Population dynamics of normal human blood inferred from somatic mutations. *Nature*. 2018;561(7724):473-478.
5. Young AL, Challen GA, Birmann BM, Druley TE. Clonal haematopoiesis harbouring AML-associated mutations is ubiquitous in healthy adults. *Nat Commun*. 2016;7:12484.
6. Jaiswal S, Fontanillas P, Flannick J, et al. Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med*. 2014;371(26):2488-2498.
7. Raaijmakers MH, Mukherjee S, Guo S, et al. Bone progenitor dysfunction induces myelodysplasia and secondary leukaemia. *Nature*. 2010;464(7290):852-857.
8. Milone G, DiMarco A, Sapienza G, et al. Red Cell Distribution Width (RDW) As Prognostic Factor in Acute Leukemia Patients Undergoing Allogeneic Hematopoietic Transplantation. *Blood*. 2017;130(Supp_1):5098.
9. Leimkuhler NB, Schneider RK. Inflammatory bone marrow microenvironment. *Hematology Am Soc Hematol Educ Program*. 2019;2019(1):294-302.
10. Hahn CN, Ross DM, Feng J, et al. A tale of two siblings: two cases of AML arising from a single pre-leukemic DNMT3A mutant clone. *Leukemia*. 2015;29(10):2101-2104.
11. Gondek LP, Zheng G, Ghiaur G, et al. Donor cell leukemia arising from clonal hematopoiesis after bone marrow transplantation. *Leukemia*. 2016;30(9):1916-1920.
12. Yasuda T, Ueno T, Fukumura K, et al. Leukemic evolution of donor-derived cells harboring IDH2 and DNMT3A mutations after allogeneic stem cell transplantation. *Leukemia*. 2014;28(2):426-428.
13. Wong TN, Miller CA, Jotte MRM, et al. Cellular stressors contribute to the expansion of hematopoietic clones of varying leukemic potential. *Nat Commun*. 2018;9(1):455.
14. Suarez-Gonzalez J, Martinez-Laperche C, Martinez N, et al. Whole-exome sequencing reveals acquisition of mutations leading to the onset of donor cell leukemia after hematopoietic transplantation: a model of leukemogenesis. *Leukemia*. 2018;32(8):1822-1826.

15. Engel N, Rovo A, Badoglio M, et al. European experience and risk factor analysis of donor cell-derived leukaemias/MDS following haematopoietic cell transplantation. *Leukemia*. 2019;33(2):508-517.

Table 1. Potential confounders responsible for the post-SCT clinical course


| | Case 1 | Case 2 |
|--|--|--|
| Conditioning regimens | Non- myeloablative: Fludarabine/Melphalan | Myeloablative: Total body irradiation/ Fludarabine/thiotepa/ ATG |
| GvHD prophylaxis | CSA + MTX | T cell depletion |
| Chronic GvHD grade | Mild | Mild-moderate |
| Acute GvHD grade | 2 | 3 |
| Post-SCT Pharmacologic immunosuppression | 1.5 years | 3 years |

GvHD= graft-versus-host disease; SCT= stem cell transplantation; CSA= cyclosporine; ATG= anti-thymocytic globulin; MTX= methotrexate


Figure legends

Figure 1. The VAF mutation profile and RDW count of two AML patients with long-term DCL
Red cell distribution width (RDW); diagnosis (Dx); variant allele frequency (VAF); donor cell leukemia (DCL).

Case 1



Case 2



SUPPLEMENTARY INFORMATION

Supplementary Methods

Human samples

Patients' bone marrow and peripheral blood samples were collected at diagnosis and relapse at the Rambam Health Care Campus, Haifa, Israel. Donors' samples were collected from peripheral blood prior to stem cell donation. Mononuclear cells were obtained from the bone marrow and peripheral blood using Ficoll separation and preserved in liquid nitrogen for further use. The use of human samples was approved by the Israel Ministry of Health (authorization No. 0023-15-RMB), in accordance with the Declaration of Helsinki. Patient clinical data are summarized in Supplementary Tables 1 & 2.

DNA purification

Mononuclear cells from the bone marrow aspiration or peripheral blood were thawed according to the standard protocol and DNA was extracted from both patients' and donors' mononuclear cells using ExgeneTM cell SV (GeneAll, cat# 106-101). DNA concentration was quantified by Qubit 3.0 fluorometer (Life Technologies, cat # Q33216). Samples that provided the DNA concentration less than 50 ng/ μ l underwent whole genome amplification using REPLI-g Mini Kit (QIAGEN, cat#150025). Saliva samples were collected from the patients and DNA was extracted using ORAgene-DNA purification Kit (DNA Genotek, cat# OG-500).

Chimerism analysis

Seventeen polymorphic short-tandem repeat (STR) markers were amplified using the AmpFISTR NGM SElect Express PCR Amplification Kit (ThermoFisher, cat # 4472193). Amplified fragments were analyzed using an ABI PRISM 3500 Genetic Analyzer (Applied Biosystems) and peak areas were quantified using Gene Mapper software (Applied Biosystems). The limit of detection of the assay was approximately 1-2%; the limit of quantitation was 5%.

Cytogenetic analysis

The karyotype was prepared from bone marrow aspirates at the metaphase stage according to standard protocols. Metaphase chromosomes were G-banded using trypsin and Giemsa and karyotypes were reported according to the International System for Human Cytogenetic Nomenclature (ISCN).

Targeted Sequencing

The Molecular Inversion Probe (MIP) Capture protocol by Hiatt et al¹ was adopted and modified. Sequencing primers and MIP backbone were as described by Hiatt et.al. We planned a MIP probe set to capture ARCH defining events. We generated the MIP panel either by microarray synthesis (LCSciences) or single oligo order (Sigma Aldrich). Probes were prepared as described by Shen et al² and Hiatt et al, respectively. Briefly: First, the probe hybridization step was performed, followed by gap-filling and ligation steps. In order to generate a final Illumina sequencing library, the final product was mixed with barcoding PCR primers, targeting the MIP backbone, and sequenced in Novaseq6000 (Illumina) 150 paired-end run.

Variant calling

To analyze the data, a dynamic, asynchronous pipeline was constructed. It started with receipt of paired-end FASTQ files and tagging each read with its UMI barcode followed by trimming the MIP's ligation and extension arms using CUTADAPT³. Trimmed FASTQ files were then aligned and mapped using BWA-MEM⁴ to a reference genome made out of broad's HG19 with only the relevant regions, each as a separate contig. Sam files were then sorted using Samtools sort⁵ and converted to the bam format. At this stage each sample went through the process termed "family collapsing", which involved reducing every family of the same MIP that held the same UMI barcodes, resulting in new FASTQs. These FASTQs were then realigned and sorted as outlined above, Tagged using Picard's AddOrReplaceReadGroups tool⁶, and processed using GATK's IndelRealigner⁷. At this point, the pipeline diverged into 2 processes for variant calling: (1) Variant calling using Varscan⁸ and (2) Variant Calling using Mutect⁹. The results of the variant calling were then processed and evaluated to only include variants that appeared in two different sequencing runs of the same sample (Supplementary Table 3). Variants were further annotated using Annovar¹⁰.

Somatic mutations were called when coverage was >1500X, Kaviar allele frequency (AF) <0.001, variant allele frequency (VAF) <0.5 and >0.005 and according to the definitions presented in Supplementary Table 4.

Clalit Database

The Clalit database includes information of patients insured by the Clalit Health Maintenance Organization (HMO) in Israel during the years 2002-2017. The Clalit dataset contains electronic medical records (EMR) of 3.45 million individuals per year on average. All data were anonymized by hashing personal identifiers and addresses followed by random sampling of data, including patient diagnoses, laboratory and medication records. This approach provided differential data analysis per patient. Diagnosis codes were acquired from both primary care and hospitalisation records, and were mapped to the ICD-9 coding system for historical reasons, with few exceptions where a partial ICD-10 coding system was used. Lab records were normalised for age and gender by subtracting raw test values from the median levels observed among all test values with matching gender and age (using a bin size of 5 years). We observed some chronological biases in lab ranges, but avoided normalising these data and instead insured patient cases and controls were matched for chronological distributions.

Defining AML cases. The EMRs were screened for all active patients ($18 < \text{age} < 100$) who were diagnosed with AML (ICD-9 code 205.0*) between the years 2003 and 2016.

The following exclusion criteria were applied:

1) We excluded patients with prior myeloid malignancies to omit secondary AML cases, which is consistent with the case selection for the genetic model. The following diagnoses were excluded if documented within 5 years prior to the AML diagnosis: essential thrombocythemia (ICD-9 238.71), low-grade myelodysplastic syndrome (MDS) (ICD-9 238.72) high-grade MDS lesions (ICD-9 238.73), MDS with 5q deletion (ICD-9 238.74), MDS, unspecified (ICD-9 238.75), polycythemia vera (ICD-9 238.4), myelofibrosis (ICD-9 289.83), chronic myelomonocytic leukemia (CMML) (ICD-9 206.10-206.22)

2) Patients who underwent any procedures performed on the bone marrow or spleen (ICD-10 code Z41) within 5 years prior to the first mentioning of the AML diagnosis code in their record. These patients were presumed to have an inaccurate AML diagnosis date or were misdiagnosed.

3) Patients who received medications suggestive of an alternative diagnosis, such as chronic myeloid leukaemia, lymphoid malignancy or acute promyelocytic leukaemia (APL):

- At any time prior to diagnosis: imatinib, dasatinib, anagrelide, hydroxycarbamide, asparaginase, pegaspargase, arsenic trioxide.
- At any time after diagnosis: imatinib, dasatinib, methotrexate, tretinoin, arsenic

trioxide.

- At any time after diagnosis, or more than single dose of mercaptopurine.

4) APL cases were excluded as they are not planned for allo-SCT.

5) Patients without a hospitalization record within 3 months prior to or 3 months after the diagnosis. This parameter was used as it is unlikely that an AML patient would not be hospitalized close to diagnosis. This filter reduced false positive cases and better defined the disease onset date.

We refined the estimated time of AML onset using the earliest time point when any of the following diagnoses appeared in patient's EMR: lymphoid leukemia (ICD-9 204), myeloid leukemia (ICD-9 205), leukemia of unspecified cell type (ICD-9 208).

A total of 875 AML cases in the training set were retained for further analysis. They were then validated by manual expert inspection of the complete records of 8 % of the cases.

To define the control set, we included all the individuals insured by Clalit that were not index cases. As for the present analysis, data were aggregated from a historical time window of 15 years, each control was associated with a randomised time point for evaluation. Using this approach, both index cases and controls represented a specific time point in the historical record of a patient, with matching calendric, age and gender distributions. Thus, 5,238,528 controls were used.

Out of the 875 AML patients we identified all those who survived more than two years from the initial diagnosis (N=174) and out of them all the ones who underwent allogeneic stem cell transplantation based on ICD-9 # 41.05 (N=71).

Laboratory features: Out of 2770 different types of lab tests, we selected the top 50 most frequent lab tests (Supplementary Table 5). For each lab measurement, we used median age/gender normalised test values and compared them to those of age/gender matched control population.

Supplementary Table 1: Clinical history of donors and recipients

| | Patient 1 | Patient 2 |
|---------------------------------|----------------------------------|--|
| Age (years) | 54 | 26 |
| Sex | M | F |
| Primary malignancy | AML | MDS |
| Cytogenetics | 46, XY | 46, XX,+8 |
| Initial chemotherapy | 7+3 | 7+3 |
| Treatment response | CR | CR |
| SCT 1 | RIC | Myeloablative haploidentical SCT |
| Source | PB | PB |
| Stem cell manipulation | NA | TCD |
| Post SCT Chimerism (STR) | 100% | 100% |
| SCT 2 | NA | Haploidentical SCT +Post SCT CTX |
| Source | | PB |
| Chimerism (STR) | NA | 100% |
| Donor- cell leukemia | | |
| Time to DCL (years) | 9 | 15 |
| Cytogenetics | 46, XX,del(6),t(6;11), (p22;q13) | 47,XY,del(20)(q11.2)+(21)/46,xy,del(20)(q11.2) |
| Chimerism (STR) | 100% | 100% |
| | Donor 1 | Donor 2 |
| Age at time of donation (years) | 52 | 73 |
| Sex | F | M |

Supplementary Table 2: CBC profile at diagnosis and relapse

| | Patient 1 | Patient 2 |
|---------------------------------|------------------|------------------|
| Diagnosis | | |
| WBC (X10 ³ /μl) | 2.03 | 2.53 |
| Hb | 10.9 | 7.5 |
| MCV | 103.7 | 96.6 |
| RDW | 17.4 | 21.3 |
| PLT (X10 ³ /μl) | 138 | 130 |
| % Blast cells in BM | 11 | 8 |
| Relapse | | |
| WBC (X10 ³ /μl) | 2.22 | 3.73 |
| Hb | 7.1 | 8.4 |
| MCV | 96.2 | 108.6 |
| RDW | 16.9 | 18.7 |
| PLT (X10 ³ /μl) | 34 | 100 |
| % Blast cells in BM | 52 | 5 |
| Donor 1 | | |
| WBC (X10 ³ /μl) | 6.17 | 6.55 |
| Hb | 12.5 | 14.6 |
| MCV | 88 | 97.6 |
| RDW | 13.2 | 14.4 |
| PLT (X10³/μl) | 173 | 206 |

Abbreviations:

- AML- acute myeloid leukemia
- BM- bone marrow
- SCT- stem cell transplantation
- CTX- cytoxan
- CR- complete remission
- DCL - donor cell leukemia
- DNR- daunorubicin
- Hb - hemoglobin
- MCV - mean corpuscular volume
- MDS - myelodysplastic syndrome
- NA - not applicable
- PB- peripheral blood
- PLT- platelets
- RDW- red cell distribution width
- RIC- reduced intensity conditioning
- SCT- stem cell transplantation
- TCD- T cell depletion
- WBC- white blood cells

Supplementary Table 3: Somatic mutations identified in recipient bone marrow (Dx) and donor peripheral blood pre-allo-SCT and at the time of DCL diagnosis

Patient 1

| Sample | Sample date | Chromosome | Position | Mutation | Type of mutation | Location | Gene | Amino Acid Change | VAF (%) |
|------------|-------------|------------|-----------|----------|------------------|----------|--------|-------------------|---------|
| Dx (BM) | 16.7.2006 | 21 | 44524456 | G>A | SNV | exonic | U2AF1 | S34F | 11.4 |
| DCL (BM) | 27.4.2014 | 21 | 44524456 | G>T | SNV | exonic | U2AF1 | S34Y | 44.86 |
| DCL (BM) | 15.1.2015 | 21 | 44524456 | G>T | SNV | exonic | U2AF1 | S34Y | 40.74 |
| Donor (PB) | 17.9.2006 | 2 | 25463238 | A>T | SNV | exonic | DNMT3A | F752Y | 2.96 |
| DCL (BM) | 27.4.2014 | 2 | 25463238 | A>T | SNV | exonic | DNMT3A | F752Y | 45.6 |
| DCL (BM) | 15.1.2015 | 2 | 25463238 | A>T | SNV | exonic | DNMT3A | F752Y | 43.3 |
| DCL (BM) | 15.1.2015 | 2 | 209113113 | G>A | SNV | exonic | IDH1 | R132C | 30 |

Patient 2

| Sample | Sample date | Chromosome | Position | Mutation | Type of mutation | Location | Gene | Amino Acid Change | VAF (%) |
|----------|-------------|------------|-----------|----------|------------------|----------|-------|-------------------|---------|
| Dx (BM) | 9.2.2000 | 21 | 44524456 | G>A | SNV | exonic | U2AF1 | S34F | 36.86 |
| DCL (PB) | 9.2.2016 | 4 | 106158275 | C>G | stopgain | exonic | TET2 | S1059X | 1.26 |
| DCL (PB) | 9.2.2016 | 20 | 31024747 | G>A | stopgain | exonic | ASXL1 | W1411X | 8.26 |
| DCL (BM) | 21.8.2017 | 20 | 31024747 | G>A | stopgain | exonic | ASXL1 | W1411X | 25.4 |

Supplementary Table 4: Mutation hotspots characterization

| Gene name | Mutation Type | Region |
|------------------|----------------------------------|--|
| ASXL1 | Frameshift/nonsense/splice-site | exon 12 |
| BCOR | Frameshift/nonsense/splice-site | whole gene |
| BCORL1 | Frameshift/nonsense/splice-site | whole gene |
| BRAF | Missense | aa range p.590-615; G469 |
| BRCC3 | Frameshift/nonsense/splice-site | whole gene |
| CALR | Frameshift | exon 9 |
| CBL | Missense | aa range p.345-434 |
| CREBBP | Frameshift/nonsense/splice-site | whole gene |
| CSF1R | Missense | L301,Y969 |
| DNMT3A | Frameshift/nonsense/splice-site | whole gene |
| DNMT3A | Missense | aa range p.292-350, p.482-614, p.634-912 |
| EZH2 | Frameshift/nonsense/splice-site | whole gene |
| EZH2 | Missense | aa range p.617-732 |
| GNAS | Missense | R201 |
| GNB1 | Missense | K57, I80 |
| IDH1 | Missense | R132 |
| IDH2 | Missense | R140, R172 |
| JAK2 | Missense/indel | V617F, aa range p.536-547 |
| KDM6A | Frameshift/nonsense/splice-site | NM_021140 |
| KRAS | Missense | G12, G13, Q61, A146 |
| NPM1 | Frameshift | exon 12 |
| NRAS | Missense | G12, G13, Q61 |
| PHF6 | Frameshift/nonsense/splice-site | whole gene |
| PPM1D | Frameshift/nonsense | exon 5, 6 |
| PRPF40B | Frameshift/nonsense/splice-site | whole gene |
| PTEN | Frameshift/nonsense/splice-site | whole gene |
| PTPN11 | Missense | aa range p.58-76, p.491-510 |
| RAD21 | Frameshift/nonsense/splice-site | whole gene |
| SF1 | Frameshift/nonsense/splice-site | whole gene |
| SF3A1 | Frameshift/nonsense/splice-site | whole gene |
| SF3B1 | Missense | aa range p.529-1201 |
| SMC1A | Missense | R96, R586 |
| SMC3 | Frameshift/nonsense/splice-site | whole gene |
| SRSF2 | Missense/deletion | P95 |
| STAG2 | Frameshift/nonsense/splice-site | whole gene |
| STAT3 | Missense | aa range p.580-670 |
| TET2 | Frameshift/nonsense/splice-site; | whole gene |
| TET2 | Missense | p.1104-1481, p.1843-2002 |
| TP53 | Frameshift/nonsense/splice-site; | whole gene |
| TP53 | Missense | aa range p.95-288, P72, R337 |
| U2AF1 | Missense | S34, R156, Q157 |

| | | |
|--------|---------------------------------|--|
| U2AF2 | Missense | aa range p.149-231, p.259-337, p.381-462 |
| ZRSR2 | Frameshift/nonsense/splice-site | whole gene |
| CEBPA | Frameshift/nonsense/splice-site | whole gene |
| RUNX1 | Frameshift/nonsense/splice-site | whole gene |
| RUNX1 | Missense | aa range p.100-440 |
| SETBP1 | missenese | 868, 870 |
| FLT3 | inframe | 580-820 |
| FLT3 | Missense | 835, 839, 841 |
| KIT | farmeshift | 416-418 |
| KIT | Missense | 816,419 |
| KMT2D | Frameshift/nonsense/splice-site | whole gene |
| NF1 | Frameshift/nonsense/splice-site | whole gene |
| WT1 | Frameshift/nonsense/splice-site | whole gene |

5. Laboratory test results included in the clinical model

| Parameter |
|--|
| Hematocrit (HCT) |
| Mean corpuscular volume (MCV) |
| Red blood cell count (RBC) |
| Hemoglobin (HGB) |
| Mean corpuscular hemoglobin (MCH) |
| Mean corpuscular hemoglobin concentration (MCHC) |
| White blood cell count (WBC) |
| Platelet count (PLT) |
| Lymphocyte percentage (LYM%) |
| Neutrophil percentage (NEUT%) |
| Eosinophil percentage (EOS %) |
| Monocyte percentage (MON%) |
| Basophil percentage (BASO %) |
| Absolute lymphocyte count (LYMP.abs) |
| Absolute neutrophil count (NEUT.abs) |
| Absolute eosinophil count (EOS.abs) |
| Absolute monocyte count (MONO.abs) |
| Basophiles (abs) |
| Mean platelet volume (MPV) |
| Red cell distribution width (RDW) |
| Creatinine - Blood |
| Glucose- Blood |
| Urea - Blood |
| Sodium |
| Potassium |
| Glutamic Oxaloacetic Transaminase |
| Glutamic Pyruvic Transaminase |
| MICR % |
| HYPO % |
| MACRO% |
| Phosphatase - Alkaline |
| Cholesterol |
| Triglycerides |
| LUC% |
| LUC |
| Cholesterol- HDL |
| Calcium - Blood |
| HYPER% |
| Uric Acid- Blood |
| Cholesterol- Ldl |
| Bilirubin Total |

| |
|------------------------------------|
| Albumin |
| Protein –Total - Blood |
| Phosphorus - Blood |
| Thyroid Stimulating Hormone (TSH) |
| Lactic Dehydrogenase (LDH) - Blood |
| Gamma Glutamyl Transpeptidase |
| Bilirubin - Direct |
| Non-HDL Cholesterol |
| PH-u |
| Specific Gravity |
| CK-CREAT. Kinase (CPK) |
| PT-INR |
| MICRO%/HYPO% |
| Vitamin B12 |
| Iron |
| PT % |
| Prothrombine time (PT- SEC) |
| Chloride (Cl) |
| Lipemic |
| Icteric |
| Hemolytic |
| Hemoglobin A1C calculated |
| CH |
| Globulin |
| Ferritin |
| T4 - free |
| APTT-sec |
| folic acid |
| PDW |
| Myeloperoxidase index (MPXI) |
| Transferrin |
| PCT |
| Cholesterol HDL ratio |
| Bilirubin indirect |
| HCT/HGB ratio |
| Creatinine Urine Sample |
| Sedimentation Rate |
| Erythrocytes |
| Leucocytes |
| C-reactive protein (CRP) |
| RDW-CV |
| M.ALBUM/CREAT ratio |
| Amylase - Blood |
| MICROALBU U SAMP |
| Protein |

| |
|--|
| Magnesium - Blood |
| Hemoglobin distribution width (HDW) |
| Fibrinogen |
| Sodium - Blood |
| Vitamin D3- 25-OH- RIA |
| Potassium - Blood |
| RDW-SD |
| Prostate specific antigen (PSA) |
| T3- free |
| Activated partial thromboplastin time (APTT-R) |
| Normoblast % |
| Estradiol (E-2) |
| Absolute normoblast count (Normoblast.abs) |
| Luteinizing hormone (LH) |

References

1. Hiatt JB, Pritchard CC, Salipante SJ, O'Roak BJ, Shendure J. Single molecule molecular inversion probes for targeted, high-accuracy detection of low-frequency variation. *Genome Res.* 2013;23(5):843-854.
2. Shen P, Wang W, Chi AK, Fan Y, Davis RW, Scharfe C. Multiplex target capture with double-stranded DNA probes. *Genome Med.* 2013;5(5):50.
3. Martin M. Cutadapt Removes Adapter Sequences from High-Throughput Sequencing Reads. *EMBnet Journal.* 2011;17:10-12.
4. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. Available at: <https://arxiv.org/pdf/1303.3997.pdf>. Accessed in 09.2019; 2013.
5. Li H, Handsaker B, Wysoker A, et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics.* 2009;25(16):2078-2079.
6. Picard Tools. Available at: <http://broadinstitute.github.io/picard>. Accessed in 09.2019.
7. McKenna A, Hanna M, Banks E, et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* 2010;20(9):1297-1303.
8. Koboldt DC, Chen K, Wylie T, et al. VarScan: variant detection in massively parallel sequencing of individual and pooled samples. *Bioinformatics.* 2009;25(17):2283-2285.
9. Cibulskis K, Lawrence MS, Carter SL, et al. Sensitive detection of somatic point mutations in impure and heterogeneous cancer samples. *Nat Biotechnol.* 2013;31(3):213-219.
10. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res.* 2010;38(16):e164.