

## KDM4C in GC lymphoma: a new piece of the epigenetic puzzle

by Karina Close and Jude Fitzgibbon

*Received: April 14, 2022.*

*Accepted: April 19, 2022.*

*Citation: Karina Close and Jude Fitzgibbon. KDM4C in GC lymphoma: a new piece of the epigenetic puzzle.*

*Haematologica. 2022 Apr 28. doi: 10.3324/haematol.2022.280898. [Epub ahead of print]*

### *Publisher's Disclaimer.*

*E-publishing ahead of print is increasingly important for the rapid dissemination of science. Haematologica is, therefore, E-publishing PDF files of an early version of manuscripts that have completed a regular peer review and have been accepted for publication. E-publishing of this PDF file has been approved by the authors. After having E-published Ahead of Print, manuscripts will then undergo technical and English editing, typesetting, proof correction and be presented for the authors' final approval; the final version of the manuscript will then appear in a regular issue of the journal. All legal disclaimers that apply to the journal also pertain to this production process.*

# ***KDM4C* in GC lymphoma: a new piece of the epigenetic puzzle**

Karina Close and Jude Fitzgibbon

Centre for Genomics and Computational Biology, Barts Cancer Institute, Queen Mary University of London, Charterhouse Sq, London EC1M 6BQ.

The group's lymphoma research is funded by Lymph & Co (Amsterdam), Blood Cancer UK (21005) and Cancer Research UK (C355/A26819) under the Accelerator Program.

In this issue of *Haematologica*, López et al.<sup>1</sup> mine whole genome sequencing (WGS) data, from germinal center (GC) lymphoma patient tumors curated as part of the International Cancer Genome Consortium (ICGC) MMML-Seq (Molecular Mechanisms in Malignant Lymphoma by Sequencing) project (n=183), to add histone demethylase *KDM4C* (4% of cases) to a burgeoning list of epigenetic regulators recurrently altered in these diseases. Both *in silico* analyses and functional reconstitution studies, in a subset of cell lines, support a tumor suppressive role of *KDM4C* in contrast to previous reports of oncogenic activity<sup>2</sup>, underlining the important contribution of cellular and molecular context to the roles of epigenetic regulators.

It is now well established that chromatin deregulation is a hallmark of cancer, seen notably in the B-cell malignancies follicular lymphoma (FL) and diffuse large B-cell lymphoma (DLBCL), where mutations arise early in the evolution of the disease, and typically affect multiple components of the chromatin machinery in the same patient. The introduction of targeted resequencing has highlighted a now familiar gamut of recurring gene mutations in *KMT2D*, *CREBBP*, *EZH2*, *ARID1A*, *EP300* and linker histones<sup>3,4</sup> and spurred efforts to understand the nature of cooperation occurring between these regulators, and their functional downstream consequences.

The findings here regarding *KDM4C* (also called *JMJD2C*), on chromosome 9p24, represents a departure, focusing on CNV (copy number variation). The gene encodes a member of the Jumonji family of demethylases, which can remove methyl groups from histone residues H3K9 and H3K36 adding to the overall complexity of the epigenetic deregulation underlying GC lymphomas (see figure) and cautions against a tunnel vision focus on well characterized marks such as H3K27 and H3K4. Indeed, the study by López et al. on *KDM4C* creates a heightened awareness of the existence of rare, and for the most part, largely inaccessible structural variants that are difficult to resolve using targeted or whole exome sequencing approaches.

Critically, it begs the questions whether *KDM4C* or the corresponding chromatin marks H3K9 and H3K36 are deregulated by other recurrent genetic or epigenetic alterations. For instance, an increase in H3K27me3 is induced not only by perturbations of the methyltransferase *EZH2* a target of SET domain gain-of-function mutations and

amplifications of chromosome 7 in GC lymphomas<sup>5,6</sup> but also by loss-of-function mutations and deletions of the corresponding demethylase *KDM6A* (also known as *UTX*) in other cancer types including leukaemias<sup>7</sup>. Indeed, loss of linker histone proteins, frequent in GC lymphomas, has been shown to result in a distinct gain of H3K36 dimethylation<sup>8</sup>. It is safe to assume that the application of WGS will uncover an increased number of diverse alterations with converging mechanistic effects to already established mutations. The translational application of epigenetic targeting therapies is exemplified by the FDA approval of EZH2 inhibitor Tazemetostat for relapsed/refractory FL. The enrichment of *EZH2* mutant patients in responders (69% *EZH2*<sup>mut</sup> vs 35% *EZH2*<sup>WT</sup>)<sup>9</sup>, represents the potential of predictive biomarkers, while also highlighting the lack of fidelity of a single genetic predictor of response, resistance to pleiotropically-acting treatments and the need to consider the modulating effects of other factors.

The contribution of epigenetic alterations to outcomes in FL is underlined by their inclusion in the clinico-genetic M7-FLIPI score, which has some predictive value of overall survival in R-CHOP-treated patients, and with five of the seven constituent genes (*EP300*, *CREBBP*, *MEF2B*, *ARID1A*, *EZH2*) representing epigenetic regulators, with either positive or negative predictive value. The holy grail, however, in an evolving treatment landscape, is a predictive model that performs independently of therapy choice, with the M7-FLIPI not informative in R-bendamustine-treated patients<sup>10</sup>. In this regard, the study by López and colleagues<sup>1</sup> makes us cogent that we need to look beyond gene mutation and be as assiduous as possible in documenting the spectrum of lesions that may be contributing to the biology of the cancer. This certainly raises challenges for companion diagnostics, but progress is being made in the ability to assess genomic sequence, translocation, and copy-number data with the aim of classifying patients into higher fidelity subtypes that offer predictive value for treatment<sup>4</sup>.

This new addition of *KDM4C* alterations to the growing list of epigenetic deregulation occurring in these diseases suggests that other rarer events have been overlooked and need to be accounted. Indeed, whether *KDM4C* plays a key role in the germinal center reaction and what contribution alterations have to malignant transformation or disease maintenance remains to be characterized. An attractive feature of leveraging epigenetic mutations for therapeutic intervention in FL is their early and clonally stable occurrence<sup>3</sup> suggesting this approach could conceivably target the common progenitor cell (CPC) population that repopulates the tumor at relapse. It would therefore be valuable to establish whether *KDM4C* alterations occur early in disease evolution.

Despite advances with agnostic treatments such as CAR-T cell and bispecific antibodies, the use of epigenetic-targeting therapies, and the profiling of epigenetic landscapes for potential biomarkers, remain at the forefront of efforts to achieve precision medicine approaches for GC lymphomas. A key challenge therefore is the elucidation of a full picture of the heterogenous drivers of the epigenetic deregulation frequent in these diseases.

## References

1. López C, Schleussner N, Bernhart S, et al. Focal structural variants revealed by whole genome sequencing disrupt the histone demethylase KDM4C in B cell lymphomas. *Haematologica*. xxx
2. Cloos PAC, Christensen J, Agger K, et al. The putative oncogene GASC1 demethylates tri- and dimethylated lysine 9 on histone H3. *Nature*. 2006;442(7100):307-311.
3. Okosun J, Bődör C, Wang J, et al. Integrated genomic analysis identifies recurrent mutations and evolution patterns driving the initiation and progression of follicular lymphoma. *Nat Genet*. 2014;46(2):176-181.
4. Wright GW, Huang DW, Phelan JD, et al. A Probabilistic Classification Tool for Genetic Subtypes of Diffuse Large B Cell Lymphoma with Therapeutic Implications. *Cancer Cell*. 2020;37(4):551-568.e14.
5. Morin RD, Johnson NA, Severson TM, et al. Somatic mutations altering EZH2 (Tyr641) in follicular and diffuse large B-cell lymphomas of germinal-center origin. *Nat Genet*. 2010;42(2):181-185.
6. Huet S, Xerri L, Tesson B, et al. EZH2 alterations in follicular lymphoma: biological and clinical correlations. *Blood Cancer J*. 2017;7(4):e555.
7. Van Haafte G, Dalglish GL, Davies H, et al. Somatic mutations of the histone H3K27 demethylase gene UTX in human cancer. *Nat Genet*. 2009;41(5):521-523.
8. Yusufova N, Kloetgen A, Teater M, et al. Histone H1 loss drives lymphoma by disrupting 3D chromatin architecture. *Nature*. 2021;589(7841):299-305.
9. Morschhauser F, Tilly H, Chaidos A, et al. Tazemetostat for patients with relapsed or refractory follicular lymphoma: an open-label, single-arm, multicentre, phase 2 trial. *Lancet Oncol*. 2020;21(11):1433-1442.
10. Pastore A, Jurinovic V, Kridel R, et al. Integration of gene mutations in risk prognostication for patients receiving first-line immunochemotherapy for follicular lymphoma: a retrospective analysis of a prospective clinical trial and validation in a population-based registry. *Lancet Oncol*. 2015;16(9):1111-1122.

## Figure Legend

Epigenetic deregulation in germinal center lymphomas: a growing number of histone proteins and their tail residues altered in these diseases.

