

Vaccine utilization and overwhelming post-splenectomy infection risk factors in two asplenia cohorts

Matthew A. Soderstrom,¹ Mechelle A. Miller M.D.,² Qing Wang,² William P. Hennrikus,³ Nora L. Watson,⁴ Ryan C. Costantino,⁵ Matthew J. Bradley,³ V. Koneti Rao⁶ and Nathan A. Boggs^{2,7}

¹Department of Internal Medicine, Brooke Army Medical Center, San Antonio, TX; ²Allergy & Immunology Service, Walter Reed National Military Medical Center, Bethesda, MD; ³Department of General Surgery, Walter Reed National Military Medical Center, Bethesda, MD; ⁴Department of Research Programs, Walter Reed National Military Medical Center, Bethesda, MD; ⁵Enterprise Intelligence and Data Solutions Program Office, Program Executive Office, Defense Healthcare Management Systems, San Antonio, TX; ⁶Laboratory of Clinical Immunology and Microbiology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD

and ⁷Department of Medicine, Uniformed Services University, Bethesda, MD, USA

Correspondence:

N. A. BOGGS - nathan.boggs@usuhs.edu

<https://doi.org/10.3324/haematol.2023.283419>

Received: May 3, 2023.

Accepted: July 12, 2023.

Early view: July 20, 2023.

©2024 Ferrata Storti Foundation

Published under a CC BY-NC license 