



THE VENO-OCCLUSIVE DISEASE OF THE LIVER

FRÉDÉRIC BARON, MANUEL DEPRez, YVES BEGUIN

Department of Medicine, Division of Hematology, and Department of Pathology; University of Liège, Liège, Belgium.

ABSTRACT

Background and Objective. The veno-occlusive disease of the liver (VOD) is a disorder caused by the non-thrombotic occlusion of the central veins of hepatic lobules. The clinical features are similar to those of intrahepatic portal hypertension (unexplained weight gain, ascites, painful hepatomegaly, jaundice). In the past, this disease was rather infrequent and was linked to the absorption of toxic agents, liver irradiation or chemotherapy. However, the intensification of treatment protocols before hematopoietic stem cell transplants has considerably increased its incidence. The strategies used for its prevention and treatment remain limited in efficacy. The present review was undertaken in order to assess progress in the diagnosis and management of this severe complication in stem cell transplantation.

Information Sources. The method used for preparing this review was an examination of 250 relevant articles or abstracts published in journals covered by Medline®.

State of Art. Despite the progress made toward the

understanding of its physiopathology and the identification of its risk factors, VOD is still one of the leading causes of morbidity and mortality during the first two months post-BMT, and therefore often constitutes a limitation for the further increment of the dose of antineoplastic drugs. This may be explained by the difficulty in making an early diagnosis of this problem, at a time when therapeutic intervention may be more effective, and, on the other hand, the lack of a well-established prevention and treatment approach for patients with VOD.

Perspectives and Conclusions. New diagnostic procedures, such as laparoscopic liver biopsy, and new therapeutic approaches, such as transjugular intrahepatic portosystemic shunting (TIPS) or defibrotide, are now being evaluated. However, additional studies will be needed to determine the most appropriate therapy for each VOD patient depending on the severity of the disease.

©1997, Ferrata Storti Foundation

Key words: stem cell transplantation, veno-occlusive disease, liver complications, anticoagulation

The veno-occlusive disease of the liver (VOD) results from a hepatotoxic lesion causing the obstruction of small intrahepatic venules, hence leading to lesions in the central zone hepatocytes and surrounding sinusoids.¹ Although the disease was first described in some Jamaicans who habitually consumed local herb infusions containing pyrrolizidine alkaloids (senecio),² nowadays VOD is above all a serious complication of bone marrow transplants (BMT). Its incidence varies considerably from 3% in pediatric centers that perform allogeneic transplants in children with thalassemia major,³ to more than 65% in centers that carry out bone marrow transplants in cases of advanced hematological disorders.⁴ However, the incidence of VOD varies among these centers, depending on their capacity to diagnose early and mild VOD. The Seattle⁵⁻⁷ and Baltimore⁸ groups have set up similar clinical diagnostic criteria for VOD (Table 1). Patients who meet modified Seattle criteria^{6,7} are retrospectively classified for severity in

three groups (mild, moderate and severe VOD) based on their outcome (Table 2). In a cohort study of 355 patients,⁹ mild, moderate and severe VOD occurred respectively in 12%, 26% and 15% of the patients, and their mortality rates before day 100 were 9%, 23% and 98%, respectively. In this article, we will review the pathophysiology and various risk factors of this disease, describe its clinical, biological and anatomical aspects, and present options for its prevention and treatment.

Pathophysiology

Although the exact sequence of events leading to the clinical manifestations of VOD is still debated, the earliest event seems to be damage to the hepatic venular and sinusoidal endothelium, which locally induces a hypercoagulable state by activating the coagulation cascade and favoring clot formation over natural anticoagulation.^{1,7,10-13} Consequently, the venular and sinusoidal lumen is reduced due to an edematous concentric subendothelial zone con-

Correspondence: Yves Beguin, MD, University of Liège, Department of Hematology, CHU Sart-Tilman, 4000 Liège, Belgium. Tel. international +32.4.3667690. Fax international +32.4.3668855.

Acknowledgements: Frédéric Baron is Research Assistant and Yves Beguin Senior Research Associate of the National Fund for Scientific Research (FNRS, Belgium) Received May 26, 1997; accepted September 22, 1997.

Table 1. Clinical criteria for the diagnosis of VOD.

1) *Seattle criteria* (see ref. #5)
Presence, before day 30 after transplant, of at least two of the following features:

1. jaundice
2. hepatomegaly and right upper quadrant pain
3. ascites and/or unexplained weight gain

2) *Baltimore criteria* (see ref. #8)
Hyperbilirubinemia ≥ 2 mg/dL (34.2 μ mol/L) before day 21 after transplant and, at least, two of the following features:

1. hepatomegaly (usually painful)
2. ascites
3. weight gain greater than 5% from baseline

3) *Modified Seattle criteria* (see refs. #6, 7)
Occurrence of two of the following events within 20 days of transplantation:

1. hyperbilirubinemia (total serum bilirubin > 2 mg/dL)
2. hepatomegaly or right upper quadrant pain of liver origin
3. unexplained weight gain (> 2% of baseline body weight) because of fluid accumulation.

Table 2. Classification of VOD according to its severity.⁹

1) *Mild VOD*
Patients have mild VOD if:

1. they show no adverse effect from liver disease
2. they require no treatment for VOD
3. their illness is self limited

2) *Moderate VOD*
Patients have moderate VOD if:

1. they have an adverse effect from liver disease
2. they require treatment for VOD (such as diuretics for fluid retention or medication to relieve pain from hepatomegaly)

2) *Severe VOD*
Patients have severe VOD if:

1. their VOD does not resolve before day 100
2. they die of VOD

taining fragmented red cells, debris and fibrillar material, thus inducing partial to complete fibrotic obliteration of the venular lamina.⁷ Systemic abnormalities of coagulation proteins have been observed, either before conditioning in patients who will later develop VOD or after high-dose chemo- or radiotherapy.^{13,14} The main issue regarding these data is discovering whether these coagulation parameters are involved in the pathogenesis of VOD or if they are merely consequences of the disease. Several cytokines, such as TNF- α , IL-1 or TGF- β have been implicated on the basis of their known

effects on endothelial cells and because of their elevated levels in patients before the onset of hepatic dysfunction associated with VOD.^{15,16} Although endothelial cells appear to be the primary target of events leading to VOD, hepatocytes also play a critical role. Centrilobular (zone 3) hepatocytes contain cytochrome P 450 enzymes that metabolize many of the chemotherapeutic agents used in conditioning regimens for BMT. Some drugs are transformed into toxic metabolites and then converted into stable metabolites by glutathione and glutathione-dependent enzymes that are present in only low concentration in centrilobular hepatocytes. Thus, when a drug that potentially gives rise to toxic metabolites is administered together with agents that reduce glutathione, there is a high probability of centrilobular hepatocyte injury and sinusoidal damage.^{7,17}

Risk factors for VOD

The occurrence of VOD is linked to a number of factors among which we can distinguish pre-transplant risk factors, risk factors associated with the conditioning regimen, and risk factors linked to the type of transplant.

Pre-transplant risk factors

The following risk factors are currently considered as being significantly associated with VOD: elevated transaminase levels before the beginning of the conditioning regimen;⁹ treatment for viral or bacterial infections at the beginning of the conditioning regimen;⁹ liver metastases¹⁸ or previous liver irradiation.⁹ Prior cumulative exposure to high doses of cytotoxic agents may contribute to these risks. Thus, a second marrow transplant preceded by a second conditioning treatment is often complicated by VOD.⁹ The contribution of the hepatitis C virus (HCV) to VOD is controversial. One study¹⁹ suggested that patients with liver disease caused by HCV infection are at a high risk of developing lethal VOD after BMT, but another report²⁰ in a selected and homogeneous population did not confirm the association.

Risk factors related to the conditioning regimen

Nearly all preparative regimens before a hematopoietic stem cell transplant can cause VOD, particularly the most intensive ones.^{9,11} Nevertheless, some conditioning protocols seem to be more toxic than others. For instance, the association of cyclophosphamide and busulfan has been shown to cause VOD more than the association of cyclophosphamide and total body irradiation.²¹ However, some recent reports do not show such an association.^{22,23} The technique of total body irradiation administered may also be important since fractionating the dose could significantly reduce the risk of VOD²⁴ in one study, but not in another.²⁵ The total

dose of radiation received by the patient is also correlated with VOD.⁹ The liver toxicity of busulfan clearly appears to be related to its pharmacokinetics: Grochow *et al.* originally correlated VOD with increased mean busulfan area under the curve of concentration vs. time (AUC) (>2012 $\mu\text{Mol}/\text{min}/\text{L}$).²⁶ A later prospective study by the same group²⁷ confirmed an increased occurrence of VOD with AUC > 1500 $\mu\text{mol}/\text{min}/\text{L}$, and showed that adaptation of the dose according to the AUC significantly reduced the risk of VOD. However, one study²⁸ was unable to correlate the risk of VOD with Busulfan pharmacokinetic parameters, and two additional papers^{29,30} reported that the adjustment of the dose according to the AUC failed to reduce VOD frequency.

Type of transplant

Contrary to what is commonly believed, multivariate analyses have shown that the incidence of VOD after allogeneic BMT is not higher than its incidence after autologous BMT.⁹ Nevertheless, the relative risk of severe VOD was found to be 2.15 times higher in mismatched family or unrelated allogeneic recipients as compared to HLA-identical sibling transplants.⁶ The use of methotrexate as prophylaxis of GVHD is associated with an increased risk of VOD as compared to the use of cyclosporine alone or together with corticosteroids.⁴

Clinical aspects

Clinical history

Usually, an unexplained weight gain is the first symptom of VOD. This weight gain, attributable to water and sodium retention by the kidney, appears within 6 to 8 days following the transplant in 95% of patients that develop VOD.¹² Two to three days later, hyperbilirubinemia of varying degrees is observed in 98% of the cases.¹² An increase in aspartate aminotransferase and alkaline phosphatase may occur simultaneously or a few days later. Most patients develop ascites and pain in the upper right quadrant, and clinical examination usually reveals a firm and painful hepatomegaly. Many patients become refractory to platelet transfusions.^{7,11-12} Renal insufficiency is also present in 50% of patients developing VOD (mainly patients with severe VOD) and 25% of them will require hemodialysis.^{12,31} Finally, patients with severe VOD can display severe encephalopathy or even become comatose.

Outcome

In 50 to 80% of the cases, a gradual improvement of the clinical condition is noted at 2 to 4 weeks after the onset of the disease.^{6,12} In the remaining 20 to 50% of the cases, patients die because of or with severe VOD. The majority of them die of multiple organ failure (MOF) with hepatic failure, but also

of renal failure, congestive heart failure, extravascular fluid effusions, and pulmonary failure requiring oxygen support and mechanical ventilation.⁶ The reasons for lung failure may include pulmonary VOD, pleural effusions and interstitial pneumonitis of infectious or non-infectious origin.^{31,32} Bleeding at gastrointestinal sites or at other sites is a significant cause of death as well. The Seattle group³³ has developed an elaborate regression model that estimates the probability of developing severe (mortal) VOD based on early measurements of serum bilirubin and the percent of weight gain. However, the sensitivity of these models is much lower than their specificity.

Diagnostic criteria

Clinical picture

Although theoretically the diagnosis of VOD should be based on the histological examination of the liver, the risk of bleeding during a liver biopsy in thrombopenic patients who are often refractory to platelet transfusions is such that the diagnosis is very often only a clinical one. The Seattle⁵ and Baltimore⁸ groups have set up similar diagnostic criteria for VOD (Table 1). Although the diagnostic accuracy of these protocols is excellent when all criteria are met, other groups have found that their sensitivity and specificity diminish in the early diagnosis of VOD when only 2 criteria are present.³⁴ However, in this case, a liver biopsy is not very sensitive, partly because of the heterogeneity of VOD lesions throughout the liver. It is not yet clear whether the modification of the Seattle criteria (Table 1) will improve their sensitivity and specificity. Patients who meet modified Seattle criteria for VOD but have another cause of liver dysfunction (Table 3) or patients who meet only one of the modified Seattle criteria all have liver dysfunction of uncertain causes. In McDonald's study of 355

Table 3. Differential diagnosis of VOD.

1) After allogeneic BMT:
1. Acute liver GVHD
2. Cyclosporine-induced hepatotoxicity
2) After autologous or allogeneic BMT:
1. Fungal infiltration of the liver
2. Viral hepatitis
3. Cholangitis lenta, e.g. during sepsis
4. Drug (trimethoprim-sulfamethoxazole, some third-generation penicillins, fluconazole and itraconazole) induced liver dysfunction
5. Constrictive pericarditis and right congestive heart failure
6. Persistent tumor infiltration of the liver

patients,⁹ 72 patients (22%) were classified as having liver disease of an uncertain cause. The reasons for this classification are shown in Table 4.

Differential diagnosis

Other liver diseases are common after BMT (Table 3). However, the presence of weight gain and fluid retention is usually sufficient to differentiate VOD from other causes of early liver dysfunction.

Acute liver GVHD causes jaundice with increased serum alkaline phosphatase and aminotransferase levels. Ascites, liver failure and encephalopathy are unusual. AGVHD usually develops between days 20 and 40 post-transplant together with skin and/or gut GVHD.³⁵ However, the differential diagnosis with VOD can be difficult when GVHD develops earlier (hyperacute GVHD) or when the onset of VOD is later than day 10. Moreover, both diseases are common after allogeneic BMT and may thus coexist. In this case, measurement of the hepatic venous pressure and/or a histologic evaluation of the liver may be helpful in determining which disease is dominant.

Fungal infiltration of the liver usually causes tender hepatomegaly, fever, and markedly elevated serum alkaline phosphatase levels³⁶ (which are unusual with VOD). However, fungi (mainly *Candida* species) can also invade blood vessels, causing hepatic infarctions or venous obstructions that produce tender hepatomegaly, ascites and signs of portal hypertension mimicking VOD.³⁷ Viral infections of the liver are unusual in the early post-transplant period because of the systematic use of acyclovir and because B and C hepatitis viruses can produce liver injury only in the presence of an intact immune system.¹¹ Cholangitis lenta follows sepsis and other causes of cytokine release. The bilirubin level can exceed 10 mg/dL, but ascites, weight gain and renal failure are unusual.

Medications used in the transplant setting can also induce liver dysfunction. Cyclosporine, in a dose-dependent fashion,³⁵ may cause cholestasis and hepatocyte necrosis and lead to gallstones.

Table 4. Reasons for assigning patients to the group of liver disease of «uncertain cause» in the McDonald's study (ref. #9).

Signs of VOD not meeting modified Seattle criteria: 35%

Other liver disease within 20 days of bone marrow infusion:

- sepsis: 28%
- GVHD: 22%
- cardiac failure: 8%
- persisting tumor infiltration of the liver: 4%

Died before liver disease could become clinically apparent: 4%.

Total parenteral nutrition, some antimicrobial drugs (such as trimethoprim-sulfamethoxazole, some third-generation penicillins, fluconazole and itraconazole) and methotrexate can produce both cholestasis as well as hepatocellular injury as well.

Constrictive pericarditis and right congestive heart failure can cause pain in the upper right quadrant, hepatomegaly, ascites, peripheral edema and pleural effusions, as well as renal failure and an increase in the concentration of serum hepatic enzymes. Both diseases can be ruled out by normal echocardiography.

Finally, in patients with refractory disease, persistent tumor infiltration of the liver should be considered.

Histology

When the diagnosis cannot be made clinically (because the timing of symptoms is modified by therapies affecting the natural evolution of the disease, or only two out of the 3 Seattle criteria are present, or clinical data suggest another cause of hepatic disorder), and aggressive therapy is required, a transjugular liver biopsy can be carried out.³⁸ This further allows for the measurement of the hepatic venous pressure gradients (HVPG), which favor VOD if greater than 10 mm Hg,^{38,39} and shows a high prognostic value. However, because of the risk of bleeding which may be fatal or preclude beneficial therapy, and because of low sensitivity in patients not meeting clinical criteria, the role of transjugular liver biopsy is still debated. It should be noted that the risk of complications after measurement of HVPG alone (without liver biopsy) is low, and this may be useful in differentiating VOD from GVHD. Another interesting approach is the laparoscopic liver biopsy.⁴⁰ This technique was recently evaluated in a group of 29 patients, including 24 BMT recipients, with hepatic dysfunction after chemotherapy. Thirty-two biopsies were obtained and all were informative. Furthermore, no procedure-related complication was noted and no patient required re-exploration.

From the histologic standpoint (Figure 1), the most distinctive feature of VOD is a thickening of the subintimal zone of central and sublobular venules, producing concentric or eccentric luminal narrowing.⁴¹⁻⁴³ In the early stages, the changes consist of a marked widening of the subendothelial zone by fragmented red cells within an edematous proliferation where fibrinogen can be identified by a special stain (Mallory's stain, Lendrum's MSB technique) or immunochemistry.⁴⁴ Thrombosis does not occur and inflammatory cells are few or absent. These obliterative lesions are focally distributed and can be obscured by the background features of acute venous outflow obstruction; they show pronounced centrilobular congestion, sinusoidal dilatation and hepatocyte loss. Therefore,




Figure 1. Masson trichrome X 250 showing early stages of VOD. Concentric luminal narrowing of centrilobular venules by subendothelial edema and accumulation of fragmented red cells, fibrinogen and hemosiderin-laden macrophages with background features of acute venous outflow obstruction.




Figure 2. Masson trichrome X 250 showing late stages of VOD. Non-specific centrilobular fibrous scarring showing an ingrowth of small vascular channels.

these early lesions can be missing or inconspicuous on routine stains, and the changes can be difficult to distinguish from sinusoidal congestion. Connective tissue stains (trichrome techniques) may help their identification.

In later stages, the subintimal lesion becomes fibrotic and can acquire an ingrowth of small vascular channels (Figure 2). Eventually the affected venules are incorporated into the centrilobular fibrous scarring and can only be identified by connective tissue stains. Change in chronic venous outflow obstruction dominates the histological picture: perivenular fibrosis, pericellular fibrosis and central-central fibrous bridges develop and cirrhosis may ultimately appear. These late changes are non-specific and can be seen in other types of chronic outflow obstruction and cirrhosis as well. Hepatocyte hyperplasia of varying degrees and peliosis hepatis have occasionally been found to accompany VOD and are particularly prominent in

azathioprine-treated renal transplant patients.⁴²

Connective tissue usually consists of a cellular portion in an enveloping framework of non-cellular substances, either of fibrous (collagen, reticular and elastic fibers), amorphous or gel types. A detailed review of the biochemical principles for staining of these substances is beyond the scope of this article.⁴⁵ Many techniques are available for the differential analysis of the connective tissues, the most frequently used being the Van Gieson and Masson trichrome techniques (Figures 1 and 2). Trichrome stains, which employ three dyes, allow for the selective demonstration of muscle, collagen fibers, fibrin and erythrocytes. Three methods selectively demonstrate fibrin: Gram-Weigert, Mallory's phosphotungstic acid, hematoxylin and Masson's trichrome modified after Lendrum. Demonstration of elastic fibers is best achieved by the following techniques: Verhoeff, orcein, Weigert's resorcin fuchsin and aldehyde fuchsin. Reticular fibers are best demonstrated by techniques that use silver salts, such as Gordon and Sweet's method or Gomori's method.

Other investigations

Other than the clinical picture and the histology, ultrasound studies may help in diagnosis by showing ascites, hepatomegaly or thickening of the gallbladder wall, which are, in any case, all non-specific or unreliable findings. Pulsed Doppler ultrasound may show a decreased or inverted portal blood flow but this is a relatively late finding in patients with established VOD.⁴⁶ A prospective study from Seattle⁴⁶ was not able to identify any sonographic feature strongly associated with VOD in the early phases of the disease, when a definitive diagnosis is most needed.

Several laboratory parameters have been studied as potential markers of VOD. Serum procollagen III⁴⁷ or its N-terminal peptide⁴⁸ have been found to be early markers of VOD even before any clinical sign of the disease. Contrary to protein S and antithrombin III, low protein C levels can discriminate between patients with or without VOD, but this is a predictive test rather than a specific marker, because the difference between the two groups is already evident before conditioning.¹³ Finally, a recent study⁴⁹ has shown that the level of plasminogen activator inhibitor 1 (PAI-1) was significantly elevated at the time of bilirubin increase in patients with VOD as compared with patients with GVHD or other causes of hepatic damage.

Prevention

Given the very high mortality rate in patients with severe VOD, it is critical to prepare effective preventive strategies during hematopoietic stem cell transplants. Preventive measures have essentially been based on four different molecules: heparin,

prostaglandin E1, pentoxifylline and ursodeoxycholic acid.

The usefulness of low-dose heparin has been shown in a prospective randomized study conducted by Attal,⁵⁰ who compared a control group and a group receiving heparin at a dose of 100 U/kg/d from day -8 until day +30 after autologous and allogeneic BMT. The incidence of VOD decreased from 13.7% in the control group to 2.5% in the heparin group. However, there was no difference in the incidence of severe VOD which was very low in both arms. In a double-blind, placebo-controlled study in 61 patients undergoing autologous or allogeneic BMT, the low molecular weight heparin enoxaparin was shown to significantly reduce the incidence and duration of VOD, but severe VOD was not reported in the placebo group.⁵¹ In addition, platelet engraftment was accelerated, platelet transfusion was reduced, and hemorrhagic events were less frequent and less severe in the treated group. Thus, although the administration of low-dose heparin appears to be safe, it remains to be shown in a large randomized trial that severe VOD can be prevented in high-risk patients.

In 1989, Gluckman⁵² reported a study in which prostaglandin E1 (PGE1) was administered at a dose of 3 µg/kg/h from the start of the preparative regimen until day 30 post-transplant in leukemic patients undergoing allogeneic BMT. The incidence of VOD decreased from 39.1% in the historical control group to 12.8% in the treated group, most notably in patients with previous hepatitis. However, others⁵³ were not able to reproduce these results in patients at high risk of VOD, and noted considerable toxicity precluding full completion of the treatment.

Pentoxifylline (PTX) is a methylxanthine analogue which can inhibit transcription of TNF-α. In a non-randomized study, the administration of PTX considerably reduced the incidence of VOD and other toxicities compared to historical controls.¹⁶ However, randomized prospective studies have failed to show any advantage of PTX in the prevention of VOD or other toxicities associated with BMT.⁵⁴

Ursodeoxycholic acid, an artificial bile acid, protects hepatocytes from damage caused by cholestasis. Essell⁵⁵ reported a low incidence of VOD in patients receiving ursodeoxycholic acid prophylactically (which was fairly well tolerated). More recently, a prospective study by the same group⁵⁶ showed that patients randomized for ursodeoxycholic acid developed significantly less VOD than patients receiving the placebo.

Finally, when high-dose radiotherapy (>12 Gy) is applied or when VOD occurs prior to completion of TBI, liver shielding may be used to reduce hepatic toxicity, although its efficacy is not demonstrated and its effect on tumor eradication remains unknown.

Treatment

While more knowledge about the pathophysiology of VOD has allowed for the proposal of such prevention, it also lays the ground for therapeutic measures, including supportive care, PGE1, r-tPA, and liver transplantation.

Supportive care is aimed at maintaining intravascular volume and renal perfusion while avoiding extravascular fluid accumulation.¹¹ This involves restriction of the sodium supply and diuretic therapy with loop diuretics or spironolactone. Perfusion of albumin concentrates or colloids can help maintain the intravascular volume, but will eventually accumulate in extravascular spaces. Therefore, preference should be given to transfusion of red cells to maintain the hematocrit above 38-40%. Several groups have advocated continuous perfusion of dopamine to maintain renal blood flow, but its real value has not been proven. Finally, when fluid accumulation cannot be controlled, hemodialysis is necessary. When ascites limit breathing, paracentesis may be required. Encephalopathy should be treated with protein restriction and oral lactulose. The use of peritoneovenous shunts can improve symptomatology, but runs a very high risk of serious complications.

There are also approaches aimed at reversing venular occlusion. There are only brief reports of PGE1 improving established moderate VOD.¹¹ Recombinant tissue plasminogen activator (r-tPA) has been used in several centers. The largest experience was reported by the Seattle⁵⁷ group in 42 patients receiving 5 to 120 mg of r-tPA over 2-4 days together with low-dose heparin. Twelve (29%) patients responded, but with a substantial risk of serious bleeding. Non-responders included those requiring supplemental oxygen, mechanical ventilation or dialysis before the start of r-tPA.

Defibrotide, a novel polydeoxyribonucleotide, has several properties of potential interest for the treatment of VOD, such as the increase of endogenous tissue plasminogen activator and the decrease of plasminogen activator inhibitor type 1. Richardson *et al.*⁵⁸ treated eight patients with severe VOD using defibrotide. Complete responses were achieved in three patients, none of whom had responded to r-tPA.

Several groups have used portosystemic shunting in order to decrease portal hypertension. Surgical shunting is possible,⁵⁹ but because few patients with severe VOD are reasonable candidates for surgery, the most interesting technique appears to be transjugular intrahepatic portosystemic stent-shunt (TIPS).⁶⁰⁻⁶² This technique consists in creating an intraparenchymal channel (kept permeable with a metal stent) between a main branch of the portal vein and a hepatic vein using a percutaneously inserted transjugular catheter. Although TIPS improved clinical status and liver function in a few

patients, more extensive experience should be accumulated on the risks and efficacy of the procedure.

Finally, as a last resort in case of life-threatening hepatic failure, orthotopic liver transplantation can be considered.^{63,64} However, very few patients who have received a liver transplant for acute VOD have become long-term survivors.¹¹

Perspectives and conclusions

VOD is a serious complication of high-dose regimens used as conditioning before BMT, for which a number of risk factors have been identified. We favor the systematic use of low-dose heparin (100 U/kg/d) or low molecular weight heparin, starting with the conditioning regimen and continuing through day 21 post-transplant, in all patients undergoing autologous or allogeneic BMT. For patients particularly at risk for developing severe VOD, we add ursodeoxycholic acid during the same period. It is of utmost importance to identify signs of VOD as early as possible, so that supportive care can be adapted accordingly. The differential diagnosis with acute GVHD or liver infections is critical because these complications may coexist, particularly in allogeneic transplant recipients, and their management is radically different and not devoid of serious side effects. In this setting, a transjugular measurement of HVPG and even a liver biopsy may be needed when clinical diagnostic criteria such as those proposed by the Seattle and Baltimore groups are not met or when concomitant disorders are suspected. Aggressive therapeutic intervention, such as r-tPA 20-40 mg continuous infusion over 3-4 days, should be initiated early (in order to avoid multiorgan failure), but should be performed only in those patients who could benefit from this in the face of rapidly progressive VOD. Insertion of TIPS by an experienced hand may be a valid option in patients with major counter-indications or no response to r-tPA, whereas the role of defibrotide remains to be determined.

References

- Shulman HM, McDonald GB, Matthews DE, et al. An analysis of hepatic venoocclusive disease and centrilobular hepatic degeneration following bone marrow transplantation. *Gastroenterology* 1980; 79:1178-91.
- Wilmot FC, Robertson GW. Senecio disease, or cirrhosis of the liver due to Senecio poisoning. *Lancet* 1920; 2:848-9.
- Lucarelli G, Galimbert M, Delfini C, et al. Marrow transplantation for thalassaemia following busulfan and cyclophosphamide. *Lancet* 1985; i:1355-7.
- Essel JH, Thompson JM, Harman GS, et al. Marked increase in veno-occlusive disease of the liver associated with methotrexate use for graft-versus-host disease prophylaxis in patients receiving busulfan/cyclophosphamide. *Blood* 1992; 79:2784-8.
- McDonald GB, Sharma P, Matthews DE, Shulman HM, Thomas ED. Venocclusive disease of the liver after bone marrow transplantation: diagnosis, incidence, and predisposing factors. *Hepatology* 1984; 4:116-22.
- McDonald GB, Hinds M, Fisher L, Schoch HG. Liver toxicity following cytoreductive therapy for marrow transplantation: risk factors, incidence, and outcome [abstract]. *Hepatology* 1991; 14:162A.
- Shulman HM, Hinterberger W. Hepatic veno-occlusive disease - liver toxicity syndrome after bone marrow transplantation. *Bone Marrow Transpl* 1992; 10:197-214.
- Jones RJ, Lee KSK, Beschoner WE, et al. Venocclusive disease of the liver following bone marrow transplantation. *Transplantation* 1987; 44:778-83.
- McDonald GB, Hinds MS, Fisher LD, et al. Venocclusive disease of the liver and multiorgan failure after bone marrow transplantation: A cohort study of 355 patients. *Ann Intern Med* 1993; 118:255-67.
- McDonald GB, Shulman HM, Wolford JL, Spencer GD. Liver disease after human marrow transplantation. *Semin Liver Dis* 1987; 7:210-29.
- Bearman SI. The syndrome of hepatic veno-occlusive disease after marrow transplantation. *Blood* 1995; 85:3005-20.
- Carreras E, Grañena A, Rozman C. Hepatic veno-occlusive disease after bone marrow transplant. *Blood Rev* 1993; 7:43-51.
- Faioni EM, Krachmalnicoff A, Bearman SI, et al. Naturally occurring anticoagulants and bone marrow transplantation: plasma protein C predicts the development of venocclusive disease of the liver. *Blood* 1993; 81:3458-62.
- Scrobohachi MI, Drouet L, Monemmasi A, et al. Liver veno-occlusive disease after bone marrow transplantation: changes in coagulation parameters and endothelial markers. *Thromb Res* 1991; 63:509-19.
- Holler E, Kolb HJ, Möller A, et al. Increased serum levels of tumor necrosis factor alpha precede major complications of bone marrow transplantation. *Blood* 1990; 75:1011-6.
- Bianco JA, Appelbaum FR, Nemunaitis J, et al. Phase I-II trial of pentoxifylline for the prevention of transplant-related toxicities following bone marrow transplantation. *Blood* 1991; 78:1205-11.
- Rappaport AM, Knoblauch M, Zelin S, Summerall J, Bras G. Experimental hepatic veno-occlusive disease. In vivo microcirculatory study. *Adv Microcirc* 1969; 2:69-79.
- Ayash LJ, Hunt M, Antman K, et al. Hepatic veno-occlusive disease in autologous bone marrow transplantation for solid tumors and lymphomas. *J Clin Oncol* 1990; 8:1699-706.
- Frickhofen N, Wiesneth M, Jainta C, et al. Hepatitis C virus infection is a risk factor for liver failure from veno-occlusive disease after bone marrow transplantation. *Blood* 1994; 83:1998-2004.
- Erer B, Angelucci E, Lucarelli G, et al. Hepatitis C virus infection in thalassaemia patients undergoing allogeneic bone marrow transplantation. *Bone Marrow Transpl* 1994; 14:369-72.
- Ringden O, Ruutu T, Remberger M, et al. A randomised trial comparing busulfan with total body irradiation as conditioning in allogeneic marrow transplant recipients with leukemia: a report from the Nordic Bone Marrow Transplantation group. *Blood* 1994; 83:2723-30.
- Devergie A, Blaise D, Attal M, et al. Allogeneic bone marrow transplantation for chronic myeloid leukemia in first chronic phase: a randomized trial of busulfan-cytosine versus cytosine-total body irradiation as preparative regimen: a report from the French society of bone marrow graft (SFGM). *Blood* 1995; 85:2263-8.
- O'Donnell MR, Long GD, Parker PM, et al. Busulfan/cyclophosphamide as conditioning regimen for allogeneic bone marrow transplantation for myelodysplasia. *J Clin Oncol* 1995; 13:2973-9.
- Deeg HJ, Sullivan KM, Buckner CD, et al. Marrow transplantation for nonlymphoblastic leukemia in first remission: toxicity and long-term follow-up of patients conditioned with single dose or fractionated total body irradiation. *Bone Marrow Transpl* 1986; 1:151-7.
- Valls A, Granena A, Carreras E, Ferrer E, Algara M. Total body irradiation in bone marrow transplantation: fractionated vs single dose. Acute toxicity and preliminary results. *Bull Cancer* 1989; 76:797-804.
- Grochow LB, Jones RJ, Brundrett RB, et al. Pharmacokinetics of busulfan: correlation with veno-occlusive disease in patients undergoing bone marrow transplantation. *Cancer Chemother Pharmacol* 1989; 25:55-61.
- Grochow LB. Busulfan disposition: the role of therapeutic monitoring in bone marrow transplantation induction regimens. *Semin Oncol* 1993; 20(Suppl 4):18-25.
- Schuler U, Schroer S, Kuhnle A, et al. Busulfan pharmacokinetics in bone marrow transplant patients: is drug monitoring warranted? *Bone Marrow Transpl* 1994; 14:759-65.
- Dix SP, Wingard JR, Mullins RE, et al. Association of busulfan area under the curve with veno-occlusive disease following BMT. *Bone Marrow Transpl* 1996; 17:225-30.
- Demirel T, Buckner CD, Appelbaum FR, et al. Busulfan, cyclophosphamide and fractionated total body irradiation for allogeneic marrow transplantation in advanced acute and chronic myelogenous leukemia: phase I dose escalation of busulfan based on targeted plasma levels. *Bone Marrow Transpl* 1996; 17:341-6.
- Zager RA, O'Quigley J, Zager RN, et al. Acute renal failure following bone marrow transplantation: a retrospective study of 272 patients. *Am J Kidney Dis* 1989; 13:210-6.
- Wingard JR, Mellits ED, Jones RJ, et al. Association of hepatic veno-occlusive disease with interstitial pneumonitis in bone marrow transplant recipients. *Bone Marrow Transpl* 1989; 4:685-9.
- Bearman SI, Anderson GL, Mori M, Hinds MS, Shulman HM, McDonald GB. Venocclusive disease of the liver: development of a model for predicting fatal outcome after marrow transplantation. *J Clin Oncol* 1993; 9:1729-36.
- Carreras E, Grañena A, Navasa M, et al. On the reliability of clinical criteria for the diagnosis of hepatic veno-occlusive disease. *Ann*

- Hematol 1993; 66:77-80.
35. Wolford JL, McDonald GB. A problem-oriented approach to intestinal and liver disease after marrow transplantation. *J Clin Gastroenterol* 1988; 20:419-33.
 36. Goodrich JM, Reed EC, Mori M, et al. Clinical features and risk factors for invasive candidal infection after marrow transplantation. *J Infect Dis* 1991; 164:731-40.
 37. Lewis JH, Patel HR, Zimmerman HJ. The spectrum of hepatic candidiasis. *Hepatology* 1982; 2:479-87.
 38. Carreras E, Granena A, Navasa M, et al. Transjugular liver biopsy in BMT. *Bone Marrow Transpl* 1993; 11:21-6.
 39. Shulman HM, Gooley T, Dudley MD, et al. Utility of transvenous liver biopsies and wedged hepatic venous pressure measurements in sixty marrow transplant recipients. *Transplantation* 1995; 59:1015-22.
 40. Iqbal M, Creger RJ, Fox RM, et al. Laparoscopic liver biopsy to evaluate hepatic dysfunction in patients with hematologic malignancies: a useful tool to effect changes in management. *Bone Marrow Transpl* 1996; 17:655-62.
 41. Shulman HM, McDonald GB, Matthews DE, et al. An analysis of hepatic venoocclusive disease and centrilobular hepatic degeneration following bone marrow transplantation. *Gastroenterology* 1980; 79:1178-91.
 42. Katzka DA, Saul SH, Jorkasky D, Sigal H, Reynolds JC, Soloway RD. Azathioprine and hepatic venoocclusive disease in renal transplant patients. *Gastroenterology* 1986; 90:446-54.
 43. Lee RG. Vascular disorders. In: Lee RG, ed. *Diagnostic liver pathology*, Saint Louis: Mosby year book INC; 1994. p. 329-31.
 44. Shulman HM, Gown AM, Nugent DJ. Hepatic venoocclusive disease after bone marrow transplantation: immunohistochemical identification of the material within occluded central venules. *Am J Pathol* 1987; 127:549-58.
 45. Bradbury P, Gordon KC. Connective tissue and stains. In: Bancroft JD, Stevens A, eds. *Theory and practice of histological techniques*, III. New York: Churchill Livingstone; 1990. p. 119-42.
 46. Hommeyer SC, Teefey SA, Jacobson AF, et al. Venocclusive disease of the liver: prospective study of US evaluation. *Radiology* 1992; 184:683-6.
 47. Heikinheimo M, Halila R, Fasth A. Serum procollagen type III is an early sensitive marker for veno-occlusive disease of the liver in children undergoing bone marrow transplantation. *Blood* 1994; 83:3036-40.
 48. Rio B, Bauduer F, Arrago JP, Zittoun R. N-terminal peptide of type III procollagen: a marker for the development of hepatic veno-occlusive disease after BMT and a basis for determining the timing of prophylactic heparin. *Bone Marrow Transpl* 1993; 11:471-2.
 49. Salat B, Holler E, Kolb HJ, et al. Plasminogen activator inhibitor-1 confirms the diagnosis of hepatic veno-occlusive disease in patients with hyperbilirubinemia after bone marrow transplantation. *Blood* 1997; 89:2184-8.
 50. Attal M, Huguet F, Rubie H, et al. Prevention of hepatic veno-occlusive disease after bone marrow transplantation by continuous infusion of low-dose heparin: a prospective, randomised trial. *Blood* 1992; 79:2834-40.
 51. Or R, Nagler A, Shpilberg O, et al. Low molecular weight heparin for the prevention of veno-occlusive disease of the liver in bone marrow transplantation patients. *Transplantation* 1996; 61:1067-71.
 52. Gluckman K, Jolivet I, Scrobohaci ML, et al. Use of prostaglandin E1 for prevention of liver veno-occlusive disease in leukaemic patients treated by allogeneic bone marrow transplantation. *Br J Haematol* 1990; 74:277-81.
 53. Bearman SI, Shen DD, Hinds MS, Hill HA, McDonald GB. A phase I/II study of prostaglandin E1 for the prevention of hepatic venoocclusive disease after bone marrow transplantation. *Br J Haematol* 1993; 84:724-30.
 54. Attal M, Huguet F, Rubie H, et al. Prevention of regimen-related toxicities after bone marrow transplantation by pentoxifylline: A prospective, randomised trial. *Blood* 1993; 82:732-6.
 55. Essell JH, Thompson JM, Harman GS, et al. Pilot trial of prophylactic ursodiol to decrease the incidence of veno-occlusive disease of the liver in allogeneic bone marrow transplant patients. *Bone Marrow Transpl* 1992; 10:367-72.
 56. Essell J, Schroeder M, Thompson J, Harman G, Halvorson R, Callander N. A randomised double-blind trial of prophylactic ursodeoxycholic acid vs placebo to prevent venoocclusive disease of the liver in patients undergoing allogeneic bone marrow transplantation [abstract]. *Blood* 1994; 84 (Suppl 1):250a.
 57. Bearman SI, Lee JL, Baron AE, McDonald GB. Treatment of hepatic venoocclusive disease with recombinant human tissue plasminogen activator and heparin in 42 marrow transplant patients. *Blood* 1997; 89:1501-6.
 58. Richardson P, Krishan A, Wheeler C, et al. The use of defibrotide in BMT-associated veno-occlusive disease [abstract]. *Blood* 1995; 86(Suppl 1):221a.
 59. Murray JA, LaBrecque DR, Gingrich RD, Pringle KC, Mitros FA. Successful treatment of hepatic venoocclusive disease in a bone marrow transplant patient with side-to-side portocaval shunt. *Gastroenterology* 1987; 92:1073-7.
 60. de la Rubia J, Carral A, Montes H, Urquijo JJ, Sanz GF, Sanz MA. Successful treatment of hepatic veno-occlusive disease in a peripheral blood progenitor cell transplant patient with a transjugular intrahepatic portosystemic stent-shunt (TIPS). *Haematologica* 1996; 81:536-9.
 61. Smith FO, Johnson MS, Scherer LR, et al. Transjugular intrahepatic portosystemic shunting (TIPS) for treatment of severe hepatic veno-occlusive disease. *Bone Marrow Transpl* 1996; 18:643-6.
 62. Fried MW, Connaghan DG, Sharma S, et al. Transjugular intrahepatic portosystemic shunt for the management of severe venoocclusive disease following bone marrow transplantation. *Hepatology* 1996; 24:588-91.
 63. Rapoport AP, Doyle HR, Starzl T, et al. Orthotopic liver transplantation for life-threatening veno-occlusive disease of the liver after allogeneic bone marrow transplant. *Bone Marrow Transpl* 1991; 8:421-4.
 64. Dowlati A, Honore P, Damas P, et al. Hepatic rejection after orthotopic liver transplantation for hepatic veno-occlusive disease or graft-versus-host disease following bone marrow transplantation. *Transplantation* 1995; 60:106-9.