

FLT3 ligand kinetic profile predicts response to treatment in patients with high-risk myelodysplastic syndrome/chronic myelomonocytic leukemia receiving CPX-351: a study from the Groupe Francophone des Myélodysplasies

Authors

Pierre Peterlin,^{1,2} Joëlle Gaschet,² Pascal Turlure,³ Marie-Pierre Gourin,³ Pierre-Yves Dumas,⁴ Sylvain Thepot,⁵ Ana Berceanu,⁶ Sophie Park,⁷ Marie-Anne Hospital,⁸ Thomas Cluzeau,⁹ Jose-Miguel Torregrosa-Diaz,¹⁰ Louis Drevon,¹¹ Rosa Sapena,¹² Fatiha Chermat,¹² Lionel Ades,¹¹ Sophie Dimicoli-Salazar,⁴ Maxime Jullien,^{1,2} Pierre Fenaux¹² and Patrice Chevallier^{1,2}

¹Clinical Hematology, Nantes University Hospital, Nantes; ²CRCI2NA UMR INSERM 1307CNRS 6075 - Nantes Université – Angers University, Nantes; ³Clinical Hematology, Limoges University Hospital, Limoges; ⁴Clinical Hematology, Bordeaux University Hospital Haut-Lévêque, Pessac; ⁵Clinical Hematology, Angers University Hospital, Angers; ⁶Clinical Hematology, Besançon University Hospital, Besançon; ⁷Clinical Hematology, Grenoble University Hospital, Grenoble; ⁸Clinical Hematology, Institut Paoli Calmettes, Marseille;

⁹Clinical Hematology, Nice University Hospital, Nice; ¹⁰Clinical Hematology, Poitiers University Hospital, Poitiers; ¹¹Clinical Hematology, Saint Louis Hospital, Paris and ¹²Groupe Francophone des Myélodysplasies, Paris, France

Correspondence:
P. PETERLIN - pierre.peterlin@chu-nantes.fr

<https://doi.org/10.3324/haematol.2024.286025>

Received: June 6, 2024.
Accepted: November 13, 2024.
Early view: November 21, 2024.

©2025 Ferrata Storti Foundation
Published under a CC BY-NC license 

Figure legends

Supplemental figure 1. Forest plot of multivariate analysis.

sFL: soluble Flt3-ligand; FLI: sustained increase from day 1 to day 22; FLD: increase until day 15, then decrease; FLL: persistent low levels; F: female; M: male; IPSS-M: Molecular International Prognostic Scoring System; L:low; H:high;VH: very high; OR: odds ratio.

Supplemental figure 1

