

Chronic lymphocytic leukemia often arises by a multiclonal selection process

Davide Bagnara^{1,2*} Andrea N. Mazzarello^{1,2*} Niccolò Cardente,¹ Stefano Vergani,^{2,3} Siddha Kasar,^{4,5} Stacey Fernandes,⁴ Gerardo Ferrer,^{2,6,7} Fabio Ghiotto,^{1,8} Jacqueline C. Barrientos,^{2,9} Jonathan E. Kolitz,^{2,9} Kanti R. Rai,^{2,9} Steven L. Allen,^{2,9} Monica Colombo,⁸ Franco Fais,^{1,8} Jennifer R. Brown,⁴ Manlio Ferrarini¹ and Nicholas Chiorazzi^{2,9}

¹Department of Experimental Medicine, University of Genoa, Genoa, Italy; ²Karches Center for Oncology Research, The Feinstein Institutes for Medical Research, Northwell Health, Manhasset, NY, USA; ³Miltenyi Biotec B.V. & Co. KG, Bergisch Gladbach, Germany; ⁴CLL Center, Dana-Farber Cancer Institute and Harvard Medical School, Boston, MA, USA; ⁵Takeda Pharmaceuticals, Cambridge, MA, USA; ⁶Josep Carreras Leukaemia Research Institute (IJC), Badalona, Spain; ⁷Centro de Investigación Biomédica en Red Cáncer (CIBERONC), Madrid, Spain; ⁸IRCCS Ospedale Policlinico San Martino, Genoa, Italy and ⁹Departments of Molecular Medicine and of Medicine, Zucker School of Medicine at Hofstra/Northwell, Hempstead, NY, USA

**DB and ANM contributed equally as first authors.*

Correspondence: D. Bagnara
davide.bagnara@unige.it

Received: August 23, 2024.

Accepted: February 12, 2025.

Early view: February 20, 2025.

<https://doi.org/10.3324/haematol.2024.286380>

©2025 Ferrata Storti Foundation

Published under a CC BY-NC license



For Supplementary Tables see Excel files

Supplementary Table 1.

Sequencing data summary of healthy donor IGH repertoire

Supplementary Table 2.

Sequencing data summary of patient with CLL IGH repertoire

Supplementary Table 3.

Summary of CRC and pADC IGHV-D-J sequences