Variables influencing the *in vitro* measurement of spontaneous aggregation of human platelets

Authors

Claudia Ghali,[1,*] Antonella Fioretti,[1,*] Mariangela Scavone,[1] Elena Bossi,[1] Bianca Clerici,[1] Simone Birocchi,[1] Evgeny Popov,[2] Marco Centola,[3] Gianmarco Podda[1] and Marco Cattaneo[4]

[1]Divisione di Medicina Generale II, ASST Santi Paolo e Carlo, Dipartimento di Scienze della Salute, Università degli Studi di Milano, Milano, Italy; [2]National Medical Research Center of Cardiology, Ministry of Health of the Russian Federation, Moscow, Russian Federation; [3]Division of Cardiology, Pio XI Hospital, ASST Brianza, Desio, Italy and [4]Fondazione Arianna Anticoagulazione, Bologna, Italy

CG and AF contributed equally as first authors.

SUPPLEMENTARY

Table 1. Demographic, clinical and laboratory characteristics of the study subjects

	Median (IQR) or n (%)
Demographics	
Sex (females)	50 (50%)
Age (years)	66 (53-76)
Clinical characteristics	
Current smokers	13 (13%)
Hypertension	34 (34%)
Dyslipidaemia	16 (16%)
Type 1 and 2 diabetes	0 (0%)
Cardiovascular disease	0 (0%)
Cerebrovascular disease	0 (0%)
Platelet dysfunction	0 (0%)
Myeloproliferative neoplasms	0 (0%)
Thrombocytopenia	0 (0%)
Chronic obstructive pulmonary disease	2 (2%)
Chronic kidney disease	5 (5%)
Treatments	
ACE inibitors or angiotensin receptor blockers	32 (32%)
Beta blockers	24 (24%)
Statins	16 (16%)
Laboratory values	
Platelets ($\times 10^9$/L)	201 (168-240)