

Beating the STATs: targeting the metabolome in acute myeloid leukemia

by Boaz Nachmias and Ofir Wolach

Received: August 11, 2025.

Accepted: August 27, 2025.

Citation: Boaz Nachmias and Ofir Wolach.

Beating the STATs: targeting the metabolome in acute myeloid leukemia.

Haematologica. 2025 Sept 4. doi: 10.3324/haematol.2025.288349 [Epub ahead of print]

Publisher's Disclaimer.

E-publishing ahead of print is increasingly important for the rapid dissemination of science.

Haematologica is, therefore, E-publishing PDF files of an early version of manuscripts that have completed a regular peer review and have been accepted for publication.

E-publishing of this PDF file has been approved by the authors.

After having E-published Ahead of Print, manuscripts will then undergo technical and English editing, typesetting, proof correction and be presented for the authors' final approval; the final version of the manuscript will then appear in a regular issue of the journal.

All legal disclaimers that apply to the journal also pertain to this production process.

Beating the STATs: targeting the metabolome in acute myeloid leukemia

Boaz Nachmias¹, Ofir Wolach^{2,3*}

¹ Department of Hematology, Hadassah Medical Center and Faculty of Medicine, Hebrew University of Jerusalem, Israel

² Institute of Hematology, Davidoff Cancer Center, Rabin Medical Center, Beilinson Hospital, Petach Tikva, Israel.

³ School of Medicine, Faculty of Medical and Health Sciences, Tel Aviv University, Tel Aviv, Israel.

***Corresponding author**

Ofir Wolach

Institute of Hematology, Davidoff Cancer Center, Rabin Medical Center, Beilinson Hospital, Petach Tikva, Israel.

In the current issue, Gil *et al.* (1) provide new insights on the role of STAT3 in regulating mitochondrial function in acute myeloid leukemia (AML). Beyond its transcriptional activity, STAT3 translocates to the mitochondria where it interacts with voltage-dependant anion channel-1 (VDAC1), a regulator of mitochondrial calcium, oxidative phosphorylation (OXPHOS) and apoptosis. Disrupting this axis impairs mitochondrial metabolism, reduces leukaemic cell viability, and leukaemic stem cell (LSC) engraftment, supporting a role for STAT3 and its downstream effectors as potential therapeutic targets in AML.

AML is an aggressive disease with high rate of relapse contributing to its poor overall survival. Increasing evidence suggests that similar to the hierarchical structure of normal hematopoiesis, a subset of leukaemia stem cells give rise to the bulk leukaemia and often mediate relapse. The seminal work by Craig Jordan as well as others, has uncovered the unique metabolic dependencies of LSC, demonstrating reliance on OXPHOS as a source for energy rather than glycolysis as in normal haematopoietic stem cells (HSC). It was further demonstrated that LSC OXPHOS relies on influx of amino acids and fatty acids, highlighting metabolic vulnerabilities and pathways of resistance (2). Key signal transduction pathways and common mutations were shown to modulate and support the reprogrammed metabolic needs of the leukaemia cells. IDH1/2 mutations were linked to mitochondrial dysregulation with enhanced OXPHOS metabolism. RAS pathway mutations were shown to upregulate MCL-1, with higher rate of fatty acid oxidation (FAO) and amino acid metabolism, as alternative OXPHOS fuel (3). Thus, linking aberrant signaling with metabolic rewiring.

Aberrant activation of the Janus kinase (JAK)/signal transducer and activator of transcription (STAT) pathway, and specifically STAT3 and STAT5 has been increasingly recognized as a key driver in the pathogenesis of both myelodysplastic syndromes (MDS) and AML. Mutations in upstream tyrosine kinases such as FLT3, BCR-ABL, KIT, and components of the RAS pathway commonly enhance STAT3/5 signaling, and functional studies have demonstrated that STAT activation is essential for the leukaemogenic potential of these mutations (4,5).

STAT3 signaling as a metabolic regulator

While STAT proteins are best known for their function as transcription factors, emerging data highlight their non-canonical roles in regulating key metabolic pathways that support leukaemic cell survival and proliferation. Early works have shown that STAT3 can localize to the mitochondria rather than the nucleus, supporting electron transfer and OXPHOS metabolism in

the context of RAS-dependent transformation (5). It was suggested that STAT3 resides in the inner mitochondrial membrane and integrates into electron transport complex I (6). Mitochondrial STAT3 requires phosphorylation at serine 727 (Ser-727), in contrast to the nuclear STAT3 which is phosphorylated on tyrosine 705 (Y705). Interestingly, while both nuclear and mitochondrial STAT3 can modulate cell metabolism, they result in divergent metabolic effects (**Figure 1**). In breast cancer cells, substitution of Ser-727 for an alanine reduced tumor growth, complex I activity, and resulted in accumulation of reactive oxygen species (ROS), suggesting that mitochondrial STAT3 supports OXPHOS potential (7). In contrast, the transcriptional effect of nuclear STAT3 was shown to upregulate HIF-1 α in epithelial tumor cells, supporting glycolysis (8). Work done in primary AML LSC, demonstrated that STAT3 supports OXPHOS *via* upregulation of its target genes including Bcl-2 (9) and MYC, which in turn upregulates the mitochondrial transporter SCL1A5 and influx of glutamine and OXPHOS metabolism. STAT3 silencing in primary AML cells resulted in cell death and reduced engraftment, which was less evident in normal hematopoietic progenitors (10). Finally, STAT3 transcriptional activity was shown to promote several key regulators of lipid metabolism supporting FAO, including peroxisome proliferator-activated receptor gamma (PPAR γ) (11) and carnitine palmitoyltransferase 1B (CPT1B). FAO was shown to support LSC as an alternative fuel, especially under therapeutic pressure of drugs that target OXPHOS, such as the Bcl-2 inhibitor venetoclax, mediating resistance and relapse (2,3). Based on these data, co-targeting STAT3 signaling and OXPHOS metabolism (e.g with azacytidine and venetoclax) may prove synergistic in AML.

In the current issue, Gil et al. provide further evidence for the pivotal role of STAT3 in mitochondrial function and survival of AML cells. The investigators utilized AML cell lines and primary patient samples to demonstrate that STAT3 is preferentially expressed and phosphorylated in the mitochondria of leukaemic cells. They next recognized candidate proteins that interact with STAT3 in the mitochondria by immunoprecipitation assays followed by mass spectrometry analysis and identified VDAC1 for further investigation. Co-immunoprecipitation assays confirmed the intimate interaction between STAT3 and VDAC1.

From a functional standpoint, pharmacologic and transcriptomic inhibition of STAT3 (with Stattic and siSTAT3, respectively) both resulted in reduced mitochondrial calcium and reduction in the mitochondrial membrane potential, an effect also replicated by selective VDAC1 inhibition. VDAC1 overexpression in the context of STAT3 inhibition partially salvaged mitochondrial calcium content suggesting that the effect of VDAC1 is downstream STAT3 activation.

Additionally, STAT3 and VDAC1 inhibition reduced OXPHOS measures such as oxygen consumption rates and ROS production with a net reduction in oxidative stress. The deleterious effects of STAT3 and VDAC1 inhibition on OXPHOS and mitochondrial calcium metabolism were also reflected by reduced mitochondrial mass in treated leukemic cells.

Finally, Gil et al. demonstrate that inhibition of STAT3 with Stattic reduced the growth and viability of LSC, including engraftment of primary AML cells in a mouse xenograft model. These data complement other pre-clinical studies demonstrating that Stattic can induce apoptosis and decrease leukaemic stem cell activity, particularly in FLT-3-mutant AML cells (12). Other STAT3 inhibitors either alone or in combination with other therapies such as venetoclax have shown efficacy in vitro and in mouse xenograft models, while clinical data is still limited (13).

One important aspect of this work is that the STAT3-VDAC1 pathway remained relevant to mitochondrial function and LSC survival in the context of venetoclax resistance, a yet unmet challenge with few therapeutic options.

In conclusion, the current study by Gil et al. as well as previous works, highlight the role of STAT3 in the metabolic adaptation of leukaemic cells. As evidence accumulates, targeting metabolic pathways regulated by STATs, either alone or in combination with existing therapies, offer a complementary approach to current treatment strategies.

References


1. Gil KB, Borg J, Pereira RM, et al. The STAT3-VDAC1 axis modulates mitochondrial function and plays a critical role in the survival of acute myeloid leukemia cells. *Haematologica*. xxx
2. Jones CL, Inguva A, Jordan CT. Targeting Energy Metabolism in Cancer Stem Cells: Progress and Challenges in Leukemia and Solid Tumors. *Cell Stem Cell*. 2021;28(3):378-393.
3. Nachmias B, Aumann S, Haran A, Schimmer AD. Venetoclax resistance in acute myeloid leukaemia-Clinical and biological insights. *Br J Haematol*. 2024;204(4):1146-1158.
4. Choudhary C, Brandts C, Schwable J, et al. Activation mechanisms of STAT5 by oncogenic Flt3-ITD. *Blood*. 2007;110(1):370-374.
5. Gough DJ, Corlett A, Schlessinger K, Wegrzyn J, Larner AC, Levy DE. Mitochondrial STAT3 supports Ras-dependent oncogenic transformation. *Science*. 2009;324(5935):1713-1716.
6. Tammineni P, Anugula C, Mohammed F, Anjaneyulu M, Larner AC, Sepuri NBV. The Import of the Transcription Factor STAT3 into Mitochondria Depends on GRIM-19, a Component of the Electron Transport Chain. *J Biol Chem*. 2013;288(7):4723-4732.
7. Zhang Q, Raje V, Yakovlev VA, et al. Mitochondrial localized Stat3 promotes breast cancer growth via phosphorylation of serine 727. *J Biol Chem*. 2013;288(43):31280-31288.
8. Demaria M, Giorgi C, Lebedzinska M, et al. A STAT3-mediated metabolic switch is involved in tumour transformation and STAT3 addiction. *Aging*. 2010;2(11):823-842.
9. Sepúlveda P, Encabo A, Carbonell-Uberos F, Miñana MD. BCL-2 expression is mainly regulated by JAK/STAT3 pathway in human CD34+ hematopoietic cells. *Cell Death Differ*. 2007;14(2):378-380.
10. Amaya ML, Inguva A, Pei S, et al. The STAT3-MYC axis promotes survival of leukemia stem cells by regulating SLC1A5 and oxidative phosphorylation. *Blood*. 2022;139(4):584-596.
11. Wang D, Zhou Y, Lei W, et al. Signal transducer and activator of transcription 3 (STAT3) regulates adipocyte differentiation via peroxisome-proliferator-activated receptor gamma (PPARgamma). *Biol Cell*. 2009;102(1):1-12.
12. Luo Y, Lu Y, Long B, et al. Blocking DNA Damage Repair May Be Involved in Stattic (STAT3 Inhibitor)-Induced FLT3-ITD AML Cell Apoptosis. *Front Cell Dev Biol*. 2021;9:637064.
13. Shastri A, Choudhary G, Teixeira M, et al. Antisense STAT3 inhibitor decreases viability of myelodysplastic and leukemic stem cells. *J Clin Invest*. 2018;128(12):5479-5488.

Figure 1: STAT3 regulates mitochondrial and nuclear metabolic pathways in AML.

(a) Schematic representation of STAT3 activity in the nucleus and mitochondria. Nuclear STAT3 promotes transcription of key metabolic genes. In the mitochondria, phosphorylated (S727) STAT3 interacts with VDAC1, supporting OXPHOS metabolism. **(b)** Targeted inhibition of STAT3 or VDAC1 disrupts mitochondrial function, leading to impaired OXPHOS and reduced leukaemic cell viability and stemness.

Abbreviations: STAT3- signal transducer and activator of transcription 3, HIF1a- hypoxia-inducible factor 1, OXPHOS-oxidative phosphorylation, BCL-2- B cell lymphoma 2, PARP- poly (ADP-ribose) polymerase, FASN- fatty acid synthase, DIDS- 4,4'-Diisothiocyanatostilbene-2,2'-disulfonic acid (VDAC1 inhibitor).

A



B

