

The “morning cup of joe” as a modifiable factor in blood transfusion

Tiffany A. Thomas and Steven L. Spitalnik

Laboratory of Transfusion Biology, Department of Pathology & Cell Biology, Columbia University, New York, NY, USA

Correspondence: S.L. Spitalnik
ss2479@cumc.columbia.edu

Received: September 26, 2025.

Accepted: October 8, 2025.

Early view: October 16, 2025.

<https://doi.org/10.3324/haematol.2025.288836>

©2026 Ferrata Storti Foundation

Published under a CC BY-NC license



Precision medicine concepts can apply to red blood cell (RBC) transfusions, as follows: transfusing the right donor unit with the right quality into the right recipient in the right clinical setting at the right time to achieve an optimal outcome. Nonetheless, this requires a more nuanced definition of “RBC quality.” Thus, “storage quality” can be defined as the ability of RBC to withstand the rigors of refrigerated storage, typically measured by storage hemolysis, 24-hour post-transfusion RBC recovery, and post-transfusion hemoglobin increment. In addition, “post-transfusion circulatory quality” can correspond to the long-term circulatory lifespan of the transfused RBC, and their ability to perfuse and oxygenate tissues, which depend, for example, on RBC deformability and 2,3-biphosphoglycerate (2,3-BPG) levels, respectively.

RBC transfusion efficacy has improved dramatically over the last 100+ years. For example, ABO blood typing improved selection of donor/recipient pairs, thereby preventing virtually all acute hemolytic transfusion reactions. Nonetheless, blood group alloimmunization still occurs, leading to delayed hemolytic transfusion reactions and difficulties in finding compatible units. In addition, improved anticoagulant and storage solutions, along with appropriate storage containers, allow for rapid RBC unit availability, efficient inventory management, and access to rare units. Identifying microbial pathogens transmitted by RBC transfusion, and developing relevant diagnostic tests, virtually eliminated transfusion-transmitted diseases, particularly in high-income countries, and pathogen-reduction technologies may soon solve this issue globally. Moreover, universal pre-storage leukoreduction (accompanied by irradiation) decreases HLA alloimmunization, febrile transfusion reactions, and transfusion-associated graft-versus-host disease. Finally, consensus statements and patient blood management approaches help to optimize utilization of RBC transfusions and/or transfusion alternatives (e.g., pre-operative iron supplementation). Given these advances, one could claim that no further improvements are necessary; nonetheless, issues do remain.

For example, we do not understand transfusion-related immunomodulation or the critical mechanisms underlying alloimmunization to RBC antigens; because of the latter, we still encounter highly alloimmunized, chronically transfused patients who become “untransfusable” because no compatible units are attainable. In addition, the “RBC storage lesion” worsens over time; thus, transfusing 42-day-old RBC differs from transfusing 7-day-old RBC, due to infusing potentially damaging supernatant constituents (e.g., extracellular vesicles, mitochondrial DNA, non-transferrin-bound iron), providing less than an optimal dose (due to decreased post-transfusion RBC recovery), and providing RBC with decreased deformability and 2,3-BPG (affecting perfusion and oxygen offloading, respectively). These issues can be partially rectified by “washing” or “rejuvenating” RBC units, but this increases cost and decreases the therapeutic dose. Nonetheless, further improvements may not matter in certain scenarios, such as transfusing several units into a “transfusion-naïve” patient who is unlikely to ever be transfused again (e.g., an elderly male patient receiving a hip replacement), or when rapidly transfusing many units into a patient experiencing a massive, acute hemorrhage (e.g., in placenta accreta). However, in chronic transfusion settings (e.g., myelodysplastic syndrome, hemoglobinopathies), providing higher quality units could permit transfusion of 10–20% fewer units and/or fewer transfusion episodes, thereby improving quality of life (e.g., fewer missed days of school, work, or leisure), minimizing adverse outcomes (e.g., transfusion reactions, central line complications), and lessening iron overload. Moreover, high-quality RBC may be important in particular high-risk situations (e.g., extremely premature neonates).¹ One way to improve overall RBC transfusion quality is to enhance understanding of the RBC storage lesion and develop approaches to ameliorate it. To this end, current research leverages: mass spectrometry-enabled metabolomics, lipidomics, and proteomics; large epidemiological studies, including “vein-to-vein databases” (coupled with screening donors for RBC-relevant genetic polymorphisms); clinical

trials with healthy volunteers and patients; and knockout and transgenic mouse models targeting RBC-relevant genes. These studies have identified multiple factors affecting RBC transfusion quality (Table 1). Moreover, although mature RBC, lacking virtually all organelles, are considered “simple,” these studies, in both normal and abnormal settings, discovered many more protein constituents and active metabolic pathways.² In addition, exposure to their environment dramatically affects RBC biology; indeed, their “exposome” includes microbiome products, pharmaceuticals (licit and illicit), diet (e.g., caffeine, vitamins, fatty acids), and the external environment (e.g., lead, smoking).³ In the current issue of *Haematologica*, Dzieciatkowska *et al.*⁴ exploited these approaches to study whether caffeine modulates RBC storage quality and post-transfusion circu-

Table 1. Examples of factors affecting red blood cell storage quality.

Factor	Effect on hemoglobin increment	Reference
Pre-storage leukoreduction of donor unit	Decrease	13
Irradiation of donor unit	Decrease	13
Donor glucose-6-phosphate dehydrogenase deficiency	Decrease	10, 13
Storage duration of donated unit	Decrease	13
Male donor & female recipient	Increase	13
Donor caffeine consumption	Decrease	4




Figure 1. Red blood cell metabolic pathways affected by caffeine exposure. Schematic diagram indicating that caffeine is both a competitive inhibitor of adenosine binding to the ADORA2b adenosine receptor, which downregulates subsequent downstream signaling through AMP-activated protein kinase, and a direct inhibitor of glucose-6-phosphate dehydrogenase, the rate-limiting enzyme in the pentose phosphate pathway. Taken together, this demonstrates that caffeine negatively affects the three major red blood cell pathways that utilize glucose (i.e., glycolysis, the Luebering-Rapoport pathway, and the pentose phosphate pathway), thereby inhibiting their synthetic capacities (downward facing red arrows) to generate energy (i.e., ATP), enhance oxygen offloading by hemoglobin (i.e., 2,3-BPG); and resist storage-induced oxidative stress (i.e., NADPH); respectively. ADORA2b: an adenosine receptor; AMPK: AMP-activated protein kinase; ATP: adenosine triphosphate; BPGM: biphosphoglycerate mutase; 1,7DMU: 1,7-dimethyluric acid; 1,3,7TMU: 1,3,7-trimethyluric acid; CYP1A2 and CYP2A6: two cytochrome p450 isoforms; LRP: Luebering-Rapoport pathway; GLUT1: glucose transporter 1; Glc-6-P: glucose-6-phosphate; G6PD: glucose-6-phosphate dehydrogenase; PPP: pentose phosphate pathway; RBC: red blood cell.

latory quality. Surprisingly, this psychoactive agent, widely consumed in coffee,⁵ negatively affects RBC transfusion quality. Although coffee is often an inducement, or reward, for blood donation, and although it may help to prevent donation-induced vasovagal reactions,⁶ the current results, coupled with caffeine's short half-life *in vivo*, suggest that modifying its use could readily improve RBC transfusion quality.

The research by Dzieciatkowska *et al.*⁴ demonstrates that caffeine affects RBC physiology through extracellular and intracellular mechanisms (Figure 1). On the RBC surface, caffeine directly blocks adenosine receptor ADORA2b, thereby inhibiting signaling through AMP-activated protein kinase.⁷ In addition, caffeine rapidly enters (and exits) RBC by passive diffusion,⁸ and binds to, and inhibits, glucose-6-phosphate dehydrogenase (G6PD), the rate-limiting enzyme of the pentose phosphate pathway (PPP). Thus, caffeine significantly affects the three major RBC pathways utilizing glucose: glycolysis (for ATP synthesis), the Luebering-Rapoport pathway (for 2,3-BPG synthesis), and the PPP (for synthesizing NADPH to ameliorate oxidative stress). Given the need to maintain energy levels during storage, RBC redirect glucose metabolism towards ATP generation, thereby limiting 2,3-BPG and NADPH synthesis; caffeine exacerbates the latter, further compromising the ability of the RBC to prevent damage from storage-induced oxidative stress (Figure 1).⁹

The study by Dzieciatkowska *et al.*⁴ also identifies relevant gene-exposome interactions. For example, caffeine inhibits wild-type G6PD, but not the common African variant associated with G6PD deficiency. Thus, caffeine does not additively or synergistically contribute to storage-induced oxidative stress of G6PD-deficient RBC, as might have been expect-

ed.¹⁰ Therefore, elaborating the exact mechanism(s) by which caffeine inhibits G6PD activity is important.¹¹ In contrast, the negative effects of certain human *ADORA2b* polymorphisms on RBC storage quality, as with the *ADORA2b*-knockout mouse, are exacerbated by caffeine's effects on (wild-type) G6PD enzyme activity. Finally, given that hepatic CYP2A6 metabolizes caffeine, its genetic polymorphisms affect caffeine clearance from plasma and, consequently, from RBC. The need for liver-dependent caffeine metabolism also explains why caffeine levels do not decrease during RBC storage.

Finally, given plasma caffeine's rapid metabolism (its $t_{1/2}$ is ~5 hours) and its rapid equilibration with RBC, short-term abstinence would dramatically decrease RBC concentrations, which could be beneficial for donated RBC (if donors would comply!). However, it is currently unknown whether there is permanent structural damage to circulating RBC¹² from any additional oxidative stress RBC may experience due to caffeine exposure, which may be maintained throughout storage; this also merits further study.

In summary, one can now think of caffeine as an extensively studied avatar of the RBC exposome. In addition, caffeine, which interacts with several polymorphic human genes (e.g., *ADORA2b*, *G6PD*, *CYP2A6*), can be considered as a modifiable factor that could improve the clinical practice of precision transfusion medicine.

Disclosures

TAT has no conflicts of interest to disclose. SLS is a member of the scientific advisory boards of Hemanext, Inc. and Alcor, Inc.

Contributions

Both authors contributed equally to this editorial.

References

1. Teofili L, Papacci P, Pellegrino C, et al. Cord red blood cell transfusions for severe retinopathy in preterm neonates in Italy: a multicenter randomized controlled trial. *EClinicalMedicine*. 2025;87:103426.
2. Haiman ZB, Key A, D'Alessandro A, Palsson BO. RBC-GEM: a genome-scale metabolic model for systems biology of the human red blood cell. *PLOS Comput Biol*. 2025;21(3):e1012109.
3. Nemkov T, Stefanoni D, Bordbar A, et al. Blood donor exposome and impact of common drugs on red blood cell metabolism. *JCI Insight*. 2021;6(3):e146175.
4. Dzieciatkowska M, Hay A, Issaian A, et al. Caffeine impairs red blood cell storage quality by dual inhibition of ADORA2b signaling and G6PD activity. *Haematologica*. 2026;111(3):861-875.
5. van Dam RM, Hu FB, Willett WC. Coffee, caffeine, and health. *N Engl J Med*. 2020;383(4):369-378.
6. Sauer LA, France CR. Caffeine attenuates vasovagal reactions in female first-time blood donors. *Health Psychol*. 1999;18(4):403-409.
7. Föllner M, Sopjani M, Koka S, et al. Regulation of erythrocyte survival by AMP-activated protein kinase. *FASEB J*. 2009;23(4):1072-1080.
8. Arnaud MJ. Pharmacokinetics and metabolism of natural methylxanthines in animal and man. In: Methylxanthines. Fredholm BB (ed). Springer Berlin Heidelberg; 2010. p. 33-91.
9. Dumaswala UJ, Zhuo L, Jacobsen DW, Jain SK, Sukalski KA. Protein and lipid oxidation of banked human erythrocytes: Role of glutathione. *Free Radic Biol Med*. 1999;27(9-10):1041-1049.
10. Francis RO, D'Alessandro A, Eisenberger A, et al. Donor glucose-6-phosphate dehydrogenase deficiency decreases blood quality for transfusion. *J Clin Invest*. 2020;130(5):2270-2285.
11. Xu H, Hu L, Liu T, et al. Caffeine targets G6PDH to disrupt redox homeostasis and inhibit renal cell carcinoma proliferation. *Front Cell Dev Biol*. 2020;8:556162.
12. Peltier S, Marin M, Dzieciatkowska M, et al. Proteostasis and metabolic dysfunction characterize a subset of storage-induced senescent erythrocytes targeted for posttransfusion clearance. *J Clin Invest*. 2025;135(9):e183099.
13. Roubinian NH, Reese SE, Qiao H, et al. Donor genetic and nongenetic factors affecting red blood cell transfusion effectiveness. *JCI Insight*. 2022;7(1):e152598.