

Unemployment in patients with histiocytic neoplasms is frequent and associated with clinical and treatment characteristics

Authors

Anne S. Reiner,¹ Dana Bossert,² Allison M. Sigler,² Jen Silvers,³
Kathleen Brewer,⁴ Diane Schriener,⁴ Priya H. Marathe,⁵ Jun J. Mao,⁶
Katherine S. Panageas¹ and Eli L. Diamond^{2,7}

¹Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY; ²Department of Neurology; Memorial Sloan Kettering Cancer Center, New York, NY; ³Histiocytosis Association, Pitman, NJ; ⁴Erdheim-Chester Disease Global Alliance, DeRidder, LA; ⁵Department of Pediatrics, Memorial Sloan Kettering Cancer Center, New York, NY; ⁶Integrative Medicine Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY and ⁷Early Drug Development Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

Correspondence:

ELI L. DIAMOND - diamone1@mskcc.org

<https://doi.org/10.3324/haematol.2025.289304>

Received: September 29, 2025.

Accepted: November 24, 2025.


Early view: December 4, 2025.

©2026 Ferrata Storti Foundation

Published under a CC BY-NC license 

Supplemental Figure 1: Distribution of employment status changes over time.
A) from enrollment to Timepoint 1. B) from enrollment to Timepoint 2.

A



B

