

Metformin: interferon's new dancing partner?

by Hans C. Hasselbalch

Received: January 22, 2026.

Accepted: February 10, 2026.

Citation: Hans C. Hasselbalch. Metformin: interferon's new dancing partner?
Haematologica. 2026 Feb 19. doi: 10.3324/haematol.2026.300445 [Epub ahead of print]

Publisher's Disclaimer.

E-publishing ahead of print is increasingly important for the rapid dissemination of science.

Haematologica is, therefore, E-publishing PDF files of an early version of manuscripts that have completed a regular peer review and have been accepted for publication.

E-publishing of this PDF file has been approved by the authors.

After having E-published Ahead of Print, manuscripts will then undergo technical and English editing, typesetting, proof correction and be presented for the authors' final approval; the final version of the manuscript will then appear in a regular issue of the journal.

All legal disclaimers that apply to the journal also pertain to this production process.

Metformin: interferon's new dancing partner ?

Hans C. Hasselbalch

Department of Hematology, Zealand University Hospital, Roskilde, Denmark

Correspondence : hans.hasselbalch@gmail.com

Disclosures: Research Grant Novartis Healthcare A/S ; Research Grant Pharmaand GmbH ; Advisory Board AOP Orphan

Introduction

In this issue of *Haematologica*, Reeves et al. report a seminal case series of 11 patients with MPNs (PV=6 and ET=5), who were intolerant of pegylated interferon-alpha2b (ropeg) (BESREMI®). Adjunct metformin treatment improved ropeg-related fatigue and/or myalgias in 10 of the 11 patients. Indeed, in 4 of them the side effects completely resolved and in some patients metformin also improved the hematological response.¹ This elegant clinical observation may have paradigm-shifting and practice-changing implications. For decades, interferon- α (IFN) has been recognized as a disease-modifying therapy in MPNs, capable of inducing molecular remissions and normalizing hematopoiesis.² Yet, treatment-limiting side effects have prevented many patients from realizing its full potential. Recently, Danish studies have shown that combination therapy with pegylated interferon alfa-2a (Peg-IFN- α 2a) and ruxolitinib (RUX) is safe and efficacious in patients with polycythemia vera (PV) and myelofibrosis (COMBI I). Most patients in this study were intolerant or refractory to Peg-IFN- α 2a monotherapy, and the majority were also refractory or intolerant to hydroxyurea.³ This combination therapy improved cell counts, reduced bone marrow cellularity and fibrosis, decreased *JAK2 V617F* burden, alleviated IFN-side effects and accordingly reduced symptom burden in a substantial proportion of the patients.³ A prospective study of this combination therapy (COMBI II) in 25 newly diagnosed PV patients confirmed the highly encouraging effects of this doublet therapy with rapid normalization of blood cell counts, low-burden *JAK2V617F* and even normalization of the bone marrow in a subset of patients.⁴ However, this combination therapy implies an increased risk of infections, which are well-known during treatment with ruxolitinib monotherapy and recorded in the COMBI-trials as well.^{3,4} Therefore, an inexpensive, safe and efficacious combinatorial approach with IFN is urgently needed to rescue the large number of patients, who are intolerant or refractory to IFN monotherapy. This excellent report by Reeves et al. opens the door to reintroducing IFN in thousands of patients worldwide who were previously deemed intolerant, by combining it with metformin - a simple, inexpensive, and well-tolerated drug with pleiotropic metabolic and anti-inflammatory effects⁵ in addition to potent anti-leukemic effects in *JAK2V617F*-positive MPNs.⁶

Metformin: An Unexpected Ally

Metformin, a cornerstone in the management of type 2 diabetes, exerts broad systemic effects beyond glucose control.^{5,6} It activates AMP-activated protein kinase (AMPK), reduces oxidative stress, inhibits mTOR signaling, and dampens chronic inflammation - all of which are relevant to MPN pathophysiology and interferon tolerance. Importantly, metformin improves mitochondrial bioenergetics and may counteract the flu-like, fatigue-related, and metabolic side effects of IFN. It also exerts antiproliferative and antineoplastic properties in MPN cells, potentially synergizing with IFN's immunomodulatory and differentiation-inducing mechanisms. These effects may explain why metformin also has been shown to protect against development of MPNs in the background population.⁷ Given this dual impact — mitigating toxicity and possibly enhancing efficacy — metformin should be considered not only as a rescue agent for IFN-intolerant patients but as a rational up-front partner.

Statins: The Second Partner in the Dance ?

Recent Danish registry-based and translational studies have shown that statins – similar to metformin – reduces the risk of developing MPNs⁸, thereby potentially reducing disease progression and improving survival in MPNs - likely through anti-inflammatory, endothelial-protective, and JAK2-signaling modulating mechanisms. Even more intriguing, statins appear to potentiate the efficacy of IFN⁹, possibly by lowering the inflammatory drive and thereby normalizing the bone marrow microenvironment. In this context, it is most intriguing that both metformin and statins possess anti-fibrotic capabilities¹⁰ and therefore, in the best of all scenarios, might also favourably and synergistically impact the development of bone marrow fibrosis in patients with MPN.

Together, IFN, metformin, and statins may form a triad of anti-inflammatory, anti-clonal, and anti-aging agents targeting key nodes in MPN pathogenesis — oxidative stress, inflammation (“inflammaging”), and metabolic dysfunction. Such a combinatorial potent triplet approach would converge on cholesterol-driven hematopoietic activation via complementary mechanisms – metformin enhancing cholesterol efflux capacity¹¹ and statins reducing LDL-driven input¹², thereby dampening inflammatory myelopoiesis and atherosclerotic inflammation. At the same time, IFN would be even more engaged and beloved by both partners and by our patients.

A New Era of Combination Therapy

Although the combination of IFN and RUX has been shown to be promising in patients with PV and MF^{2,3}, it needs to be validated in well-designed prospective studies to confirm its superiority to monotherapy, which has not been confirmed in patients with myelofibrosis. The combination of IFN with metabolic modulators such as metformin and statins could represent the next major therapeutic advance in the field. These agents are safe, inexpensive, and globally available, and their pleiotropic effects align perfectly with the chronic inflammatory and metabolic hallmarks of MPNs. It is now imperative to design prospective trials to validate metformin's ability to rescue IFN-intolerant patients, evaluate its potential as a first-line adjunct to IFN and explore synergistic regimens combining IFN, metformin, and statins for optimal molecular and clinical outcomes.

Conclusion and Perspectives

The work by Reeves et al reminds us that scientific progress often begins with clinical observation. Metformin's capacity to make interferon tolerable again could transform MPN therapy, enabling more patients to receive a treatment capable of deep molecular remissions and potential cure. Future studies should map the molecular crosstalk between metabolism, inflammation, and the malignant clone. In this regard, it is highly intriguing to note that metformin possesses strong anti-atherosclerotic effects and suppresses mobilization of hematopoietic stem cells and progenitors (HSPCs), thereby dampening myelopoiesis, which is a key driver of atherosclerotic inflammation and plaque formation. Metformin exerts these effects by upregulating the cholesterol transporter Abca1 in hematopoietic cells. By these effects metformin may act in synergy with statins, which reduce HSPCs by lowering LDL and modulating mobilizing cytokine pathways.^{11,12} The time has come to explore "the right partners for interferon to dance with" — and both metformin and statins may truly be great companions (Figure 1). It is prime time for revisiting these two old drugs for the treatment of MPNs.

References

1. Reeves BN, Song J, Leier A, et al. Metformin alleviates side effects and supports the resumption of interferon therapy in polycythemia vera and essential thrombocythemia. *Haematologica*. xxx
2. Hasselbalch HC, Silver RT. New Perspectives of Interferon-alpha2 and Inflammation in Treating Philadelphia-negative Chronic Myeloproliferative Neoplasms. *Hemasphere*. 2021;5(12):e645.
3. Sørensen AL, Mikkelsen SU, Knudsen TA, et al. Ruxolitinib and interferon- α 2 combination therapy for patients with polycythemia vera or myelofibrosis: a phase II study. *Haematologica*. 2020;105(9):2262-2272.
4. Sørensen AL, Skov V, Kjær L, et al. Combination therapy with ruxolitinib and pegylated interferon-alfa2a in newly diagnosed patients with polycythemia vera. *Blood Adv*. 2024;8(20):5416-5425.
5. Rehman A, Satyam SM, El-Tanani M, et al. Metformin Beyond Diabetes: A Precision Gerotherapeutic and Immunometabolic Adjuvant for Aging and Cancer. *Cancers*. 2025;17(15):2466.
6. Machado-Neto JA, Fenerich BA, Scopim-Ribeiro R, et al. Metformin exerts multitarget antileukemia activity in JAK2V617F-positive myeloproliferative neoplasms. *Cell Death Dis*. 2018;9(3):311.
7. Kristensen DT, Øvlisen AK, Jakobsen LHK, et al. Metformin use and risk of myeloproliferative neoplasms: a Danish population-based case-control study. *Blood Adv*. 2024; 8(16):4478-4485.
8. Kristensen DT, Kisbye Øvlisen A, Hjort Kyneb Jakobsen L, et al. Use of statins and risk of myeloproliferative neoplasms: a Danish nationwide case-control study. *Blood Adv*. 2023;7(14):3450-3457.

9. Davidsen ID, Larsen MK, Löffler L, et al. Statins enhance the efficacy of pegylated interferon- α 2 in Philadelphia-negative chronic myeloproliferative neoplasms. *Blood Adv.* 2025;9(23):6189-6198.
10. Wu M, Xu H, Liu J, et al . Metformin and Fibrosis: A Review of Existing Evidence and Mechanisms. *J Diabetes Res.* 2021;2021:6673525.
11. Lee MKS, Cooney OD, Zhu Y, et al. Metformin suppresses atherosclerosis by dampening extramedullary myelopoiesis. *Am J Physiol Endocrinol Metab.* 2025 ;329(3):E420-E432.
12. Cimato TR, Palka BA, Lang JK, Young RF. LDL cholesterol modulates human CD34+ HSPCs through effects on proliferation and the IL-17 G-CSF axis. *PLoS One.* 2013(8):e73861.

Figure 1. The Interferon “Dancing Partners” in MPN Therapy. Schematic illustration showing pegylated interferon- α 2 (IFN) at the center of a square of combination partners. **Left** : A healthy lifestyle with exercise to improve metabolism and immune fitness is the key preventive medicine across all disciplines to dampen chronic inflammation, thereby enhancing IFN-tolerability, reducing fatigue and prohibiting accelerated immune- and inflammaging. **Top**: Metformin improves metabolic resilience, reduces oxidative stress, and enhances IFN tolerance. **Right** : Statins exert anti-inflammatory and endothelial-stabilizing effects, potentially enhancing IFN efficacy. **Base**: The malignant clone and inflammatory microenvironment are the targets of the synergistic triad as shown in the combination trials with IFN and ruxolitinib. Together, the combination acts to reduce “inflammaging,” restore hematopoietic homeostasis, and hopefully will improve long-term outcomes.

