

Balancing remission durability and treatment burden in older acute myeloid leukemia. Comment on: "Long-term follow-up of oral decitabine/cedazuridine plus venetoclax for older or unfit patients with newly diagnosed acute myeloid leukemia"

by Yushan Jin, Shiyuan Li and Si Li

Received: July 10, 2026.

Accepted: July 15, 2026.

Citation: Yushan Jin, Shiyuan Li and Si Li. Balancing remission durability and treatment burden in older acute myeloid leukemia. Comment on: "Long-term follow-up of oral decitabine/cedazuridine plus venetoclax for older or unfit patients with newly diagnosed acute myeloid leukemia".

Haematologica. 2026 July 23. doi: 10.3324/haematol.2026.301606 [Epub ahead of print]

Publisher's Disclaimer.

E-publishing ahead of print is increasingly important for the rapid dissemination of science.

Haematologica is, therefore, E-publishing PDF files of an early version of manuscripts that have completed a regular peer review and have been accepted for publication.

E-publishing of this PDF file has been approved by the authors.

After having E-published Ahead of Print, manuscripts will then undergo technical and English editing, typesetting, proof correction and be presented for the authors' final approval, the final version of the manuscript will then appear in a regular issue of the journal.

All legal disclaimers that apply to the journal also pertain to this production process.

Balancing remission durability and treatment burden in older acute myeloid leukemia. Comment on: "Long-term follow-up of oral decitabine/cedazuridine plus venetoclax for older or unfit patients with newly diagnosed acute myeloid leukemia"

Yushan Jin ^{1,2}, Shiyuan Li^{1*}, Si Li^{1*}

¹ The First Affiliated Hospital of Dalian Medical University, Dalian, China

² College of Laboratory Medicine, Dalian Medical University, Dalian, China

*** Corresponding authors:**

Shiyuan Li, The First Affiliated Hospital of Dalian Medical University, Dalian, China, E-mail: shuaili2015@outlook.com

Si Li, The First Affiliated Hospital of Dalian Medical University, Dalian 116011, China, E-mail: jys18804048313@outlook.com

Authors' contributions

All authors contributed equally to the conception and drafting of the manuscript and approved the submitted version.

Disclosure

The authors declare no conflict of interest.

Dear Editor,

Htut TW and colleagues address an important question in hematology: Long-term follow-up of oral decitabine/cedazuridine plus venetoclax for older or unfit patients with newly diagnosed acute myeloid leukemia.¹ The article is valuable because it tackles a clinically recognizable problem and provides a useful basis for discussion. The key issue, however, is how long-term outcomes with oral decitabine/cedazuridine plus venetoclax should be interpreted in older or unfit acute myeloid leukemia patients with competing risks. This distinction matters because readers may otherwise move too quickly from an interesting observation to a clinical or methodological conclusion that the present evidence does not fully establish. We therefore believe the article should be read not only for its main result, but also for the assumptions that determine how far that result can travel.

The first issue is that older or unfit AML populations are shaped by early mortality, treatment interruption, infections, transfusion dependence, and subsequent therapy, all of which can complicate survival interpretation. The second is that long-term follow-up is valuable, but durability should be reported together with time on therapy, dose reductions, hospitalization burden, and quality-of-life or functional outcomes when available.^{2, 3} These concerns do not diminish the contribution of the study, but they define the conditions under which the findings can be applied. A strong letter should make this boundary visible, because the most important risk for readers is not that the result is uninteresting, but that its practical meaning is broader than the data can support. In this sense, the central question is not whether the work is useful; it is what exact decision, hypothesis, or future study the work can legitimately support.

This distinction has practical consequences for interpretation. If the study is read as definitive, readers may focus on the headline conclusion and overlook design features that shape the estimate, such as patient selection, timing of measurement, endpoint definition, missing data, operator behavior, or treatment pathway after enrollment. If it is read as a carefully bounded contribution, however, the same findings become more useful because they identify where the evidence is strong and where replication or refinement is still required. That framing is especially important for studies likely to influence clinical language before they change clinical practice.

A third issue is that subgroup analyses by baseline fitness, cytogenetic or molecular risk, and early response depth would help clinicians identify who is most likely to benefit.⁴ The next study should therefore be designed around a small number of prespecified questions: which subgroup is most likely to benefit from the proposed interpretation, which outcome should be considered decisive, and what level of follow-up, technical quality, or data completeness would make the conclusion reproducible. Answering these questions would help move the field from an interesting report to a usable clinical or methodological standard. Such a design would also make negative or neutral findings more informative, because it would show whether the original signal fails generally or only under specific biological, technical, or operational conditions.

The findings should also be discussed with attention to negative and neutral results. A focused letter is most useful when it protects readers from a one-directional interpretation of promising data. If the reported strategy performs well only under selected conditions, that limitation should be visible early. If the observed effect is clinically attractive but operationally difficult, implementation barriers should be treated as part of the evidence rather than as a secondary concern. This is not a conservative posture for its own sake. It is a way to keep promising results actionable without turning them into recommendations before the necessary comparative or external evidence exists.

We would therefore encourage future work to report the elements that determine transportability. For clinical studies, these include baseline risk, co-interventions, follow-up intensity, competing events, safety, and patient-centered outcomes. For biomarker, registry, artificial-intelligence, or methodological studies, they include calibration, missingness, data provenance, annotation quality, temporal drift, and external validation. Making these elements explicit would allow readers to distinguish a robust finding from a finding that is promising but context-dependent.

In summary, Long-term follow-up of oral decitabine/cedazuridine plus venetoclax for older or unfit patients with newly diagnosed acute myeloid leukemia is a meaningful contribution to the literature.¹ The regimen may be useful, but its value depends on balancing remission durability against treatment burden in a frail population. Its greatest value lies in opening a practical discussion about how the reported evidence should be used, verified, and communicated. Clearer boundaries can strengthen, rather than weaken, the impact of a well-conducted study, because they show readers exactly what should change now and what still requires confirmation.

Reference

- [1] Htut TW, Keketjian J, Short N, et al. Long-term follow-up of oral decitabine/cedazuridine plus venetoclax for older or unfit patients with newly diagnosed acute myeloid leukemia. *Haematologica*. 2026 July 2. doi: 10.3324/haematol.2026.301126. [Epub ahead of print]
- [2] DiNardo CD, Erba HP, Freeman SD, Wei AH. Acute myeloid leukaemia. *Lancet*. 2023;401(10393):2073-2086.
- [3] DiNardo CD, Jonas BA, Pullarkat V, et al. Azacitidine and venetoclax in previously untreated acute myeloid leukemia. *N Engl J Med*. 2020;383(7):617-629.
- [4] Short NJ, Kantarjian H. Choosing between intensive and less intensive front-line treatment approaches for older patients with newly diagnosed acute myeloid leukaemia. *Lancet Haematol*. 2022;9(7):e535-e545.