

43. Andersen MK, Pedersen-Bjergaard J, Kjeldsen L, Dufva IH, Brondum-Nielsen K. Clonal Ph-negative hematopoiesis in CML after therapy with imatinib mesylate is frequently characterized by trisomy 8. *Leukemia* 2002;16:1390-3.
44. Cashman J, Henkelman D, Humphries K, Eaves C, Eaves A. Individual BFU-E in polycythemia vera produce both erythropoietin dependent and independent progeny. *Blood* 1983;61:876-84.
45. Levin J, Cocault L, Demerens C, Challier C, Pauchard M, Caen J, et al. Thrombocytopenic c-mpl(-/-) mice can produce a normal level of platelets after administration of 5-fluorouracil: the effect of age on the response. *Blood* 2001;98:1019-27.
46. Berlin NI, Louis R, Wasserman and the history of polycythemia vera. *Mt Sinai J Med* 1995;62:206-15.
47. Berk PD, Goldberg JD, Silverstein MN, Weinfeld A, Donovan PB, Ellis JT, et al. Increased incidence of acute leukemia in polycythemia vera associated with chlorambucil therapy. *N Engl J Med* 1981;304:441-7.
48. Najean Y, Deschamps A, Dresch C, Daniel MT, Rain JD, Arrago JP. Acute leukemia and myelodysplasia in polycythemia vera. A clinical study with long-term follow-up. *Cancer* 1988;61:89-95.
49. Silverstein MN. Postpolycythemia myeloid metaplasia. *Arch Intern Med* 1974;134:113-7.
50. Randi ML, Barbone E, Fabris F, Varotto L, Macri C, Girolami A. Post-polycythemia myeloid metaplasia: experience with a large cohort of patients. *J Med* 1994;25:363-9.
51. Streiff MB, Smith B, Spivak JL. The diagnosis and management of polycythemia vera in the era since the Polycythemia Vera Study Group: a survey of American Society of Hematology members' practice patterns. *Blood* 2002;99:1144-9.
52. Silver RT. A new treatment for polycythemia vera: recombinant interferon α . *Blood* 1990;76:664-5.
53. Silver RT. Imatinib mesylate reduces phlebotomy requirements in polycythemia vera. *Blood* 2002;100:41a[abstract].
54. Sterkers Y, Preudhomme C, Lai JL, Demory JL, Caulier MT, Watel E, et al. Acute myeloid leukemia and myelodysplastic syndromes following essential thrombocythemia treated with hydroxyurea: high proportion of cases with 17p deletion. *Blood* 1998;91:616-22.
55. Bernasconi P, Boni M, Cavigliano PM, Calatroni S, Brusamolino E, Passamonti F, et al. Acute myeloid leukemia (AML) having evolved from essential thrombocythemia (ET): distinctive chromosome abnormalities in patients treated with pipobroman or hydroxyurea. *Leukemia* 2002;16:2078-83.

FLT3 inhibition as tailored therapy for acute myeloid leukemia

The clinical success of the specific tyrosine kinase inhibitor, STI571^{1,2} (otherwise known as Glivec® or imatinib, Novartis Pharma), has fostered onco-hematologic research worldwide to develop new molecularly targeted forms of therapy. The target of imatinib is preferentially BCR-ABL, an intracellular oncogenic tyrosine kinase that shares several homologies with the class III receptor tyrosine kinase (RTK) family, whose members include the FLT3, KIT, FMS, and PDGF receptors.^{3,4} Most of these RTKs are implicated, either in mutated or wild-type conformations, in the constitutive activation and proliferation of human leukemias, especially acute myeloid leukemia (AML).^{3,5}

Particular interest has been aroused by the relatively high frequency of FLT3 receptor mutations found in AML. The FLT3 receptor has several structural domains, including 5 immunoglobulin-like domains in the extracellular regions, a juxtamembrane (JM) domain, 2 kinase domains (TK1 and TK2) separated by a kinase insert (KI) domain, and a C-ter-

minal domain in intracellular regions.⁴ Ligand binding to the RTK extracellular domain leads to receptor dimerization, stabilizes an open conformation of the catalytic domain (A-loop) for adenosine triphosphate (ATP) and substrate binding, and enables transphosphorylation of the A-loop. The subsequent phosphorylation of tyrosine residues accompanies RTK activation. Consequently, the FLT3 receptor leads to induction of fundamental intracellular signaling pathways, which in turn regulate both cell proliferation and apoptosis.³

The FLT3 receptor has been found to be frequently targeted in AML and somewhat less commonly in myelodysplastic syndromes (MDS) by two different types of genetic alteration.

Firstly, an internal tandem duplication (ITD) of the JM domain-coding sequence of the *FLT3* gene (FLT3/ITD) is found in 20% to 41% of adult and pediatric patients with *de novo* or secondary AML, as well as in about 3% of patients with MDS,⁶⁻¹⁰ as shown also by Moreno *et al.*¹¹ in this issue of *Haematologica*. These mutations constitutively activate the receptor, and are strongly associated with hyperleukocytosis, poor response to therapy and dismal prognosis. Among the distinctive forms of AML, higher frequencies of FLT3 alterations have been detected in acute promyelocytic leukemia (APL)¹² and in AMLs with apparently normal karyotype.⁶⁻¹¹ The reasons underlying these associations are currently unclear. When transplanted in murine hematopoietic progenitors, the mutant FLT3 receptor causes cellular transformation and produces a myeloproliferative syndrome, even though this does not by itself appear sufficient to cause acute leukemia.^{13,14} An additional length mutation affecting the tyrosine kinase domain in exon 20 has been recently described.¹⁵

Secondly, point mutations in the FLT3 receptor have been reported to occur in 3% to 8% of AML patients, mostly at a specific site in the gene (D835 and I836)¹⁶ (Figure 1). Although not apparently associated with either leukocytosis or worse prognosis, these point mutations result in similar deregulatory activity on the receptor and cause its constitutive activation and tend to worsen disease-free survival. Furthermore, the point mutations occur independently of FLT3/ITD. Taken together, these observations indicate that *FLT3* currently appears to be the most frequently mutated gene and constitutively activated receptor in AML.

The FLT3 receptor as a candidate target for tailored therapy in acute myeloid leukemia

Following the remarkable success of STI571, several researchers have pointed to the FLT3 receptor, or its signal transduction pathway,¹⁷ as a possible specific target for tailored therapy. Several tyrosine kinase inhibitors, which were not originally developed with FLT3 as the intended target, have been reported to inhibit the FLT3 receptor on AML cell lines or prima-




Figure 1. Schematic representation of the FLT3 receptor kinase domains with point mutations and some types of ITD. Schematic representation of the FLT3 receptor kinase domains: Genomic, mRNA and protein structure of the FLT3 receptor. Right side of the figure: DNA: schematic illustration of exons 10, 11, 12 and 17 of the FLT3 receptor gene. RNA, messenger RNA representation: extra cellular domain (ECD); transmembrane domain (TM); juxtamembrane domain (JM); tyrosine kinase domain (TK1); kinase insert (KI); tyrosine kinase domain2 (TK2); C terminal region (COOH). Protein: representation of protein domains of functional FLT3 receptor. The kinase domain is not drawn to scale. Middle of the figure: two types of activating mutations in FLT3 are associated with AML; the first type consists of ITDs of amino acids in the JM domain resulting in constitutive tyrosine kinase activation. WT, wild-type amino acid sequence of the region usually involved in ITDs;¹⁻⁶ some derivative ITDs sequences with the point of insertion (arrows). The second type of mutation consists of point mutations in the so-called *activation loop* of the second tyrosine kinase domain. Mutations at two specific residues, aspartic acid (D835) and/or isoleucine (I836), also result in constitutive FLT3 activation. Amino acid substitutions are depicted; arrows indicate the points of insertion of representative ITD sequences. Modified from: Gilliland DG, Griffin JD. The roles of FLT3 in hematopoiesis and leukemia. *Blood* 2002;100:1532-42.

ry blast cells. These include the tyrphostin AG1295¹⁸ or AG1296,¹⁹ and herbimycin A (HA).^{17,20,21}

Following on from their preliminary observations on the inhibitory activity of HA, Minani *et al.*²⁰ suggested that the mechanism behind the cytotoxicity of this ansamycin derivative, which is now known to target Hsp90, could be mediated by the inhibition of phosphorylation of ITD-FLT3. Exposure to another Hsp90 inhibiting HA, namely radicicol,²² was able to dissociate ITD-FLT3 from the Hsp90 chaperone complex, activating blast cell apoptosis. More recently, inhibition of FLT3 has been reported in pre-clinical studies of other RTK inhibitors, namely CEP-701 or KT-5555,²³⁻²⁵ PKC412,²⁶ and CT53518.²⁷ Moreover, in an interesting accompanying review that appeared in the same issue of *Cancer Cell*,²⁸ Sawyers reported a fourth inhibitor, SU11248,²⁹ that is now being evaluated in a clinical trial (in which we are directly involved). All four molecules have been designed to bind the receptor specifically and compete with ATP in its ATP-binding pocket. CEP-701 exerts an inhibitory action on the TrkA receptor tyrosine

kinase.²³ The inhibitory activity of PKC412 extends to RTK other than FLT3, such as protein kinase C, VEGFR2, PDGFR, c-Kit, and FMS.²⁶

Similarly, Millenium's compound, CT53518 and Sugen's SU11248 (the latter derived from the chemical research and development of SU5416)^{28,29} both have further inhibitory effects on PDGFR and c-kit.²⁷

With four FLT3 inhibitors heading into the clinic, is it possible to predict which patients stand to gain most benefit from these drugs?

Assuming that FLT3 is a suitable target for molecular therapy in AML, we can postulate that enrolment in clinical trials should be proposed for those patients with FLT3 receptor alterations in their leukemic cells. Furthermore, treatment could be extended to those AML patients refractory or resistant to conventional chemotherapy and/or with associated or concomitant mutated c-Kit and PDGFR translocation.³⁰⁻³³ Based on pre-clinical studies, we can expect that AML patients with FLT3 alterations may show some clinical response.

Are all AML patients likely to respond to FLT3 inhibitors?

In CML Ph⁺ patients in chronic phase, the molecular target of imatinib is ABL, present either in its P210 form (as in the BCR-ABL fusion protein) or in its wild-type form. Furthermore, the BCR-ABL oncogene is absent from normal stem cells but present in its double isoform (b2-a2 and b3-a2, with very rare exceptions)^{34,35} on the leukemic cells, offering a unique molecular target for imatinib's selectivity. On the other hand, FLT3 receptors carrying ITD exhibit more heterogeneous forms of mutation, giving rise to several individual structural forms of the receptor. It is likely that it will be difficult for the inhibitor drugs to recognize, bind and inhibit all these forms. Based on the different chemical formulations of the four inhibitors, and again learning from the ABL/imatinib interaction experience,³⁶ we can hypothesize that the different inhibitors are likely to show different combining activities with each form of the mutated FLT3 receptor. We may expect various degrees of clinical responsiveness among patients, simply based on the specific interactions of each inhibitor with variable forms of FLT3 ITD. We could also envisage peculiar, individual degrees of sensitivity to the same inhibitor among patients sharing the same molecular defect. In this respect, it is important to consider that, distinct from CML, AMLs comprise a spectrum of genetically heterogeneous diseases which may account for different sensitivities to the inhibitor.

Is it possible to predict resistance to these inhibitors? And could there be "intrinsic" resistance pre-dating the start of therapy?

It is now clear that resistance to imatinib is mostly due to point mutations in or around the ABL ATP binding pocket. Despite recent observations suggesting that *polyclonal* resistance to imatinib could be present prior to the initiation of therapy in a few CML patients,³⁷ in the majority of cases cellular resistance to imatinib is acquired during treatment. Multiple independent mutant clones seem to emerge during treatment with imatinib, along with the ABL point mutations that have been detected in relapsed cases,³⁷ and a clonal selection of these cells would confer refractoriness that could be defined as *acquired resistance*. On the other hand, the mutated FLT3 receptor form is present from the onset of disease in the majority of AML patients and ITD or point mutations are associated exclusively with the leukemic clone: either the ITD or the point mutation, or a combination of the two, may confer a proliferative advantage to the blast cell. If this is the case, resistance to FLT3 inhibitors could be present from the onset of disease (i.e. *intrinsic* resistance). Drug inhibitory pressure could then foster a clonal selection of cells with a second or further mutation, conferring further resistance (*acquired* resistance).

From the co-crystal structure of the ABL kinase

bound to imatinib^{36,38,39} it has been postulated that *acquired* drug resistance³⁵ depends on a reduced interaction between imatinib and the mutated ABL form, the latter being due to a single amino acid substitution. To overcome this obstacle, a mutated form of imatinib³⁶ and alternative inhibitors, such as PD173955,^{36,40} have been designed. PD173955 is able to bind the ABL ATP pocket in the so-called *on* conformation, which is supposed to be more accessible to the drug. At present, however, there are no data on the co-crystal structure of the FLT3 receptor either in its wild-type conformation or in its mutated form. Furthermore, no information is available about the co-crystal structure of the FLT3 receptor associated with any of the inhibitors. In the absence of information regarding the structure whereby the receptor binds to the inhibitors, one can hardly speculate on *intrinsic* resistance.

In conclusion, unlike CML, AML with FLT3 ITD or point mutations are characterized by multiple genetic forms. This variety of forms hampers prediction of success rates for the majority of patients treated with any of the four new FLT receptor inhibitors. Only clinical trials will reveal whether this treatment can provide a viable option for such patients. On the other hand, extensive molecular characterization of FLT3 gene status will be required to determine whether a given alteration type is more susceptible to targeted inhibition. Our personal preference would be to include relapsed AML patients carrying FLT3 receptor ITD and/or point mutations in initial trials. Combined with extensive and multiparametric genetic characterization of enrolled cases, this would favor rapid identification of responders. Mechanism(s) of resistance could then be investigated and hopefully identified in subsequent studies.

Giovanni Martinelli, Pier Paolo Piccaluga,
Francesco Lo Coco*
Institute of Hematology and Medical Oncology "L.
e A. Seràgnoli", University of Bologna;
*Department of Cellular Biotechnologies and
Hematology, "La Sapienza" University, Rome, Italy

Funding

This work and GM were supported by the Associazione Italiana per la Ricerca sul Cancro (A.I.R.C.); MURST 40%; MURST: Ateneo di Bologna 2001 and 2002; A.I.L.; and the Italian C.N.R target. We thank Bianchini Michele for technical assistance with drawing the picture.

References

1. Druker B. Signal transduction inhibition: results from phase I clinical trials in chronic myeloid leukemia. *Semin Hematol* 2001;38:9-14.
2. Kantarjian H, Sawyers C, Hochhaus A, Guilhot F, Schiffer C, Gambacorti-Passerini C, et al. Hematologic and cytogenetic responses to imatinib mesylate in chronic myelogenous leukemia. *N Engl J Med* 2002;346:645-52.

3. Gilliland DG, Griffin JD. The roles of FLT3 in hematopoiesis and leukemia. *Blood* 2002;100:1532-42.
4. Matthews W, Jordan CT, Wiegand GW, Pardoll D, Lemischka IR. A receptor tyrosine kinase specific to hematopoietic stem and progenitor cell-enriched populations. *Cell* 1991;65:1143-52.
5. Blume-Jensen P, Hunter T. Oncogenic kinase signalling. *Nature* 2001;411:355-65.
6. Kondo M, Horibe K, Takahashi Y, Matsumoto K, Fukuda M, Inaba J, et al. Prognostic value of internal tandem duplication of the FLT3 gene in childhood acute myelogenous leukemia. *Med Pediatr Oncol* 1999; 33:525-9.
7. Thiede C, Steudel C, Mohr B, Schaich M, Schakel U, Platzbecker U, et al. Analysis of FLT3-activating mutations in 979 patients with acute myelogenous leukemia: association with FAB subtypes and identification of subgroups with poor prognosis. *Blood* 2002;99:4326-35.
8. Liang DC, Shih LY, Hung IJ, Yang CP, Chen SH, Jaing TH, et al. Clinical relevance of internal tandem duplication of the FLT3 gene in childhood acute myeloid leukemia. *Cancer* 2002;94:3292-8.
9. Meshinchi S, Woods WG, Stirewalt DL, Sweetser DA, Buckley JD, Tjoa TK, et al. Prevalence and prognostic significance of Flt3 internal tandem duplication in pediatric acute myeloid leukemia. *Blood* 2001;97:89-94.
10. Schnittger S, Schoch C, Dugas M, Kern W, Staib P, Wuchter C, et al. Analysis of FLT3 length mutations in 1003 patients with acute myeloid leukemia: correlation to cytogenetics, FAB subtype, and prognosis in the AMLCG study and usefulness as a marker for the detection of minimal residual disease. *Blood* 2002;100:59-66.
11. Moreno I, Barragan E, Rueda E, Roman J, Fernandez P, Leon P, et al. Incidence and prognostic value of FLT3 internal tandem duplication and D835 mutations in acute myeloid leukemia. *Haematologica* 2003;88:21-6.
12. Noguera NI, Breccia M, Divona M, Diverio D, Costa V, De Santis S, et al. Alterations of the FLT3 gene in acute promyelocytic leukemia: association with diagnostic characteristics and analysis of clinical outcome in patients treated with the Italian AIDA protocol. *Leukemia* 2002;16:2185-9.
13. Kelly LM, Kutok JL, Williams IR, Boulton CL, Amaral SM, Curley DP, et al. PML/RAR α and FLT3-ITD induce an APL-like disease in a mouse model. *Proc Natl Acad Sci USA* 2002;99:8283-8.
14. Kelly LM, Liu Q, Kutok JL, Williams IR, Boulton CL, Gilliland DG. FLT3 internal tandem duplication mutations associated with human acute myeloid leukemias induce myeloproliferative disease in a murine bone marrow transplant model. *Blood* 2002;99:310-8.
15. Spiekermann K, Schoch C, Haferlach T, Hiddemann W, Schnittger S. A new and recurrent activating length mutation in exon 20 of the FLT3 gene in acute myeloid leukemia. *Blood* 2002;100:3423-5.
16. Yamamoto Y, Kiyoi H, Nakano Y, Suzuki R, Kodera Y, Miyawaki S, et al. Activating mutation of D835 within the activation loop of FLT3 in human hematologic malignancies. *Blood* 2001; 97:2434-9.
17. Naoe T, Kiyoe H, Yamamoto Y, Minami Y, Yamamoto K, Ueda R, et al. FLT3 tyrosine kinase as a target molecule for selective antileukemia therapy. *Cancer Chemother Pharmacol* 2001;48 Suppl 1:S27-30.
18. Levis M, Tse KF, Smith BD, Garrett E, Small D. A FLT3 tyrosine kinase inhibitor is selectively cytotoxic to acute myeloid leukemia blasts harboring FLT3 internal tandem duplication mutations. *Blood* 2001;98:885-7.
19. Tse KF, Novelli E, Civin CI, Bohmer FD, Small D. Inhibition of FLT3-mediated transformation by use of a tyrosine kinase inhibitor. *Leukemia* 2001;15:1001-10.
20. Minami Y, Kiyoi H, Yamamoto Y, Yamamoto K, Ueda R, Saito H, et al. Selective apoptosis of tandemly duplicated FLT3-transformed leukemia cells by Hsp90 inhibitors. *Leukemia* 2002;16: 1535-40.
21. Zhao M, Kiyoi H, Yamamoto Y, Ito M, Towatari M, Omura S, et al. In vivo treatment of mutant FLT3-transformed murine leukemia with a tyrosine kinase inhibitor. *Leukemia* 2000; 14:374-8.
22. Kurebayashi J, Otsuki T, Kurosumi M, Soga S, Akinaga S, Sonoo H. A radicicol derivative, KF58333, inhibits expression of hypoxia-inducible factor-1 α and vascular endothelial growth factor, angiogenesis and growth of human breast cancer xenografts. *Jpn J Cancer Res* 2001;92:1342-51.
23. Levis M, Allebach J, Tse KF, Zheng R, Baldwin BR, Smith BD, et al. A FLT3-targeted tyrosine kinase inhibitor is cytotoxic to leukemia cells in vitro and in vivo. *Blood* 2002;99:3885-91.
24. George DJ, Dionne CA, Jani J, Angeles T, Murakata C, Lamb J, et al. Sustained in vivo regression of Dunning H rat prostate cancers treated with combinations of androgen ablation and Trk tyrosine kinase inhibitors, CEP-751 (KT-6587) or CEP-701 (KT-5555). *Cancer Res* 1999;59:2395-401.
25. Dionne CA, Camoratto AM, Jani JP, Emerson E, Neff N, Vaught JL, et al. Cell cycle-independent death of prostate adenocarcinoma is induced by the trk tyrosine kinase inhibitor CEP-751 (KT6587). *Clin Cancer Res* 1998;4:1887-98.
26. Weisberg E, Boulton C, Kelly LM, Manley P, Fabbro D, Meyer T, et al. Inhibition of mutant FLT3 receptors in leukemia cells by the small molecule tyrosine kinase inhibitor PKC412. *Cancer Cell* 2002;1:433-43.
27. Kelly LM, Yu JC, Boulton CL, Apatira M, Li J, Sullivan CM, et al. CT53518, a novel selective FLT3 antagonist for the treatment of acute myelogenous leukemia (AML). *Cancer Cell* 2002;1:421-32.
28. Sawyers CL. Finding the next Gleevec: FLT3 targeted kinase inhibitor therapy for acute myeloid leukemia. *Cancer Cell* 2002;1:413-5.
29. O'Farrell AM, Abrams T, Yuen H, Ngai T, Louie S, Wong L, et al. SUGEN compounds SU5416 and SU11248 inhibit Flt3 activity: therapeutic application in AML. American Society of Hematology 43rd Annual Meeting, Orlando, Florida 2001. abstract #497.
30. Golub TR, Barker GF, Lovett M, Gilliland DG. Fusion of PDGF receptor β to a novel ets-like gene, tel, in chronic myelomonocytic leukemia with t(5;12) chromosomal translocation. *Cell* 1994;77:307-16.
31. Apperley JF, Gardembas M, Melo JV, Russell-Jones R, Bain BJ, Baxter EJ, et al. Response to imatinib mesylate in patients with chronic myeloproliferative diseases with rearrangements of the platelet-derived growth factor receptor β . *N Engl J Med* 2002; 347:481-7.
32. Baxter EJ, Hochhaus A, Bolufer P, Reiter A, Fernandez JM, Senent L, et al. The t(4;22)(q12;q11) in atypical chronic myeloid leukaemia fuses BCR to PDGFRA. *Hum Mol Genet* 2002;11: 1391-7.
33. Baxter EJ, Kulkarni S, Vizmanos JL, Jaju R, Martinelli G, Testoni N, et al. Novel translocations that disrupt the platelet-derived growth factor receptor B (PDGFRB) gene in BCR-ABL negative myeloproliferative disorders. *Br J Haematol* 2002;in press.
34. Martinelli G, Amabile M, Giannini B, Terragna C, Ottaviani E, Soverini S, et al. Novel types of bcr-abl transcript with breakpoints in BCR exon 8 found in Philadelphia positive patients with typical chronic myeloid leukemia retain the sequence encoding for the DBL- and CDC24 homology domains but not the pleckstrin homology one. *Haematologica* 2002;87:688-94.
35. Roumiantsev S, Shah NP, Gorre ME, Nicoll J, Brasher BB, Sawyers CL, et al. Clinical resistance to the kinase inhibitor STI-571 in chronic myeloid leukemia by mutation of Tyr-253 in the Abl kinase domain P-loop. *Proc Natl Acad Sci USA* 2002;99: 10700-5.
36. Nagar B, Bornmann WG, Pellicena P, Schindler T, Veach DR, Miller WT, et al. Crystal structures of the kinase domain of c-Abl in complex with the small molecule inhibitors PD173955 and imatinib (STI-571). *Cancer Res* 2002;62:4236-43.
37. Shah NP, Nicoll JM, Nagar B, Gorre ME, Paquette RL, Kuriyan J, et al. Multiple BCR-ABL kinase domain mutations confer polyclonal resistance to the tyrosine kinase inhibitor imatinib (STI571) in chronic phase and blast crisis chronic myeloid leukemia. *Cancer Cell* 2002;2:117-25.
38. Marx J. Molecular biology. Cancer fighter's modus operandi revealed. *Science* 2000;289:1857-9.
39. Schindler T, Bornmann W, Pellicena P, Miller WT, Clarkson B, Kuriyan J. Structural mechanism for STI-571 inhibition of abelson tyrosine kinase. *Science* 2000;289:1938-42.
40. Wisniewski D, Lambek CL, Liu C, Strife A, Veach DR, Nagar B, et al. Characterization of potent inhibitors of the Bcr-Abl and the c-kit receptor tyrosine kinases. *Cancer Res* 2002;62:4244-55.