

Peripheral blood or bone marrow cells in reduced-intensity or myeloablative conditioning allogeneic HLA identical sibling donor transplantation for multiple myeloma

Gösta Gahrton, Simona Iacobelli, Giuseppe Bandini, Bo Björkstrand, Paolo Corradini, Charles Crawley, Ute Hegenbart, Gareth Morgan, Nicolaus Kröger, Anton Schattenberg, Stefan O. Schonland, Leo F. Verdonck, Lisa Volin, Theo de Witte, Dietger Niederwieser for the Myeloma Subcommittee of the EBMT

Addendum

List of centers and reporters that have contributed to this study:

F. Bonifazi, Hospital San Orsola, Bologna, Italy [240]; L. Verdonck, University Medical Centre Utrecht, Utrecht, The Netherlands [239]; T. Ruutu, Helsinki University Central Hospital, Helsinki, Finland [515]; A. Schattenberg, University Medical Center St. Radboud, Nijmegen, The Netherlands [237]; P. Dreger, University of Heidelberg, Heidelberg, Germany [524]; G. Morgan, Royal Marsden Hospital, Sutton, UK [218]; A. Goldstone, University College London Hospital, London, United Kingdom [224]; N.H. Russell, Nottingham City Hospital, Nottingham, UK [717]; J. Sierra, Hospital Santa Creu I Sant Pau, Barcelona, Spain [260]; A. Zander, University Hospital Eppendorf, Hamburg, Germany [614]; J. Cornelissen, D.den Hoed Cancer Centre/AZR, Rotterdam, The Netherlands [246]; P. von dem Borne, Leiden University Hospital, Leiden, The Netherlands [203]; M. Boccadoro, Azienda Ospedaliera S. Giovanni Batista, Torino, Italy [696]; N. Milpied, Hopital Haut-Leveque, Pessac, France [267]; J.L. Harousseau, Hotel Dieu, Nantes, France [253]; J. Apperley, Hammersmith Hospital, London, UK [205]; M. Michallet, Hopital E. Herriot, Lyon, France [671]; A. Bacigalupo, Ospedale San Martino, Genova, Italy [217]; D. Bunjes, Universität Ulm, Ulm, Germany [204]; R. Scime, Ospedale V. Cervello, Palermo, Italy [392]; A. Torres Gomez, Cordoba Hospital – Reina Sofia, Córdoba, Spain [238]; A. Ferrant, Cliniques Universitaires St. Luc, Brussels, Belgium [234]; D. Blaise, Institut Paoli Calmettes, Marseille, France [230]; P. Jacobs, Constantiaberg Med-Clin, Cape Town, South Africa [772]; S. Amadori, Univ. Tor Vergata, St. Eugenio Hospital, Rome, Italy [756]; A. Bosi, Ospedale di Careggi, Firenze, Italy [304]; I. Franklin, Glasgow Royal Infirmary, Glasgow, Scotland, UK [244]; J.Y. Cahn, Hopital A. Michallon, Grenoble, France [270]; D. Caballero, Hospital Clinico, Salamanca, Spain [727]; A.M. Carella, IRCCS, Casa Sollievo della Sofferenza, San Giovanni Rotondo, Italy [526]; L. Ahlberg, University Hospital Linköping, Linköping, Sweden [740]; I. Majolino, Ospedale S. Camillo, Rome, Italy [287]; L. Jebavy, Charles University Hospital, Hradec Králové, Czech Republic [729]; R. Hamladji, Centre Pierre et Marie Curie, Alger, Algeria [703]; E. Carreras, Hospital Clinic, Barcelona, Spain [214]; A. Gratwohl, Kantonsspital, Basel, Switzerland [202]; E. Liakopoulou, Christie NHS Trust Hospital, Manchester, United Kingdom [780]; P. Di Bartolomeo, Ospedale Civile, Pescara, Italy [248]; M. Falda, Azienda Ospedaliera S. Giovanni, Torino, Italy [231]; L. Kanz, Medizinische Klinik, Tübingen, Germany [223]; J. Finke, University Hospital Freiburg, Freiburg, Germany [810]; C. Craddock, Univ. Hosp. Birmingham NHST Trust, Birmingham, UK [387]; R. Haas, Heinrich Heine Universität, Düsseldorf, Germany [390]; A. Iriondo, Hospital Universitario 'Marqués de Valdecilla',

Santander, Spain [242]; M. Boogaerts, University Hospital of Leuven, Leuven, Belgium [209]; K. Remes, Turku University Central Hospital, Turku, Finland [225]; G. Lambertenghi Deliliers, Ospedale Maggiore di Milano, Milano, Italy [265]; G. Meloni, Univ. "La Sapienza", Rome, Italy [232]; T. Littlewood, The Oxford Radcliffe Hospital, Oxford, United Kingdom [255]; M. L.A. Noens, University Hospital Gent, Gent, Belgium [744]; C.A. de Souza, Univ. Est. de Campinas/TMO/UNICAMP, Campinas SP, Brazil [374]; P. Iacopino, Azienda Ospedaliera, Reggio Calabria, Italy [587]; H. Schouten, University Hospital Maastricht, Maastricht, The Netherlands [565]; M. Bregni, Istituto Scientifico H.S. Raffaele, Milano, Italy [813]; H. Greinix, AKH Vienna, Vienna, Austria [227]; J. van der Lelie, Academic Medical Centre, Amsterdam, The Netherlands [247]; J. Hansz, K. Marcinkowski University of Medical Science, Poznan, Poland [730]; T. Barbui, Ospedale Bergamo, Bergamo, Italy [658]; J. Gribben, St. Batholomew's and the Royal London Hospital, London, UK [768]; E. Morra, Ospedale di Niguarda Ca'Granda, Milano, Italy, [294]; M. Fernandez, Clinica Puerta de Hierro, Madrid, Spain [728]; P. Ljungman, Huddinge University Hospital, Huddinge, Sweden [212]; D. Niederwieser, University of Leipzig, Leipzig, Germany [389]; B. Bradley, University of Bristol / Dept. of Transplantation Sciences, Bristol, UK [386]; A. Fassas, G.P. Gen. Hosp. of Thessaloniki, Exokhi, Greece [561]; J.H. Bourhis, Institut Gustave Roussy, Villejuif, France [666]; G.J. Ossenkoppele, VU University Medical Center, Amsterdam, The Netherlands [588]; M. Aglietta, Cattedra di Oncologia Medica, Candiolo Torino, Italy [377]; S. Slavin, Hadassah University Hospital, Jerusalem, Israel [258]; V. Leblond, Pitie-Salpetriere, Paris, France [262]; H.-J. Kolb, Klinikum Grosshadern, München, Germany [513]; J. Snowden, Royal Hallamshire Hospital, Sheffield, United Kingdom [778]; M. Sanz, Hospital Universitario La Fe, Valencia, Spain [663]; G. Gurman, Ankara University, Ibn Sina Hospital, Ankara, Turkey [617]; S. Mackinnon, Royal Free Hospital and School of Med., London, UK [216]; M. Attal, Hopital de Purpan, Toulouse, France [624]; C. Crawley, Addenbrookes Hospital, Cambridge, United Kingdom [566]; J.A. Lui Yin, Manchester Royal Infirmary, Manchester, United Kingdom [601]; M. Petrini, Dipart. di Oncol., dei trapianti e delle nuove tecnol. in med., Pisa, Italy [132]; G. Muftic, GKT school of Medicine London, United Kingdom [763]; A. Buzyn, Hôpital Necker, Paris, France [160]; E. Gluckman, Hopital St. Louis, Paris, France [207]; N. Patton, Canterbury Health Laboratory, Christchurch, New Zealand [798]; G. Ehninger, Universitätsklinikum Dresden, Dresden, Germany [808]; K. Kolbe, Johannes-Gutenberg-University, Mainz, Germany [786]; G. Visani, Pesaro Hospital, Pesaro, Italy [529]; R. Giustolisi, University of Catania, Catania, Italy [792]; D. Bron, Institut Jules Bordet, Brussels, Belgium [215]; N. Jacobsen, Rigshospitalet, Copenhagen, Denmark [206]; J. Passweg, Hopital Cantonal Universitaire, Geneva, Switzerland [261]; C. Cordonnier, Hopital Henri Mondor, Creteil, France [252]; P. Zachee, AZ

Stuivenberg, Antwerp, Belgium, [339]; T. Masszi, St. László Hospital, Hungary [739]; A. Schwarzer, Alfred Hospital, Melbourne, Australia [595]; R. Fanin, Udine University Hospital, Udine, Italy [705]; J. Fernandez-Ranada, Hospital de la Princesa, Madrid, Spain [236]; B. Lioure, Hopital de Hauteperre, Strasbourg, France [672]; J.P. Jouet, Hopital Claude Huriez, Lille, France [277]; J.M. Davies, Western General Hospital, Edinburgh, Scotland, UK [228]; F. Benedetti, University of Verona, Verona, Italy [623]; E. Deconinck, Hopital Jean Minjot, Besancon, France [233]; D. Selleslag, A.Z. Sint-Jan, Brugge, Belgium [506]; B. Rio, Hotel Dieu, Paris, France [222]; G. Karianakis, Diagnostic and Therapeutic Centre of Athens, Athens, Greece [643]; D. Beelen, University Hospital, Essen, Germany [259]; L. Contu, Univ. di Cagliari, Cagliari, Italy [811]; H. Horst, Christian-Albrechts-University, Kiel, Germany [256]; M.M. Martelli, University of Perugia, Perugia, Italy [794]; D. Milligan, Birmingham Heartlands Hospital, Birmingham, UK [284]; J. Sanchez de Toledo Codina, Hospital Vall d'Hebron, Barcelona, Spain [527]; S. Schey, Guy's Hospital, London, United Kingdom [721]; P. Coser, Hospital San Maurizio, Bolzano, Italy [299]; S. Kalayoglu-Besisik, University of Istanbul, Istanbul, Turkey [760]; P. Mazza, Hospedale Nord, Taranto, Italy [332]; G. Gastl, University Hospital Innsbruck, Austria [271]; M. Schapira, Universitaire Vaudois Centre Hospitalier, Lausanne, Switzerland [820]; F. Guilhot, Hopital La Miletrie, Poitiers, France [264]; H. Tilly, Centre Henri Becquerel, Rouen, France [941]; J. Rowe, Rambam Medical Centre, Haifa, Israel [345]; F. Rodeghiero, S. Bortolo Hospital, Vicenza, Italy [797]; B. Simonsson, University Hospital, Uppsala, Sweden [266]; E. Alessandrino, Policlinico San Matteo IRCCS, Pavia, Italy [286]; L. Garderet, Hopital Saint Antoine, Paris, France [213]; A. Leon Lara, Hospital del SAS, Cádiz Jerez de la Frontera, Spain [612]; J.M. Morelada Jimenez, Hospital Morales Meseguer, Murcia, Spain [735]; P. Pimentel, Inst. Portugues Oncologia Centro do Porto, Porto, Portugal [291]; M. Brune, Sahlgrenska University Hospital, Goeteborg, Sweden [289]; R. Duarte Palomino, L'Hospital Universitari de Bellvitge, Barcelona, Spain [759]; R. Schots, University Hospital VUB, Brussels, Belgium [630]; A.E. Hunter, Leicester Royal Infirmary NHS Trust, Leicester, United Kingdom [713]; J. M. de Pablos Gallego, Hospital Univ. Virgen de las Nieves, Granada, Spain [559]; N. Gratecos, Hôpital de l'Arget, Nice, France [523]; V. Koza, Charles University Hospital, Pilsen, Czech Republic [718]; J. Marsh, St. George's Hospital, London, United Kingdom [539]; P. Leoni, Clinica di Ematologia, Ospedale di Torrette, Ancona-Torrette, Italy [788]; J. Garcia-Conde, Hospital Clinico Universitario, Valencia, Spain [282]; S. Proctor, Royal Victoria Infirmary, Newcastle-upon-Tyne, United Kingdom [276]; E. Kansu, Hacettepe University, Ankara, Turkey [292]; R. Schwerdtfeger, Deutsche Klinik für Diagnostik, Wiesbaden, Germany [311]; W. Linkesch, Karl Franzens University Graz, Graz, Austria [308]; A. Ganser, Medical School of Hannover, Hannover, Germany [295]; G. Bries, Virga Jesse Hospital, Hasselt, Belgium [632]; J.L. Diez-Martin, Hospital G.U. Gregorio Maranon, Madrid, Spain [819]; O. de Weerd, St. Antonius Hospital, Nieuwegein, The Netherlands [200]; W. Rösler, University of Erlangen, Erlangen, Germany [809]; R. Arnold, Charite, Campus Mitte Med. Klinik, Berlin, Germany [807]; A. Nagler, Tel-Aviv University Chaim Sheba Medical Center, Tel-Hashomer, Israel [754]; A. Wahlin, Umea University Hospital, Umea, Sweden [731]; V. Hofmann, Onkozentrum Klinik Im Parke, Zürich, Switzerland [700]; S. Lenhoff, University Hospital, Lund, Sweden [283]; W. Wiktor-Jedrzejczak, The Medical University of Warsaw, Warsaw, Poland [954]; H. Wandt, Klinikum Nürnberg, Nürnberg, Germany [625]; N. Harhalakis, Evangelismos Hospital, Athens, Greece [622]; A. Gallamini, Az. Ospedaliera S. Croce e Carle, Cuneo, Italy [606]; G. Cooke, St James' University Hospital, Leeds, United Kingdom [254]; N. Novitzky, UCT Medical School, Cape Town, South Africa [512]; R.E. Clark, Royal Liverpool University Hospital, Liverpool, United Kingdom [501]; L. Brinch, Rikshospitalet, The National Hospital, Oslo, Norway [235]; F.G.C. Jones, Belfast City hospital, Belfast Northern Ireland, United Kingdom [268]; J. Bay, Centre Jean Perrin, Clermont-Ferrand, France [273]; V. Rizzoli, Univ. di Parma, Parma, Italy, [245]; J.C. Garcia-Ruiz, Hospital de Cruces, Barakaldo (Bilbao) (Vizcaya), Spain [393]; A. Fauser, Klinik für K.M.T. & Hämato-Onkologie Idar-Oberstein, Germany [592]; S. Killick, Royal Bournemouth Hospital, Bournemouth, United Kingdom [765]; S. Cagiran, Ege University Medical School, Bornova-Izmir, Turkey [628]; J. Pretnar, University Med.Center, Ljubljana, Slovenia [640]; D. Carrera Fernandez, Hospital Covadonga, Oviedo, Spain [642]; A. Vitek, Inst. of Hematology and Blood Transf., Prague, Czech Republic [656]; G. Rossi, Spedali Civili – Brescia, Brescia, Italy [288]; D. Lutz, Elisabethinen-Hospital, Linz, Austria [594]; A. Lange, Lower Silesian Centre for Cellular Transplantation, Wroclaw, Poland [538]; A. Yalcin, Gulhane Military Medical Academy, Ankara, Turkey [372]; M. Koenigsmann, Otto-von-Guericke-Universität, Magdeburg, Germany [359]; J. Besalduch, Hospital Universitari Son Dureta, Palma de Mallorca, Spain [722]; E. Iannitto, Univ. di Palermo, Palermo, Italy [814]; K. Sulek, Military Medical Academy, Warsaw, Poland [816]; J.F. Tomas, Fundación Jimenez Díaz, Madrid, Spain [309]; K. Wilson, University of Wales College of Medicine, Cardiff Wales, UK [303]; W. Siegert, Charite-Virchow Klinikum d.Humboldt-Univ., Berlin, Germany [293]; E. Holler, University Regensburg, Regensburg, Germany [787]; A. Martinez-Rubio, Hospital Infantil La Paz, Madrid, Spain [734]; U. Tidefelt, Örebro University Hospital, Örebro, Sweden [738]; M. Bayik, Marmara Universitesi Hastanesi, Istanbul, Turkey [714]; S. Morandi, Centro Trapianti Midollo Osseo, Medicina II, Cremona, Italy [226]; R.P. Herrmann, Royal Perth Hospital, Perth Western Australia, Australia [710]; L. Feldman, Antartida Hospital Privado, Buenos Aires, Argentina [742]; Y. Beguin, University of Liege, Liege, Belgium [726]; B. Metzner, Klinikum Oldenburg, Oldenburg, Germany [749]; A. Zambelli, Fondazione S. Maugeri, Pavia, Italy [771]; B. Marianska, Institute of Haematology and Blood Transfusion, Warsaw, Poland [693]; W. Heit, Ev. Krankenhaus, Essen, Germany [784]; U. Schanz, University Hospital, Zürich, Switzerland [208]; A. Gabbas, Ospedale 'San Francesco', Nuoro, Italy [793]; C. Favre, University of Pisa, Pisa, Italy [795]; A. Hellmann, Medical University of Gdansk, Gdansk, Poland [799]; A. Uss, Hospital No.9, Minsk, Belarus [801]; A. Alabdulaaly, Riyadh Armed Forces Hospital, Riyadh, Saudi Arabia [818]; Dr. A. M. Peny, Francois Baclesse, Caen, France [951]; M. Sandherr, Klinikum Augsburg II Medizinische Klinik, Augsburg, Germany [152]; R. de Bock, AZ Middelheim, Antwerp, Belgium [783]; O. Reman, Centre Hospitalier Universitaire, Caen, France [251]; B. Labar, University Hospital Centre – Rebro, Zagreb, Croatia [302]; G. Leone, Università Cattolica S. Cuore, Rome, Italy [307]; M. Bargetzi, Kantonsspital Aarau, Aarau, Switzerland [316]; G. Marotta, Policlinico Le Scotte, Siena, Italy [321]; T. Fitzsimons, University of Glasgow, Glasgow Scotland, United Kingdom [325]; A. Santoro, Transplantation Unit, Milano Rozzano, Italy [354]; G.W. van Imhoff, Univ. Medical Center Groningen, Groningen, The Netherlands [546]; M. van Marwijk Kooy, Isala Kliniecken Zwolle, Zwolle, The Netherlands [548]; C. von Schilling, Klinikum Rechts der Isar III Med Klinik der TU, Munich, Germany [558] 1; O.

Vinante, Civiv Hospital, Noale (VE), Italy [563]; L. Gugliotta, Arcispedale S. Maria Nuova, Reggio Emilia, Italy [660]; K.R. Romeril, Wellington Hospital, Wellington, New Zealand [581]; J. Kienast, University of Münster, Münster, Germany [680]; P. Lederlin, CHRU Nancy, Nancy, France [249]; M. Freund, Universität Rostock, Rostock, Germany [585]; J. Vorlicek, University Hospital Brno, Brno, Czech Republic [597]; A. Neubauer, Philipps Universitaet Marburg, Marburg, Germany [645]; V. Liso, Universita degli Studi di Bari, Bari, Italy [649]; M. Boasson, CHRU, Angers, France [650]; E. Pogliani, Ospedale San Gerardo, Monza, Italy [544]; E. le Gall, CHRU de Rennes-Hôpital Sud Clinique Med. Infantile, Rennes, France [661]; T. de Revel, Hôpital Percy, Clamart, France [665]; J. Holowiecki, Silesian Medical Academy, Katowice, Poland [677]; J.L. Gil, Hospital Universitario 'Puerta del Mar', Cádiz, Spain [679]; M. Gonzalez Valentin, Hospital Carlos Haya, Malaga, Spain [576].