


Functional characterization of twelve natural *PROS1* mutations associated with anticoagulant protein S deficiency

Begoña Hurtado,^{1,5} Xavier Muñoz,^{1,5} M^a Carme Mulero,¹ Gemma Navarro,² Pere Domènech,³ Pablo García de Frutos,⁴ Mercè Pérez-Riba¹ and Núria Sala¹

From the Centre de Genètica Mèdica i Molecular, IDIBELL, L'Hospitalet de Llobregat, Barcelona, Spain;¹ Institut Municipal d'Assistència Sanitària (IMAS), Hospital del Mar, Barcelona (GN);² Unitat de Trombosi i Hemostàsia, Hospital Universitari de Bellvitge, L'Hospitalet de Llobregat, Barcelona;³ Department of Neurological and Vascular Research, Institute for Biomedical Research of Barcelona, IIBB-CSIC-IDIBAPS, Barcelona⁴ Present address: Laboratori de Recerca Translacional, Institut Català d'Oncologia (ICO)-IDIBELL, L'Hospitalet de Llobregat, Barcelona⁵

Citation: Hurtado B, Muñoz X, Mulero C, Navarro G, Domènech P, de Frutos PG, Pérez-Riba M and Sala N. Functional characterization of 12 natural *PROS1* mutations associated with anticoagulant protein S deficiency *Haematologica* 2008 Apr; 93(4):XXX-XXX

DOI:10.3324/haematol.12090



Supplementary Figure S1. Standard curve for PS measurements with the total and free PS ELISA assay. 150, 100, 50, 12.5 and 6.25 ng of purified human plasma PS (hPSc) in phosphate-buffered saline were measured with each assay.