



# Transcriptional activation of the cardiac homeobox gene CSX1/NKX2-5 in a B-cell chronic lymphoproliferative disorder

Xinying Su,<sup>1,5</sup> Véronique Della-Valle,<sup>1,5</sup> Eric Delabesse,<sup>3</sup> Zahia Azgui,<sup>2</sup> Roland Berger,<sup>1</sup> Hélène Merle-Béral,<sup>2,4</sup> Olivier A Bernard,<sup>1,5</sup> and Florence Nguyen-Khac<sup>1,2,4</sup>

<sup>1</sup>INSERM, E0210, Hôpital Necker, Paris;

<sup>2</sup>Service d'Hématologie Biologique, Groupe Hospitalier Pitie-Salpetriere, Paris; <sup>3</sup>Laboratoire d'Hématologie, CHU Purpan, Toulouse; <sup>4</sup>Université Pierre et Marie Curie, Paris VI; <sup>5</sup>Université Rene Descartes, Paris V, France

Citation: Su CX, Della-Valle V, Delabesse E, Azgui Z, Berger R, Merle-Béral H, Bernard OA and Nguyen-Khac F. Transcriptional activation of the cardiac homeobox gene CSX1/NKX2-5 in a B-cell chronic lymphoproliferative disorder. *Haematologica* 2008; doi: 10.3324/haematol.12595



| %GFP+      | round I | round II |
|------------|---------|----------|
| MIE        | 84      | 0        |
| HOXA9      | 75      | 84,52    |
| NKX2-5     | 84,35   | 84,41    |
| HOX11/TLX1 | 35,86   | 92,15    |

**Supplementary Figure S2.** Immunophenotype and GFP fluorescence analyses of cells present at round.