


**The MDM2 -309 T/G promoter single nucleotide polymorphism does not alter disease characteristics in chronic lymphocytic leukemia (CLL)**

Thorsten Zenz,<sup>1</sup> Sonja Häbe,<sup>1</sup> Axel Benner,<sup>2</sup> Dirk Kienle,<sup>1</sup> Hartmut Döhner<sup>1</sup> and Stephan Stilgenbauer<sup>1</sup>


<sup>1</sup>Department of Internal Medicine III, University of Ulm, Ulm, Germany; <sup>2</sup>Division of Biostatistics, German Cancer Research Center, Heidelberg, Germany

Citation: Zenz T, Häbe S, Benner A, Kienle D, Döhner H, Stilgenbauer S. The MDM2 -309 T/G promoter single nucleotide polymorphism does not alter disease characteristics in chronic lymphocytic leukemia. *Haematologica* 2008 doi: 10.3324/haematol.12738




**Supplementary Figure S1.** DHPLC pattern of single (A) and multiple (normalized graph) (B) samples with different MDM2 309 SNP sequences. The lower part of the figure (C) shows the respective sequences.

**A**



**B**



**Supplementary Figure S2. (A)** Quantitative gene expression and MDM2-SNP309: mRNA levels of MDM2 and P53 stratified by genotype. **(B)** MDM2-SNP309 has no influence on overall survival in CLL: Kaplan-Meier blot for overall survival in a cohort of patients with CLL ( $p=0.96$ ).