


Blocking the APRIL circuit enhances acute myeloid leukemia cell chemosensitivity

Désirée Bonci,¹ Maria Musumeci,¹ Valeria Coppola,¹ Antonio Addario,¹ Concetta Conticello,² Michael Hahne,³ Massimo Gulisano,⁴ Francesco Grignani,⁵ and Ruggero De Maria^{1,2}

¹Dept. of Hematology, Oncology and Molecular Medicine, Istituto Superiore Sanità, Rome, Italy; ²Mediterranean Institute of Oncology, Catania, Italy; ³Institut de Génétique Moléculaire de Montpellier, France; ⁴IOM Ricerca, Catania, Italy; ⁵Patologia Generale, Dipartimento di Medicina Clinica e Sperimentale, Perugia University, Policlinico Monteluce, Perugia, Italy

Citation: Bonci D, Musumeci M, Coppola V, Addario A, Conticello C, Hahne M, Gulisano M, Grignani F, and De Maria R. Blocking the APRIL circuit enhances acute myeloid leukemia cell chemosensitivity. *Haematologica* 2008. doi: 10.3324/haematol.13035



Online Supplementary Figure S1. APRIL expression and targeting in AML cells. (A) Flow cytometry analysis of AML blasts of M3 and M5 subtypes labeled with anti-APRIL antibody as compared to isotype control staining. One sample of two for each AML subtype is shown. (B) M2 blasts were treated with 5 μ g TACI-Fc and 5 μ g BCMA-Fc (BCMA/TACI-Fc) or 10 μ g Fn14-Fc recombinant soluble receptors and stained after 24h with annexin-V/7AAD. Data are mean \pm s.d. of three independent experiments in triplicate.