


A high number of losses in 13q14 chromosome is associated with a worse outcome and biological differences in patients with B chronic lymphoid leukemia (B-CLL)

José-Ángel Hernández,¹ Ana-Eugenia Rodríguez,² Marcos González,³ Rocío Benito,² Celia Fontanillo,² Virgilio Sandoval,⁴ Mercedes Romero,⁵ Guillermo Martín-Núñez,⁶ Alfonso García de Coca,⁷ Rosa Fisac,⁸ Josefina Galende,⁹ Isabel Recio,¹⁰ Francisco Ortuño,¹¹ Juan-Luis García,² Javier de las Rivas,² Norma-Carmen Gutiérrez,^{2,3} Jesús-Fernando San Miguel,^{2,3} and Jesús-María Hernández^{2,3}

¹Servicios de Hematología, Hospital Infanta Leonor, Madrid; ²IBMCC, Centro de Investigación del Cáncer, Universidad de Salamanca-CSIC; ³Hospital Clínico Universitario de Salamanca; ⁴Hospital Virgen Blanca, León; ⁵Hospital del Río Hortega, Valladolid; ⁶Hospital Virgen del Puerto, Plasencia, Cáceres; ⁷Hospital Clínico Universitario, Valladolid; ⁸Hospital General de Segovia; ⁹Hospital del Bierzo, León; ¹⁰Hospital Nuestra Señora de Sonsoles, Ávila, an ¹¹Hospital Morales Meseguer, Murcia, Instituto de Estudios de Ciencias de la Salud de Castilla y León (IECSCYL), Unidad de Investigación, Hospital Universitario de Salamanca, Spain

Citation: Hernández J-A, Rodríguez A-E, González M, Benito R, Fontanillo C, Sandoval V, Romero M, Martín-Núñez G, de Coca AG, Fisac R, Galende J, Recio I, Ortuño F, García J-L, de las Rivas J, Gutiérrez N-C, San Miguel J-F, and Hernández J-M. A high number of losses in 13q14 chromosome is associated with a worse outcome and biological differences in patients with B chronic lymphoid leukemia (B-CLL). *Haematologica* 2009; doi:10.3324/haematol.13862



Online Supplementary Figure S1. Overall survival (Figure 1sa) and time to first therapy (Figure 1sb) of patients with B-CLL and 13q14 deletion as the sole aberration and a percentage of FISH losses < 50 %, 51-79 % or \geq 80 %.