


IDH1 mutations in patients with myelodysplastic syndromes are associated with an unfavorable prognosis

Felicitas Thol,¹ Eva M. Weissinger,¹ Jürgen Krauter,¹ Katharina Wagner,¹ Frederik Damm,¹ Martin Wichmann,¹ Gudrun Göhring,² Christiane Schumann,³ Gesine Bug,⁴ Oliver Ottmann,⁴ Wolf-Karsten Hofmann,³ Brigitte Schlegelberger,² Arnold Ganser,¹ and Michael Heuser¹

¹Department of Hematology, Hemostasis, Oncology, and Stem Cell Transplantation, Hannover Medical School, Hannover, Germany; ²Institute of Cellular and Molecular Pathology, Hannover Medical School, Hannover, Germany; ³Department of Hematology and Oncology, University Hospital Mannheim, Mannheim, Germany, and ⁴Department of Internal Medicine III, University of Frankfurt, Frankfurt, Germany

Citation: Thol F, Weissinger EM, Krauter J, Wagner K, Damm F, Wichmann F, Göhring G, Schumann C, Bug G, Ottmann O, Hofmann W-K, Schlegelberger B, Ganser A, and Heuser M. IDH1 mutations in patients with myelodysplastic syndromes are associated with an unfavorable prognosis. *Haematologica* 2010;95(10):1668-1674. doi:10.3324/haematol.2010.025494

Sample Description



Online Supplementary Figure S1. Description of patient samples. MDS, myelodysplastic syndrome; AML, acute myeloid leukemia