

Combined CD8⁺ and CD4⁺ adenovirus hexon-specific T cells associated with viral clearance after stem cell transplantation as treatment for adenovirus infection

Maarten L. Zandvliet,¹ J.H. Frederik Falkenburg,² Ellis van Liempt,² Louise A. Veltrop-Duits,³ Arjan C. Lankester,³ Jayant S. Kalpoe,⁴ Michel G.D. Kester,² Dirk M. van der Steen,² Maarten J. van Tol,³ Roel Willemze,² Henk-Jan Guchelaar,¹ Marco W. Schilham,³ and Pauline Meij¹

¹Department of Clinical Pharmacy and Toxicology; ²Department of Hematology; ³Department of Pediatrics, and ⁴Department of Medical Microbiology, Leiden University Medical Center, Leiden, The Netherlands

Citation: Zandvliet ML, Falkenburg JH, van Liempt E, Veltrop-Duits LA, Lankester AC, Kalpoe JS, Kester MG, van der Steen DM, van Tol MJ, Willemze R, Guchelaar HJ, Schilham MW, and Meij P. Combined CD8⁺ and CD4⁺ adenovirus hexon-specific T cells associated with viral clearance after stem cell transplantation as treatment for adenovirus infection. *Haematologica* 2010;95(11):1943-1951. doi:10.3324/haematol.2010.022947

Online Supplementary Table S1. Characterization of HAdV hexon CD8⁺ epitopes recognized by T-cell lines as determined by peptide-MHC I tetramer staining or recognition of specific hexon peptides.


Donor	Peptide	HAdV hexon CD8 ⁺ epitopes recognized			Newly identified		
		Previously described	AA	HLA restriction	Peptide	AA	HLA restriction
1	MPN	320-329		B35			
	IPY	705-713		B35	IPF	873-882	B35
2	YVL	917-926		A2			
	MPN	320-329		B35	IPF	873-882	B35
	IPY	705-713		B35			
3	YVL	917-926		A2			
	KPY	114-124		B7			
	MPN	320-329		B7	FRK	585-593	B7
	TYF	37-45		A24			
4	TDL	886-894		A1	YSY	78-86	B63
5	TYF	37-45		A24			
6	TDL	886-894		A1	ETY	36-45	B52
					YSY	78-86	B63

Online Supplementary Table S2. Conservation of HAdV hexon CD8⁺ epitopes between different HAdV species.

Previously described HAdV hexon CD8 ⁺ epitopes							
HAdV species	HAdV serotype	AA 37-45	AA 114-124	AA 320-329	AA 705-713	AA 886-894	AA 917-926
C	5	TYFSLNNKF	KPYSGTAYNAL	M P N R P N Y I A F	I P Y L D G T F Y	T D L G Q N L L Y	Y V L F E V F D V V
A	12	---T-G---	-----S-	A---A-----	-----	-----M--	-----
B	35	---N-G---	-----S-	-----G-	-----	-----M--	-L-----
C	2	-----	-----	-----	-----	-----	-----
D	30	----G---	-----S-	-----G-	-----	-----M--	-L-----
E	4	----G---	-----S-	-----G-	-----	-----M--	--V-----
F	40	----G---	-----S-	A-----G-	V-----	-----M--	-----

Newly identified HAdV hexon CD8 ⁺ epitopes							
HAdV species	HAdV serotype	AA 36-45	AA 78-86	AA 585-593	AA 873-882		
C	5	ETYFSLNNKF	YSYKARFTL	FRKDVNVMVL	IPFSSNFMSM		
A	12	D---T-G---	-----	-----I-	-----		
B	35	D---N-G---	---V-Y--	-----	-----		
C	2	-----	-----	-----	-----		
D	30	D---G---	-----	-----I-	-----		
E	4	D---G---	---V-Y--	-----I-	-----		
F	40	D---G---	---V---	-----I-	-----		

HAdV hexon sequences were compared between serotype 5 (AP_000211), serotype 12 (AP_000121), serotype 35 (AP_000585), serotype 2 (AP_000175), serotype 30 (ABA00012), serotype 4 (AAO24097), and serotype 40 (NP_040862). The sequence derived from serotype 5 is depicted for each HAdV hexon CD8⁺ epitope, and amino acids that differ are shown for the other serotypes.



Online Supplementary Figure S1. Confirmation of newly identified HAdV hexon CD8⁺ T-cell epitopes. Intracellular IFN γ staining of HAdV-specific T-cell lines after restimulation with minimal MHC class I restricted HAdV hexon peptides. In addition, the HAdV-specific T-cell line derived from donor 2 was stained with IPF/HLA-B35 tetramers. To confirm HLA-restriction of the other epitopes, IFN γ release was determined by ELISA upon incubation with autologous or partially HLA-matched allogeneic EBV-LCL without peptide (white bars), or loaded with the relevant minimal HAdV hexon peptide (black bars).