


p-ERK1/2 is a predictive factor of response to erythropoiesis-stimulating agents in low/int-1 myelodysplastic syndromes

Emilie Frisan,^{1,2,3,5} Patrycja Pawlikowska,^{1,2,3} Cecile Pierre-Eugène,^{1,2,3,4} Vivian Viallon,^{5,6} Laure Gibault,^{5,7} Sophie Park,^{1,2,3,5,8} Patrick Mayeux,^{1,2,3} François Dreyfus,^{1,2,3,5,8} François Porteu,^{1,2,3} Michaëla Fontenay^{1,2,3,4,5}

¹Institut Cochin, Département d'Immunologie-Hématologie, ²INSERM U1016, ³Centre National de la Recherche Scientifique, Unité mixte de recherche 8104; ⁴Assistance Publique-Hôpitaux de Paris, Hôpital Broca-Cochin-Hôtel-Dieu, Service d'hématologie biologique; ⁵Université Paris Descartes, Faculté de médecine; ⁶Assistance Publique-Hôpitaux de Paris, Hôpital Cochin, Département d'Epidémiologie et de Biostatistiques; ⁷Assistance Publique-Hôpitaux de Paris, Hôpital Cochin, Service d'Anatomo-pathologie; ⁸Assistance Publique-Hôpitaux de Paris, Hôpital Cochin, Service de Médecine Interne, Unité fonctionnelle d'hématologie, Paris, France

Citation: Frisan E, Pawlikowska P, Pierre-Eugène C, Viallon V, Gibault L, Park S, Mayeux P, Dreyfus F, Porteu F, and Fontenay M. p-ERK1/2 is a predictive factor of response to erythropoiesis-stimulating agents in low/int-1 myelodysplastic syndromes. *Haematologica* 2010;95(11):1964-1968. doi:10.3324/haematol.2010.024349

Online Supplementary Figure S1. Flow chart.



Online Supplementary Figure S2. Decreased amplification rate and increased apoptosis of erythroid progenitors from MDS non-responsive to ESA. (A) Growth curves of *in vitro* cultured erythroid progenitors for 14 days (controls: diamonds, MDS responsive to ESA: square, MDS non-responsive to ESA: triangle). (B) CD34⁻-derived BFU-E at day 4 of the liquid culture. (C) Apoptosis measured by annexinV labeling at day 14 of the liquid culture.

