


Reassessment of small lymphocytic lymphoma in the era of monoclonal B-cell lymphocytosis

Sarah E. Gibson,¹ Steven H. Swerdlow,¹ Judith A. Ferry,² Urvashi Surti,^{1,3} Paola Dal Cin,⁴ Nancy Lee Harris,² and Robert P. Hasserjian²

¹Department of Pathology, University of Pittsburgh School of Medicine, Pittsburgh, PA; ²Department of Pathology, Massachusetts General Hospital, Boston, MA; ³Pittsburgh Cytogenetics Laboratory, Magee-Women's Hospital of UPMC, Pittsburgh, PA, and ⁴Center for Advanced Molecular Diagnostics, Brigham and Women's Hospital, Boston, MA, USA

Citation: Gibson SE, Swerdlow SH, Ferry JA, Surti U, Dal Cin P, Harris NL, and Hasserjian RP. Reassessment of small lymphocytic lymphoma in the era of monoclonal B-cell lymphocytosis. *Haematologica* 2011;96(8):1144-1152 doi:10.3324/haematol.2011.042333



Online Supplementary Figure S1. Examples of lymph nodes involved by CLL/SLL in patients with $<5 \times 10^9/L$ peripheral blood monoclonal B-cells. (A) Enlarged axillary lymph node diffusely involved by SLL with a residual germinal center (arrowhead); hematoxylin-eosin (H&E), original magnification x20). (B) Small cervical lymph node diffusely involved by SLL with numerous scattered proliferation centers (H&E, original magnification x20). (C) Enlarged axillary lymph node with focal interfollicular/intersinus involvement by CLL/SLL cells (arrowhead) (H&E, original magnification x100). (D) Occasional proliferation centers are present (H&E, original magnification x400). Immunohistochemical stains for CD20 (E), CD3 (F), and CD5 (G) highlight the infiltrate of CD20⁺, CD5^{dim} SLL cells (original magnification x400).