

High pentraxin 3 level predicts septic shock and bacteremia at the onset of febrile neutropenia after intensive chemotherapy of hematologic patients

Matti Vänskä,^{1,*} Irma Koivula,^{2,*} Sari Hämäläinen,² Kari Pulkki,³ Tapio Nousiainen,⁴ Esa Jantunen,^{4,5} and Auni Juutilainen^{1,5}

¹Department of Medicine, Kuopio University Hospital, Kuopio; ²Department of Medicine, Division of Infectious Diseases, Kuopio University Hospital, Kuopio; ³Department of Clinical Chemistry, University of Eastern Finland and Eastern Finland Laboratory Centre, Kuopio; ⁴Department of Medicine, Division of Hematology, Kuopio University Hospital, Kuopio; ⁵Institute of Clinical Medicine/Internal Medicine, University of Eastern Finland, Kuopio, Finland

*MV and IK contributed equally to this manuscript.

Citation: Vänskä M, Koivula I, Hämäläinen S, Pulkki K, Nousiainen T, Jantunen E, and Juutilainen A. High pentraxin 3 level predicts septic shock and bacteremia at the onset of febrile neutropenia after intensive chemotherapy of hematologic patients. *Haematologica* 2011;96(9):1385-1389. doi:10.3324/haematol.2011.044925

Online Supplementary Table S1. Association of the baseline characteristics with PTX3 on day 0 in 100 hematologic patients with febrile neutropenia after chemotherapy. The median of PTX3 on day 0 from the beginning of fever was 9.0 µg/L.

| | PTX3 d0 below median (N=50) | PTX3 d0 above median (N=50) | P value |
|---|-----------------------------------|-----------------------------------|---------|
| Male (n, % of all patients) | 29 (58%) | 32 (64%) | 0.539 |
| Age (years) (mean, range) | 51 (18-70) | 52 (22-66) | 0.748 |
| ≥ 60 years (n, % of patients) | 17 (34%) | 15 (30%) | 0.668 |
| Duration of neutropenic fever (d) (mean, range) | 4.5 (1-19) | 6.3 (1-25) | 0.174 |
| Duration of neutropenia (d) (mean, range) | 12.8 (4-30) | 11.4 (3-80) | 0.149 |
| Comorbidities | | | |
| Cardiovascular disease | 12 (24%) | 7 (14%) | 0.202 |
| Diabetes | 2 (4%) | 4 (8%) | 0.400 |
| Other | 12 (24%) | 9 (18%) | 0.461 |
| Treatment | | | |
| BEAM | 13 (26%) | 19 (38%) | 0.198 |
| BEAC | 4 (8%) | 11 (22%) | 0.050 |
| HD-MEL | 8 (16%) | 13 (26%) | 0.220 |
| IAT | 11 (22%) | 3 (6%) | 0.021 |
| HD-AraC-Ida | 8 (16%) | 0 (0%) | 0.003 |
| IdAraC-Ida | 5 (10%) | 4 (8%) | 0.727 |
| Mito-HDAraC | 1 (2%) | 0 (0%) | 0.315 |

BEAM: carmustine, etoposide, cytarabine and melphalan; BEAC: carmustine, etoposide, cytarabine and cyclophosphamide; HD-MEL: high-dose melphalan; IAT: idarubicin, cytarabine, thioguanine; HD-AraC-Ida: high-dose cytarabine, idarubicin; IdAraC-Ida: intermediate dose cytarabine, idarubicin; Mito-HDAraC: mitoxantrone, high-dose cytarabine