


Cord blood-derived CD34⁺ hematopoietic cells with low mitochondrial mass are enriched in hematopoietic repopulating stem cell function

Damia Romero-Moya,¹ Clara Bueno,¹ Rosa Montes,¹ Oscar Navarro-Montero,¹ Francisco J. Iborra,² Luis Carlos López,³ Miguel Martín,¹ and Pablo Menéndez^{1,4,5}

¹GENyO (Centre for Genomics and Oncological Research: Pfizer / University of Granada / Andalusian Government), Granada;

²Department of Molecular and Cellular Biology, Centro Nacional de Biotecnología, Consejo Superior de Investigaciones Científicas, Madrid; ³Centro de Investigación Biomédica, Instituto de Biotecnología, Universidad de Granada; ⁴Josep Carreras Leukemia Research Institute, Barcelona; and ⁵Institució Catalana de Recerca i Estudis Avançats, (ICREA), Barcelona, Spain

©2013 Ferrata Storti Foundation. This is an open-access paper. doi:10.3324/haematol.2012.079244



Online Supplementary Figure S1. (A) Levels of reactive oxidative species (ROS) measured by MitoSox staining in CD34⁺ Mito^{Low} cells and Mito^{High} cells. (B) Quantitative reverse transcriptase polymerase chain reaction analysis of HIF1 α and Meis1 in CD34⁺ Mito^{Low} cells relative to CD34⁺ Mito^{High} cells. The dotted horizontal line represents the gene expression level in CD34⁺ Mito^{High} cells. β -actin was used as a housekeeping gene.