

Errata Corrige

European guidelines for empirical antibacterial therapy for febrile neutropenic patients in the era of growing resistance: summary of the 2011 4th European Conference on Infections in Leukemia

Diana Averbuch,¹ Christina Orasch,² Catherine Cordonnier,³ David M. Livermore,⁴ Małgorzata Mikulska,⁵ Claudio Viscoli,⁵ Inge C. Gyssens,^{6,7,8} Winfried V. Kern,⁹ Galina Klyasova,¹⁰ Oscar Marchetti,² Dan Engelhard,¹ and Murat Akova;¹¹ on behalf of ECIL4, a joint venture of EBMT, EORTC, ICHS, ESGICH/ESCMID and ELN

¹Pediatric Infectious Diseases Unit, Hadassah-Hebrew University Medical Center, Jerusalem, Israel; ²Infectious Diseases Service, Department of Medicine, Lausanne University Hospital, Switzerland; ³APHP-Henri Mondor Hospital, Hematology Department and Université Paris Est -Créteil, France; ⁴Norwich Medical School, University of East Anglia, Norwich, UK; ⁵Division of Infectious Diseases, University of Genova, IRCCS San Martino-IST, Genoa, Italy; ⁶Department of Medicine and Nijmegen Institute for Infection, Inflammation and Immunity (N4i), Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands; ⁷Department of Medical Microbiology and Infectious Diseases, Canisius Wilhelmina Hospital, Nijmegen, The Netherlands; ⁸Hasselt University, Diepenbeek, Belgium; ⁹Center for Infectious Diseases and Travel Medicine, Department of Medicine, University Hospital, Albert-Ludwigs University, Freiburg, Germany; ¹⁰National Research Center for Hematology, Moscow, Russia; and ¹¹Department of Medicine, Section of Infectious Diseases, Hacettepe University School of Medicine, Ankara, Turkey

©2013 Ferrata Storti Foundation. This is an open-access paper. doi:10.3324/haematol.2014.103770

An incorrect version of figure 1 appeared on December 2013 Issue, page 1831. The correct version of figure 1 is published below.




Figure 1. Recommended strategies in various circumstances when using an escalation approach.