
The clinical significance of patients' sex in chronic lymphocytic leukemia

Daniel Catovsky,1 Rachel Wade,2 and Monica Else1


¹Haemato-Oncology Research Unit, Division of Molecular Pathology, The Institute of Cancer Research, London; and ²Clinical Trial Service Unit, Oxford, UK.

©2014 Ferrata Storti Foundation. This is an open-access paper. doi:10.3324/haematol.2013.101378 Manuscript received on November 19, 2013. Manuscript accepted on March 18, 2014. Correspondence: daniel.catovsky@icr.ac.uk


SUPPLEMENTARY FIGURE 1: Overall survival by sex in each individual randomized trial




1A: CLL1 (363 patients)



1B: CLL2 (286 patients)




1C: CLL3 (395 patients)




1D: LRF CLL4 (777 patients)

SUPPLEMENTARY FIGURE 2: Overall survival by sex within sub-groups




2A: Overall survival by sex within each disease stage (all randomized trials)




2B: Overall survival by sex within each age group (all randomized trials)

SUPPLEMENTARY FIGURE 3: Overall survival with response to treatment, and progression-free survival, by sex



3A: Overall survival by sex and response to first-line treatment (all randomized trials)



3B: Progression-free survival by sex (LRF CLL4 trial only)