

Rituximab and three dexamethasone cycles provide responses similar to splenectomy in women and those with immune thrombocytopenia of less than two years duration

James B. Bussel,¹ Christina S. Lee,¹ Caroline Seery,¹ Allison A. Imahiyerobo,¹ Michaela V. Thompson,¹ Diane Catellier,² Ithamar G. Turenne,¹ Vivek L. Patel,^{1,3} Paul A. Basciano,¹ Rebecca L. Elstrom,^{1,4} and Waleed Ghanima^{1,5}

¹Departments of Pediatrics and Medicine, Division of Hematology/Oncology, Weill Cornell Medical College, New York, NY, USA; ²Research Triangle Institute, Chapel Hill, NC, USA; ³Albert Einstein College of Medicine, MD PhD program, Bronx, NY, USA; ⁴Genentech, South San Francisco, CA, USA; and ⁵Department of Medicine, Østfold Hospital Trust, Fredrikstad, Norway

©2014 Ferrata Storti Foundation. This is an open-access paper. doi:10.3324/haematol.2013.103291

Manuscript received on January 7, 2014. Manuscript accepted on April 17, 2014.

Correspondence: jbussel@med.cornell.edu

Supplemental Table 1. Genotypes of H β 1-tubulin single nucleotide polymorphisms on 54 patients

Total: n=54	<u>A/A</u>	<u>A/B</u>	<u>B/B</u>
Any Response	32/44 (73%)	9/10 (90%)	0
CR	27/32	7/9	0
PR	5/32	2/9	0
Response Duration >12 months	26/32 (81%)	8/9 (89%)	0
CR	24/27	7/7	0
PR	2/5	1/2	0
Relapse (<50x10 ⁹ /L)	10/32 (31%)	2/9 (22%)	0
CR	6/27	0	0
PR	4/5	1/2	0
NR	12/44 (27%)	1/10 (10%)	0

A= Wild Type Allele

B = SNP Allele