

# Tyrosine kinase inhibitors improve long-term outcome of allogeneic hematopoietic stem cell transplantation for adult patients with Philadelphia chromosome positive acute lymphoblastic leukemia

Eolia Brissot,<sup>1,2,3,4</sup> Myriam Labopin,<sup>1,2,3</sup> Marielle M. Beckers,<sup>5</sup> Gérard Socié,<sup>6</sup> Alessandro Rambaldi,<sup>7</sup> Liisa Volin,<sup>8</sup> Jürgen Finke,<sup>9</sup> Stig Lenhoff,<sup>10</sup> Nicolaus Kröger,<sup>11</sup> Gert J. Ossenkoppele,<sup>12</sup> Charles F. Craddock,<sup>13</sup> Ibrahim Yakoub-Agha,<sup>14</sup> Günhan Gurman,<sup>15</sup> Nigel H. Russell,<sup>16</sup> Mahmoud Aljurf,<sup>17</sup> Michael N. Potter,<sup>18</sup> Armon Nagler,<sup>19</sup> Oliver Ottmann,<sup>20</sup> Jan J. Cornelissen,<sup>21</sup> Jordi Esteve,<sup>22</sup> and Mohamad Mohty<sup>1,2,3</sup>

<sup>1</sup>Université Pierre et Marie Curie, Paris, France; <sup>2</sup>INSERM, UMRs 938, Paris, France; <sup>3</sup>Service d'Hématologie Clinique et de Thérapie Cellulaire, Hôpital Saint Antoine, APHP, Paris, France; <sup>4</sup>CHRU Hôtel-Dieu, Nantes, France; <sup>5</sup>University Hospital Leuven, Leuven, Belgium; <sup>6</sup>Hôpital Saint Louis, Paris, France; <sup>7</sup>Azienda Ospedaliera Papa Giovanni XXIII, Bergamo, Italy; <sup>8</sup>Helsinki University Central Hospital, Finland; <sup>9</sup>University of Freiburg Medical Center, Germany; <sup>10</sup>Lund University Hospital, Sweden; <sup>11</sup>University Hospital Hamburg-Eppendorf, Hamburg, Germany; <sup>12</sup>VU University Medical Center, Amsterdam, The Netherlands; <sup>13</sup>Queen Elizabeth Hospital, Birmingham, UK; <sup>14</sup>CHRU Lille, France; <sup>15</sup>Ankara University, Faculty of Medicine, Ankara, Turkey; <sup>16</sup>Nottingham University Hospital, Nottingham, UK; <sup>17</sup>King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia; <sup>18</sup>The Royal Marsden NHS Foundation Trust, London, UK; <sup>19</sup>Chaim Sheba Medical Center, Tel Hashomer, Israel; <sup>20</sup>Goethe-University Frankfurt, Germany; <sup>21</sup>Erasmus Medical Center-Daniel den Hoed Cancer Center, Rotterdam, The Netherlands; <sup>22</sup>Hospital Clinic Institut d'Investigacions Biomèdiques August Pi i Sunyer, Barcelona, Spain

©2015 Ferrata Storti Foundation. This is an open-access paper. doi:10.3324/haematol.2014.116954

Manuscript received on September 8, 2014. Manuscript accepted on December 15, 2014.

Correspondence: mohamad.mohty@inserm.fr

**Table S1- Patient characteristics**

Characteristics	TKIs before transplantation		p-value
	No	Yes	
<b>Age at allo-SCT (years), median (range)</b>	40 (20-58)	43 (18-70)	0.005
<b>WBC at diagnosis (no.X10<sup>9</sup>/L), median (range)</b>	21 (0.7-363)	21 (0.3-640)	0.2
<b>Interval from diagnosis to allo-SCT (days), median (range)</b>	152 (80-432)	160 (67-503)	0.24
<b>Interval from diagnosis to CR1 (days), median (range)</b>	43 (21-188)	43 (15-251)	0.51
<b>Interval from CR1 to allo-SCT (days), median (range)</b>	101 (23-188)	100 (15-393)	0.70
<b>Transplant year, median (range)</b>	2002 (00-10)	2007 (00-10)	<0.0001
<b>Patient gender, no. (%)</b>			
Male	49 (59)	211 (54)	
Female	34 (41)	179 (46)	0.41
<b>Donor gender, no. (%)</b>			
Male	54 (65)	252 (65)	
Female	29 (35)	138 (35)	0.94
<b>Female donor/ male R, no. (%)</b>			
No	62 (75)	324 (83)	
Yes	21 (25)	66 (17)	0.07
<b>Extramedullary disease, no. (%)</b>			
No	62 (81)	310 (85)	
Yes	15 (20)	54 (15)	0.31
<b>Type of donor, no. (%)</b>			
MSD	45 (54)	179 (46)	
MUD	38 (46)	211 (54)	0.17
<b>Stem cell source, no. (%)</b>			
Bone marrow	28 (34)	81 (21)	
Peripheral blood	55 (66)	309 (79)	0.01
<b>Conditioning regimen, no. (%)</b>			
MAC	77 (93)	298 (76)	
RIC	6 (7)	92 (24)	0,001
<b>MRD positive at allo-SCT, no. (%)</b>			
no	40 (73)	215 (63)	
yes	15 (27)	125 (37)	0.17
<b>ATG, no. (%)</b>			
no	56 (70)	254 (66)	
yes	24 (30)	131 (34)	0.49

TKIs: tyrosine kinase inhibitors; WBC: white blood cells; CR1: first complete remission; allo-SCT: hematopoietic stem cell transplantation; HLA: human leucocyte antigen; MSD: matched sibling donor; MUD: matched unrelated donor MAC: myeloablative conditioning; RIC: reduced intensity conditioning; MRD: minimal residual disease; ATG: antithymocyte globulin

**Table S2. Univariate analysis of TKIs post allo-SCT, aGVH and cGVHD on LFS, OS, RI and NRM at 5 years (Cox model with time dependent variable)**

variable	LFS		OS		RI		NRM	
	HR 95%CI	P-value	HR 95%CI	P-value	HR 95%CI	P-value	HR 95%CI	P-value
TKIs post-transplant in prophylaxis*	0.49 0.31-0.79	0.003	0.42 0.24-0.72	0.002	0.53 0.29-0.96	0.04	0.44 0.21-0.96	0.04
Acute GVHD $\geq$ II*	1.09 0.84-1.40	0.52	1.47 1.12-1.92	0.01	0.52 0.35-0.76	0.001	2.63 1.80-3.86	0.0000
Chronic GVHD*	1.15 0.86-1.54	0.36	1.19 0.88-1.60	0.25	0.52 0.34-0.78	0.002	3.80 2.31-6.26	0.0000

LFS: leukemia free survival; OS: overall survival; RI: relapse incidence; NRM: non relapse mortality; TKIs: tyrosine kinase inhibitors

\*time dependant variable

**Table S3. Univariate analysis of TKIs post allo-SCT on aGVH and cGVHD (Cox model with time dependant variable)**

	p	HR	95% CI
Acute GVHD $\geq$ II*	0.03	0.21	0.05-0.86
Chronic GVHD*	0.007	0.6	0.41-0.87

\*time dependant variable

**Table S4. Multivariate analysis for aGVHD and cGVHD**

variable	aGVHD		cGVHD	
	HR 95%CI	P-value	HR 95%CI	P-value
TKIs before transplant	1.52 0.92-2.52	0.10	1.46 0.94-2.27	0.09
Age at transplant (years)	1.00 0.98-1.01	0.83	1.01 0.99-1.02	0.29
WBC at diagnosis	1.00 0.999-1.002	0.61	0.997 0.995-1.000	0.02
Interval from diagnosis to transplant (d)	0.99 0.997-1.001	0.39	1.000 0.997-1.002	0.66
Female R vs male R	1.01 0.70-1.44	0.97	0.96 0.69-1.34	0.81
Female D to male R	0.84 0.51-1.39	0.49	1.31 0.89-1.93	0.17
MUD vs MSD	1.21 0.81-1.82	0.35	1.17 0.83-1.64	0.37
RIC vs MAC	0.83 0.52-1.32	0.42	0.77 0.51-1.17	0.23
ATG	0.85 0.56-1.28	0.44	0.57 0.40-0.82	0.002
TKIs post-transplant (all reasons) *	0.21 0.05-0.85	0.03	0.73 0.49-1.09	0.12
aGVHD $\geq$ II*	—		1.97 1.47-2.65	0.00001

HR: hazard ratio; TKIs: tyrosine kinase inhibitors; WBC: white blood cells; R: recipient; D: donor ; MUD: matched unrelated donor; MSD: matched sibling donor; RIC: reduced intensity conditioning; MAC: myeloablative conditioning; ATG: antithymocyte globulin, TKIs: tyrosine kinase inhibitors, aGVHD: acute graft-versus-host disease

\*Time dependant variable