

Regulatory T cells and progenitor B cells are independent prognostic predictors in lower risk myelodysplastic syndromes

Josephine D. Kahn, Martine E.D. Chamuleau, Theresa M. Westers, Peter M. Van de Ven, Linda van Dreunen, Margot van Spronsen, Gert J. Osenkoppela, and Arjan A. van de Loosdrecht*

Correspondence: *a.vandeloosdrecht@vumc.nl*
doi:10.3324/haematol.2014.116657

Supplementary table 1. Data on T cell subsets, NK cells and progenitor B cells.

Peripheral blood (PB): % represents percentage of lymphocytes

Bone marrow (BM): % represents percentage of leukocytes

	Median	Range
<i>PB T lymphocyte subpopulations</i>		
% CD4 ⁺ CD45RA ⁺⁺ CD27 ⁺ (CD4 naïve)	33.1	7,2 – 72.9
% CD4 ⁺ CD45RA ⁺⁺ CD27 ⁻ (CD4 effector)	0.9	0.0 – 26.3
% CD8 ⁺ CD45RA ⁺⁺ CD27 ⁺ (CD8 naïve)	18.2	2.0 – 55.0
% CD8 ⁺ CD45RA ⁺⁺ CD27 ⁻ (CD8 effector)	25.6	2.1 – 71.2
<i>PB regulatory T cells</i>		
% CD4 ⁺ CD25 ^{hi} FoxP3 ⁺	5.8	2.3 – 10.3
<i>PB NK cells</i>		
% CD3 ⁺ CD16 ⁺ CD56 ^{dim}	5.5	0.9 – 18.0
<i>BM progenitor B cells</i>		
% CD10 ⁺ CD34 ⁺	9.7	1.0 – 40.0