

Predictive models for ocular chronic graft-versus-host disease diagnosis and disease activity in transplant clinical practice

Lauren M. Curtis,¹ Manuel B. Datiles III,² Seth M. Steinberg,³ Sandra A. Mitchell,⁴ Rachel J. Bishop,² Edward W. Cowen,⁵ Jacqueline Mays,⁶ John M. McCarty,⁷ Zoya Kuzmina,¹ Filip Pirsli,¹ Daniel H. Fowler,¹ Ronald E. Gress¹ and Steven Z. Pavletic¹

¹Experimental Transplantation and Immunology Branch, National Cancer Institute (NCI), National Institutes of Health (NIH), Bethesda, MD; ²National Eye Institute, NIH, Bethesda, MD; ³Biostatistics and Data Management Section, NCI, NIH, Rockville, MD; ⁴Outcomes Research Branch, Division of Cancer Control and Population Sciences, NCI, NIH, Rockville, MD; ⁵Dermatology Branch, NCI, NIH, Bethesda, MD; ⁶National Institutes of Dental and Craniofacial Research, NIH, Bethesda, MD; and ⁷Bone Marrow Transplant Program, Massey Cancer Center, Virginia Commonwealth University Medical Center, Richmond, VA, USA

©2015 Ferrata Storti Foundation. This is an open-access paper. doi:10.3324/haematol.2015.124131

Manuscript received on January 21, 2015. Manuscript accepted on June 16, 2015.

Correspondence: pavletis@mail.nih.gov

Supplemental data:

#1: Supplemental Table #1: Univariate analysis evaluating any association between diagnosis of ocular cGVHD and three types of GVHD prophylaxis at time of transplant

Type of GVHD prophylaxis	Patients	Ocular cGVHD present (n,%)	Ocular cGVHD absent (n,%)	<i>p</i>
Cyclosporine -based	Cyclosporine-based GVHD prophylaxis n=83	57 (69)	26 (31)	0.14
	Non-cyclosporine based prophylaxis n=121	95 (79)	26 (21)	
Tacrolimus -based	Tacrolimus-based GVHD prophylaxis n=80	63 (79)	17 (21)	0.32
	Non-tacrolimus based GVHD prophylaxis n=124	89 (72)	35 (28)	
T-cell depletion (ATG, ex-vivo T-cell depletion, alemtuzumab)	T-cell depleting GVHD prophylaxis n=33	25 (76)	8 (24)	1.00
	Non-T-cell depleting GVHD prophylaxis n=171	127 (74)	44 (26)	

Note: 6 (3%) patients with missing GVHD prophylaxis data.