

Haematologica  
HAEMATOL/2015/126300  
Version 3  
ANGIOIMMUNOBLASTIC T-CELL LYMPHOMA IS THE MOST  
COMMON T-CELL LYMPHOMA IN TWO DISTINCT FRENCH  
INFORMATION DATA SETS

Disclosures: The authors have no conflicts of interest to disclose

Contributions: Conception and design: Laurence de Leval, Marie Parrens, Philippe Gaulard, Collection and assembly of data: all authors Data analysis and interpretation: Laurence de Leval, Marie Parrens, Fabien Le Bras, Jean-Philippe Jais, Virginie Fattacioli, Corinne Haioun, Christian Gisselbrecht, Philippe Gaulard Manuscript writing: all authors Final approval of manuscript: all authors Financial support: Laurence de Leval, Philippe Gaulard Administrative support: Laurence de Leval, Philippe Gaulard Provision of study materials or patients: Laurence de Leval, Marie Parrens, Fabien Le Bras, Antoine Martin, Laurence Lamant, Richard Delarue, Françoise Berger, Flavie Abion, Céline Bossard, Marie-Christine Copin, Danielle Canoni, Frédéric Charlotte, Gandhi Damaj, Peggy Dartigues, Bettina Fabiani, Albane Ledoux-Pilon, Karine Montagne, Thierry Molina, Martine Patey, Patrick Tas, Michel Pech, Barbara Petit, Tony Petrella, Jean-Michel Picquenot, Thérèse Rousset, Marie-Christine Rousselet, Isabelle Soubeyran, Sylvie Thiebault, Olivier Tournilhac, Luc Xerri, Christian Gisselbrecht, Corinne Haioun, Georges Delsol, Philippe Gaulard