

Haematologica
HAEMATOL/2015/133736
Version 3

THE ANTI-TUMORAL EFFECT OF LENALIDOMIDE IS INCREASED
IN VIVO BY HYPOXIA-INDUCIBLE FACTOR (HIF)-1 α INHIBITION IN
MYELOMA CELLS

Paola Storti, Denise Toscani, Irma Airoidi, Valentina Marchica, Sophie Maiga, Marina Bolzoni, Elena Fiorini, Nicoletta Campanini, Eugenia Martella, Cristina Mancini, Daniela Guasco, Valentina Ferri, Gaetano Donofrio, Franco Aversa, Martine Amiot, and Nicola Giuliani

Disclosures: N.G. research funding from Celgene Italy and Janssen Pharmaceutical

Contributions: PS and DT performed all the key cell culture experiments, analyzed data and contributed to write the manuscript; I.A. performed the in vivo studies; V.M., M.B., E.F., D.G., V.F. and S.M. contributed to cell culture experiments, data analysis and western blot; C.C., E.M. and C.M.: performed the histological analysis of bone biopsies; G.D. performed the lentivirus transduction; F.A.: vital reagents and edited the manuscript; M.A. contributed to design the study; N.G.: design the study, analyzed data and wrote the manuscript.