

Haematologica
HAEMATOL/2015/137711
Version 3

Loss of B cells and their precursors is the most constant feature of
GATA-2 deficiency in childhood myelodysplastic syndrome

Michaela Novakova, Marketa Zaliova, Martina Sukova, Marcin Wlodarski, Ales Janda, Eva Fronkova, Vit Campr, Katerina Lejhancova, Ondrej Zapletal, Dagmar Pospisilova, Zdenka Cerna, Tomas Kuhn, Peter Svec, Vendula Pelkova, Zuzana Zemanova, Gitte Kerndrup, Marry van den Heuvel-Eibrink, Vincent van der Velden, Charlotte Niemeyer, Tomas Kalina, Jan Trka, Jan Stary, Ondrej Hrusak, and Ester Mejstrikova

Disclosures: none

Contributions: M.N. performed FC, GATA-2 and KREC-TREC analysis, wrote the manuscript, M.Z. and M.W. analyzed GATA-2 sequencing, M.S., A.J., K.L., O.Z., D.P., Z.C., T. Ku. and P.S. collected patient data and sent the samples, E.F. designed and analyzed KREC-TREC analysis, V.C. and G.K. performed histopathological analyses, V.P. performed flow cytometry, Z.Z. performed cytogenetic analyses, M.v.d.H.E and V.v.d.V. designed the flow cytometrical parameters, T.K. designed and performed flow cytometry, J.T. collected the material and designed the research, J.S. and C.N. designed the research, O.H. designed the research, performed flow cytometry, analyzed data and wrote the manuscript, E.M. designed the research, analyzed data, wrote the manuscript. All authors revised and approved the manuscript.