

Haematologica
HAEMATOL/2016/143628
Version 3

Umbilical cord blood transplantation is a suitable option for
consolidation of acute myeloid leukemia with FLT3-ITD

Craig E. Eckfeldt, Nicole Randall, Ryan M. Shanley, Sophia Yohe, Nelli
Bejanyan, Michelle Dolan, Erica D. Warlick, Michael R. Verneris,
Claudio G. Brunstein, John E. Wagner, Daniel J. Weisdorf, and
Celalettin Ustun

Disclosures: no conflict of interest related to this article Research reported in this publication was supported by NIH grant P30 CA77598 utilizing the Biostatistics and Bioinformatics Core shared resource of the Masonic Cancer Center, University of Minnesota and by the by the National Center for Advancing Translational Sciences of the National Institutes of Health Award Number UL1TR000114. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health

Contributions: CU: conceived the idea, did literature search, data collection, analyzed data, wrote the article, edited. CEE, NR: data collection, did literature search, analyzed data, wrote the article, edited. SY,MD, NB, EW, MRV, CGB, JEW, DJW: analyzed data, wrote the article, edited. RMS: data collection, analyzed data, wrote the article, edited.