

# Non-Hodgkin lymphoma in the developing world: review of 4539 cases from the International Non-Hodgkin Lymphoma Classification Project

Anamarija M. Perry,<sup>1</sup> Jacques Diebold,<sup>2</sup> Bharat N. Nathwani,<sup>3</sup> Kenneth A. MacLennan,<sup>4</sup> Hans K. Müller-Hermelink,<sup>5</sup> Martin Bast,<sup>6</sup> Eugene Boilesen,<sup>7</sup> James O. Armitage,<sup>1</sup> and Dennis D. Weisenburger<sup>3</sup>

<sup>1</sup>Department of Pathology, University of Manitoba, Winnipeg, Manitoba, Canada; <sup>2</sup>Department of Anatomic Pathology and Cytology, Hotel-Dieu, University Denis Diderot, Paris, France; <sup>3</sup>Department of Pathology, City of Hope National Medical Center, Duarte, CA, USA; <sup>4</sup>Section of Pathology and Leeds Institute of Molecular Medicine, St. James University Hospital, Leeds, UK; <sup>5</sup>Institute of Pathology, University of Würzburg, Germany; <sup>6</sup>Department of Internal Medicine, University of Nebraska Medical Center, Omaha, NE, USA; and <sup>7</sup>Center for Collaboration on Research Design and Analysis, College of Public Health, University of Nebraska Medical Center, Omaha, NE, USA

©2016 Ferrata Storti Foundation. This is an open-access paper. doi:10.3324/haematol.2016.148809

Received: May 4, 2016.

Accepted: June 23, 2016.

Pre-published: June 27, 2016.

Correspondence: [dweisenburger@coh.org](mailto:dweisenburger@coh.org)

---

**Supplemental Table 1. Cases of pediatric (<19 years) non-Hodgkin lymphoma by region**

	SEEU (N)	CSA (N)	NAF/ME/IN (N)	SAF (N)	FE (N)	Developing Regions (N)	Developed Regions (N)
<b>B-cell lymphomas</b>							
Diffuse large B-cell lymphoma	2	10	10	2	2	26	2
Follicular lymphoma, all grades	0	4	3	0	0	7	0
Marginal zone B-cell lymphoma, MALT type	0	2	0	0	0	2	0
Marginal zone lymphoma, nodal/splenic	0	3	1	0	0	4	0
Burkitt lymphoma	2	11	8	2	6	29	1
Primary mediastinal B-cell lymphoma	0	1	3	0	1	5	0
High-grade B-cell lymphoma, Burkitt-like	1	0	2	0	0	3	0
Precursor B-lymphoblastic leukemia/lymphoma	4	16	2	0	0	22	1
Unclassifiable high-grade B-cell lymphoma	2	1	1	0	1	5	
Subtotal	11	48	30	4	10	103	4
<b>T- and NK-cell lymphomas</b>							
Peripheral T-cell lymphoma, not otherwise specified	0	2	2	0	1	5	2
Precursor T-lymphoblastic leukemia/lymphoma	2	4	8	9	6	29	3
Extranodal NK/T-cell lymphoma, nasal type	0	0	0	0	1	1	0
Anaplastic large T-cell lymphoma, ALK+ and ALK-	0	4	2	0	4	10	3
Mycosis fungoides	0	1	0	0	0	1	0
Peripheral T-cell lymphoma, other types	0	2	1	0	0	3	0
Subtotal	2	13	13	9	12	49	8
<b>Total</b>	13	61	43	13	22	152	12

Abbreviations: SEEU, Southeastern Europe; CSA, Central/South America; NAF/ME/IN, North Africa, Middle East and India; SAF, Southern Africa; FE, Far East

**Supplemental Table 2.** Rare or unclassifiable cases of T-cell lymphoma by region.

<b>T-cell Lymphomas</b>	<b>SEEU (N)</b>	<b>CSA (N)</b>	<b>NAF/ME/IN (N)</b>	<b>SAF (N)</b>	<b>FE (N)</b>	<b>Developing Regions (N)</b>	<b>Developed Regions (N)</b>
Enteropathy-associated T-cell lymphoma	2	3	3	1	0	9	4
Hepatosplenic T-cell lymphoma	0	1	0	2	0	3	0
Subcutaneous panniculitis-like T-cell lymphoma	1	2	0	0	0	3	0
Unclassifiable high-grade T-cell lymphoma	1	7	4	2	0	14	1
<b>Total</b>	<b>4</b>	<b>13</b>	<b>6</b>	<b>5</b>	<b>0</b>	<b>28</b>	<b>5</b>

Abbreviations: SEEU, Southeastern Europe; CSA, Central/South America; NAF/ME/IN, North Africa, Middle East and India; SAF, Southern Africa; FE, Far East

### **Supplemental Table 3. Study participants**

#### North America:

RD Gascoyne and JM Connors (British Columbia Cancer Agency, Vancouver, Canada)

#### Western Europe:

TA Lister and A Norton (St. Bartholomew's Hospital, London, United Kingdom);  
B Coiffier and F Berger (Centre Hospitalier Universitaire de Lyon, Lyon, France);  
F Cavalli and E Pedrinis (Ente Ospedaliero Cantonale, Locarno, Switzerland); W  
Hiddemann (Klinikum der Ludwig-Maximilian, University of Munich, Munich,  
Germany); A Schauer (University of Göttingen, Göttingen, Germany); G Ott  
(Robert Bosch Hospital, Stuttgart, Germany); T Rüdiger (University of Würzburg,  
Würzburg, Germany)

#### Southeastern Europe:

S Gasparov, A Skrtic, M Dominis, S Dzebro Rocic, P Korac, and S Ostojic  
Kolonic (University Hospital Merkur, Zagreb, Croatia); S Dotlic, I Ilic, I Aurer, S  
Basic Kinda, I Radman, and B Labar (University Hospital Center Zagreb, Zagreb,  
Croatia); G Aralica, C Tomasovic-Loncaric and V Pejisa (University Hospital  
Dubrava, Zagreb, Croatia); D Dujmovic (University of Zagreb Medical School,  
Zagreb, Croatia); N Jonjic, K Lucin, A Duletic Nacinovic and T Valkovic (Faculty  
of Medicine, University of Rijeka, Rijeka, Croatia); I Karaman, D Lozic, S  
Madunic, and M Jakic Bubalo (University Hospital Center, Split, Croatia); J Rajc  
and J Sincic-Petricicevic (University Hospital Center, Osijek, Croatia); B Coha

(General Hospital Slavonski Brod, Slavonski Brod, Croatia); R Babok-Flegaric (General Hospital Varazdin, Varazdin, Croatia); Z Nemet-Lojan (General Hospital Koprivnica, Koprivnica, Croatia); B Fetica, A. Cucuianu, I Berindan-Neagoe, Lj Petrov, and P Buiga Rares (Oncology Institute "Ion Chiricuta", Cluj-Napoca, Romania); G Petrusevska, A Stojanovic, L Cevrevska, D Dukovski, S Trajkova, N Basheska, M Klimper, J Petrovska and M Petrusevska (University of "Ss Cyril and Methodius", Skopje, Macedonia).

Central/South America:

MI Narbaitz, S Pavlosvsky and F Sackmann (Academia Nacional de Medicina, Buenos Aires, Argentina); V Piana de Andrade and C Mello (Hospital AC Camargo, Sao Paulo, Brazil); V Martinez and ME Cabrera (Instituto Anatomia Patologica and Hospital del Salvador, University of Chile, Santiago, Chile); HM Kirsch (Laboratorio de Patologia, Guatemala City, Guatemala); C Barrionuevo (Instituto Nacional de Enfermedades Neoplasticas, Lima, Peru).

North Africa/Middle East/India:

N Boudjerra (CPMC – Hopital Mustapha, Algiers, Algeria); J Audouin (Hotel-Dieu, Paris, France); EA Ishak (Cairo University, Cairo, Egypt); A Abou-Elella (Lancaster General Health, Hershey, PA, USA); J Martin (King Faisal Specialist Hospital & Research Centre, Riyadh, Saudi Arabia); L Temmim (Kuwait Cancer Control Center, Al Shwaikh, Kuwait); K Naresh (Tata Memorial Hospital, Mumbai, India).

Southern Africa:

P Close and P Jacobs (University of Cape Town, Cape Town, South Africa); Y Perner and W Bezwoda (University of the Witwatersrand, Johannesburg, South Africa); R Hohle, L Levy and A Coutts (University of Zimbabwe, Harare, Zimbabwe)

Far East:

F Ho, R Liang and W Au (University of Hong Kong, Queen Mary Hospital, Pokfulam, Hong Kong); ESR Hardjolutito (Fakultas Kedokteran Universitas Indonesia, Jakarta, Indonesia); X Li (Cancer Hospital, Fudan University, Shanghai, China); P Wannakrairot and T Intragumtornchai (Chulalongkorn University, Bangkok, Thailand).