Haematologica HAEMATOL/2016/149112 Version 3

The immunophenotypical fingerprint of patients with primary antibody deficiencies is partially present in their asymptomatic first-degree relatives.

Delfien J.A. Bogaert, Marieke De Bruyne, Veronique Debacker, Pauline Depuydt, Katleen De Preter, Carolien Bonroy, Jan Philippe', Victoria Bordon, Bart N. Lambrecht, Tessa Kerre, Andrea Cerutti, Karim Y. Vermaelen, Filomeen Haerynck, and Melissa Dullaers

Disclosures: D.B. is a PhD fellow of the Research Foundation Flanders (FWO), a government granting agency. T.K. and K.V. are senior clinical investigators of the FWO. A.C. has an FWO-Odysseus grant. This research was funded by the University Hospital Ghent Spearhead Research Programs and the Jeffrey Modell Foundation.

Contributions: Delfien Bogaert, first author, recruited the study subjects, performed the experiments, did the statistical analyses and drafted the initial manuscript. Marieke De Bruyne assisted in sample handling and critically reviewed the manuscript. Veronique Debacker assisted in sample handling and in performing experiments. Pauline Depuydt and Katleen De Preter performed unsupervised computational clustering analyses of the data and critically reviewed and revised the manuscript. Carolien Bonroy and Jan Philippe supervised routine laboratory analyses and critically reviewed and revised the manuscript. Victoria Bordon, Bart Lambrecht, Tessa Kerre, and Karim Vermaelen carried out initial clinical and laboratory workup of patients and critically reviewed and revised the manuscript. Filomeen Haerynck, shared last author, conceptualized the study, carried out initial clinical and laboratory workup of patients and critically reviewed and revised the manuscript. Melissa Dullaers, shared last author, conceptualized the study, supervised the experiments and statistical analyses, and critically reviewed and revised the manuscript. All authors have approved the final manuscript as submitted.