

Haematologica  
HAEMATOL/2016/155184  
Version 4

Obstetric antiphospholipid syndrome: early variations of angiogenic factors are associated with adverse outcomes. The NOH-ANGIO observational cohort study

Éva Cochery-Nouvellon, Érick Mercier, Sylvie Bouvier, Jean-Pierre Balducci, Isabelle Quéré, Antonia Perez-Martin, Eve Mousty, Vincent Letouzey, and Jean-Christophe Gris

Disclosures: Financial support was provided by Nîmes University hospital via an internal funding scheme. The authors have no competing conflict of interest to disclose.

Contributions: J.-C. G., É. C.-N., É. M.1, S. B., and J.-P. B. contributed to the conception and design of the study. J.-C. G., J.-P. B., A. P.-M., I.Q., É.M.7, and V.L. acquired the samples whereas É.C.-N., É. M.1, and S.B. performed biological analysis. J.-C. G., S.B., J.-P.B., I.Q., A. P.-M., E.M.7 and V.L. contributed to data analysis and to interpretation. J.-C. G. drafted the manuscript and all authors revised it critically and approved the final version (J.-C. G., É. C.-N., É. M.1, S.B., J.-P. B., A. P.-M., I.Q., É.M.7, V.L.).