

Haematologica  
HAEMATOL/2016/153270  
Version 3

Increased peroxisome proliferator-activated receptor-gamma activity  
reduces imatinib uptake and efficacy in chronic myeloid leukemia  
mononuclear cells

Jueqiong Wang, Liu Lu, Chung H. Kok, Verity A. Saunders, Jarrad M.  
Goyne, Phuong Dang, Tamara M. Leclercq, Timothy P. Hughes, and  
Deborah L. White

Disclosures: T.P.H. received research funding from Ariad, CSL, Novartis, and BMS and received honoraria from and participated on advisory boards of Ariad, Pfizer, Novartis, and BMS. D.L.W. received research funding from Ariad, CSL, Novartis, and BMS and received honoraria from and participated on advisory boards of Novartis and BMS. The remaining authors declare no competing financial interests.

Contributions: J.W. performed the study, analyzed data, and wrote the manuscript; L.L. performed the study and reviewed the manuscript, C.H.K. analyzed data, and co-wrote the manuscript; V.A.S. performed the study, analyzed data, and reviewed the manuscript; J.M.G. performed the study and reviewed the manuscript; P.D. performed the study and reviewed the manuscript; T.M.L. reviewed the manuscript; T.P.H. and D.L.W. initiated and directed the study, and reviewed the manuscript. All authors read and approved the final manuscript.