## Risk of cardiovascular events and pulmonary hypertension following splenectomy – a Danish population-based cohort study from 1996-2012

Marianne Rørholt, 1 Waleed Ghanima, 2,3 Dora Körmendiné Farkas 4 and Mette Nørgaard 4

<sup>1</sup>Department of Research and Otorhinolaryngology, Østfold Hospital Trust, Norway; <sup>2</sup>Department of Research and Medicine, Østfold Hospital Trust, Norway; <sup>3</sup>Institute of Clinical Medicine, University of Oslo, and Department of Haematology Oslo University Hospital, Norway and <sup>4</sup>Department of Clinical Epidemiology, Aarhus University Hospital, Denmark

©2017 Ferrata Storti Foundation. This is an open-access paper. doi:10.3324/haematol.2016.157008

Received: September 22, 2016.

Accepted: May 31, 2017.

Pre-published: June 1, 2017.

Correspondence: mn@clin.au.dk

		CODES USED TO DEFINE COHORTS, COMORBIDITIES AND OUTCOMES
Splenectomised		
	Splenectomy	JMA 00, 10, 11, 20, JMA 99, JMA 96
Indications		
	Traumatic rupture of the spleen	S36.0 (ICD-10), 865.9 (ICD-8)
	Hematopoietic cancer	C81–C96 (ICD-10); 200-207, 275.59 (ICD-8)
	Hereditary hemolytic anaemias	D55–D58 (ICD-10); 282 (ICD-8)
	ITP	D69.3 (ICD-10); 287.10 (ICD-8)
	Other or nonspecific thrombocytopenia	D69.4, D69.5, D69.6 (ICD-10); 287.11, 287.18, 287.19, 287.29 (ICD-8)
	Splenic diseases and splenomegaly	D73, R16.1, R16.2 (ICD-10); 289.4, 782.89 (ICD-8)
	Abdominal cancers	C16, C18 (ICD-10); 151, 153 (ICD-8)
	Other or unknown indications	Various codes (ICD-10); various codes (ICD-8)
Outcomes		
	Pulmonary hypertension	127.0, 127.2, 127.8, 127.9 (ICD-10); 426 (ICD-8)
	Coronary artery disease	120.0-125.9 (ICD-10); 410-414 (ICD-8)
	Myocardial infarction	I21 (ICD-10); 410 (ICD-8)
	Stroke, ischemic	I63, I64 (ICD-10); 432-434 (ICD-8)
	Stroke, haemorrhagic	I61, (ICD-10); 432 (ICD-8)
Comorbidities		
	COPD	J41-J47 (ICD-10); 490-493, 518 (ICD-8)
	Pulmonary Embolus	I26 (ICD-10); 450.99 (ICD-8)
	Heart failure	I50, I11.0 (ICD-10); 428.99 (ICD-8)
	Obesity (BMI)	E66 (ICD-10); 277 (ICD-8)