

## Myelodysplasia and liver disease extend the spectrum of RTEL1 related telomeropathies

Regulator of telomere elongation helicase 1 (RTEL1) is a DNA helicase involved in telomere maintenance.<sup>1,2</sup> Germline biallelic *RTEL1* variants have been previously reported in a subset of patients with dyskeratosis congenita (DC) and its severe variant Hoyeraal-Hreidarsson syndrome (HH).<sup>3-6</sup> Furthermore, germline heterozygous *RTEL1* variants have been linked to a subset of patients

with pulmonary fibrosis.<sup>2,7,8</sup> We have undertaken sequencing analysis (whole exome and targeted<sup>9</sup>) of *RTEL1*, using genomic DNA extracted from peripheral blood of 429 patients from our international bone marrow failure registry which includes DC, HH, aplastic anemia (AA), and familial myelodysplasia/leukemia (MDS/AML). This has revealed that 35 out of the 429 patients have *RTEL1* variants (Table 1). Based on the minor allele frequency in the population reported on the Exome Aggregation Consortium database (ExAC – <http://exac.broadinstitute.org/>), the type of variant (missense, nonsense and indels), telomere length, the

**Table 1.** *RTEL1* variants identified in 35 index cases.

	Index cases	Diagnosis	Age (years)	Gender	Length T/S ratio (percentile)	DNA Change	Protein Change	ExAC Frequency <sup>9</sup>	CADD	
<b>Biallelic</b>										
Likely pathogenic	1	AA	36	F	0.46 (1 <sup>st</sup> )	c.2942G>A (homozygous)	p.R981Q	1 in 119934	28.4	
	2	AA	12	F	0.49 (<10 <sup>th</sup> )	c.3286G>T (homozygous)	p.G1096W	1 in 118164	26.2	
	3	DC	14	M	0.4 (<1 <sup>st</sup> )	c.2300G>A (homozygous)	p.R767Q	NR	27.7	
	4	DC	15	F	0.64 (<10 <sup>th</sup> )	c.2785_2787delCAG (heterozygous)	p.Q929del	NR	12.9	
Bystander						c.2992C>T (heterozygous)	p.R998*	2 in 119914 (1 in 59,957)	37	
	5	HH	2	F	0.45 (<1 <sup>st</sup> )	c.1716C>G (homozygous)	p.I572M	NR	24.9	
<b>Heterozygous</b>										
<b>LOF</b>										
Likely pathogenic	6	DC	77	M	0.5 (<10 <sup>th</sup> )	c.3028C>T	p.R1010*	10 in 119716 (1 in 11,972)	34	
	7	MDS	45	F	0.47 (<10 <sup>th</sup> )	c.3028C>T	p.R1010*	10 in 119716 (1 in 11,972)	34	
	8	MDS	55	M	0.73 (<50 <sup>th</sup> )	c.2992C>T	p.R998*	2 in 119914 (1 in 59,957)	37	
Unknown significance	9	MDS	54	M	0.48 (<10 <sup>th</sup> )	c.3012_3028del	p.Q1005Kfs*80	NR	34	
	10	AML	23	F	1.14 (>50 <sup>th</sup> )	c.3464C>T	p.T1155M	5 in 117408 (1 in 23,482)	12.94	
	11	MDS	20	F	0.95 (<50 <sup>th</sup> )	c.2965G>C	p.E989Q	3 in 119938 (1 in 39,980)	22	
	12	DC	10	M	0.39 (<1 <sup>st</sup> )	c.2723C>G	p.P908R	1 in 119146	0.001	
	13	DC	24	M	0.99 (>50 <sup>th</sup> )	c.208C>T	p.R70C	10 in 120456 (1 in 12,046)	25.3	
	14	AA	24	F	0.65 (<10 <sup>th</sup> )	c.2941C>T	p.R981W	6 in 119930 (1 in 19,988)	33	
	15	AA	6	F	1.04 (>50 <sup>th</sup> )	c.2351C>T	p.A784V	6 in 118274 (1 in 19,712)	2.88	
	16	AA	28	F	1.56 (>90 <sup>th</sup> )	c.3595G>A	p.G1199R	4 in 107372 (1 in 26,843)	5.246	
	17	DC	8	F	0.98 (>90 <sup>th</sup> )	c.1603A>G	p.I535V	NR	9.212	
	18	DC	18	M	0.95 (<50 <sup>th</sup> )	c.3430G>A	p.V1144M	NR	23.6	
	19	AA	28	M	0.56 (<10 <sup>th</sup> )	c.4129A>G <sup>†</sup>	p.T1377A	NR	1.406	
	20	AA	10	F	1.01 (>50 <sup>th</sup> )	c.3608G>A	p.S1203N	NR	23.5	
	Bystander	21	DC	16	F	0.47 (<10 <sup>th</sup> )	c.1991G>T	p.G664V	NR	26.1
		22	DC	18	M	1.21 (>50 <sup>th</sup> )	c.2618G>A	p.G873D	249 in 19124 (1 in 77)	10.29
		23	DC	37	M	0.64 (<10 <sup>th</sup> )	c.2516G>T	p.S839I	126 in 71024 (1 in 564)	17.05
		24	AA	4	F	1 (>50 <sup>th</sup> )	c.3047C>T	p.P1016L	184 in 119184 (1 in 648)	10.85
		25	DC	50	F	0.94 (<50 <sup>th</sup> )	c.3047C>T	p.P1016L	184 in 119184 (1 in 648)	10.85
26		DC	NA	M	NA	c.3047C>T	p.P1016L	184 in 119184 (1 in 648)	10.85	
27		MDS/AML	61	F	0.63 (<10 <sup>th</sup> )	c.3128A>G	p.Q1043R	151 in 118626 (1 in 786)	0.276	
28		DC	4	M	0.85 (<50 <sup>th</sup> )	c.3992G>A <sup>†</sup>	p.R1331Q	120 in 101400 (1 in 845)	12.7	
29		DC	3	M	1.51 (>90 <sup>th</sup> )	c.3992G>A <sup>†</sup>	p.R1331Q	120 in 101400 (1 in 845)	12.7	
30		HH	0	M	0.69 (<50 <sup>th</sup> )	c.2734G>C	p.V912L	85 in 117986 (1 in 1,388)	6.325	
31		DC	54	M	0.47 (<10 <sup>th</sup> )	c.4159C>T <sup>†</sup>	p.P1387S	71 in 110950 (1 in 1,563)	24.7	
32		AA	7	M	1.34 (>50 <sup>th</sup> )	c.4159C>T <sup>†</sup>	p.P1387S	71 in 110950 (1 in 1,563)	24.7	
33		DC	31	M	0.64 (<10 <sup>th</sup> )	c.1261C>G	p.Q421E	71 in 120318 (1 in 1,695)	24.2	
34		AA	34	M	0.6 (<10 <sup>th</sup> )	c.1261C>G	p.Q421E	71 in 120318 (1 in 1,695)	24.2	
35		DC	3	M	1.44 (>50 <sup>th</sup> )	c.3121G>A	p.D1041N	43 in 118650 (1 in 2,759)	14.34	

NR: not reported; CADD PHRED: combined annotation dependent depletion score; AA: aplastic anemia; AML: acute myeloid leukemia; DC: dyskeratosis congenita; HH: Hoyeraal-Hreidarsson syndrome; MDS: myelodysplasia. <sup>†</sup>variant is not in the canonical transcript ENST00000508582 seen in ExAC, but is found in ENST00000482936. Centiles for T/S ratios, established from a healthy control population (n = 202), are as follows: 99<sup>th</sup> centile = 1.99, 90<sup>th</sup> centile = 1.47, 50<sup>th</sup> centile = 0.96, 10<sup>th</sup> centile = 0.68, 1<sup>st</sup> centile = 0.46. Telomeres are considered short if they are at or below the 10<sup>th</sup> centile, and very short if they are at or below the 1<sup>st</sup> centile. <sup>‡</sup>For each of the rare variants reported (less than 10 heterozygotes), the ethnicity of our patient matched at least one reported on the ExAC. NB: Six index cases harbour variants in other known disease genes: index cases 11, 15, 28, 31, 33 and 35 harbour variants in *TERT* (heterozygous c.3197C>T; p.P1066L and c.322C>T; p.R108C), *DNAJC21* (homozygous c.793G>T; p.Q265\*), *TERT* (heterozygous c.1336\_1337insC; p.R446Pfs93\* and c.329G>C; p.G110A), *TERT* (homozygous c.3150G>C; p.K1050N) and *TERC* (heterozygous c.205C>T), *TINF2* (heterozygous c.838A>G; p.K280Q), and *DKC1* (hemizygous c.914A>C; p.K314T), respectively.

Combined Annotation Depletion (CADD) score,<sup>10</sup> and segregation as well as information found in literature, we classified these variants into four different groups: (1) biallelic variants, (2) heterozygous loss of function (LOF) variants, (3) heterozygous missense variants of unknown significance (VUS) and (4) heterozygous missense bystander variants.

As a result, we have further defined the relationship between variants in the *RTEL1* gene and this spectrum of disease. The initial disease association was made when biallelic *RTEL1* variants were shown to cause early onset of a severe form of DC and HH.<sup>3,5,6,11</sup> Here, we describe five new biallelic families (Figure 1A and 1B), where the variants are believed to be disease-causing in four (Families 1-4). Two families presented with AA, two with DC, and one with HH (Table 1 and Table S1). Interestingly, in one of these the index case presented

with AA in adulthood. In the HH family (Family 5), we believe the homozygous *RTEL1* variant is a bystander as it did not segregate with disease, being homozygous in both the index case, with severe disease in infancy, and in the asymptomatic 25-year-old mother.

Three disease causing heterozygous LOF *RTEL1* variants (nonsense and frameshift deletion) were found in patients from four unrelated families (Figure 1B and 1C), one of whom presented with DC, and the others with MDS and/or liver disease. Therefore, this extends the phenotypes associated with heterozygous loss of function *RTEL1* variants to include late onset of MDS and liver disease (Figure 1C and Table 2). This combination of hematological and liver disease is very reminiscent of that established for heterozygous variants in another telomere related gene, *TERT*,<sup>12</sup> which can also present with a severe early onset disease when the variants are biallelic.<sup>13</sup>

**Table 2.** Characteristics of families with *RTEL1* LOF variants.

Family	Individuals	Age at study (years)	Gender	Clinical status	Nucleotide	Amino acid	Variant status	Clinical features/diagnosis
6	I-1	NA	F	asymptomatic	NA	NA	NA	None
	I-2	NA	M	asymptomatic	NA	NA	NA	None
	II-1	NA	F	asymptomatic	NA	NA	NA	None
	II-2	77	M	affected	c.3028C>T	p.R1010*	Heterozygous	DC, lacy skin pigmentation, pancytopenia, pulmonary fibrosis, cirrhosis
	II-3	NA	F	asymptomatic	NA	NA	NA	None
	III-1	NA	M	asymptomatic	NA	NA	NA	None
	III-2	54	F	asymptomatic	c.3028C>T	p.R1010*	Heterozygous	None
	III-3	52	F	affected	c.3028C>T	p.R1010*	Heterozygous	Liver disease (non-specific hepatic changes)
	III-4	49	F	asymptomatic	Wild type	Wild type	Wild type	None
	III-5	47	M	asymptomatic	c.3028C>T	p.R1010*	Heterozygous	None
	III-6	NA	F	asymptomatic	NA	NA	NA	None
	IV-1	20	F	asymptomatic	Wild type	Wild type	Wild type	None
	IV-2	17	F	asymptomatic	Wild type	Wild type	Wild type	None
	IV-3	16	F	asymptomatic	c.3028C>T	p.R1010*	Heterozygous	None
	IV-4	23	F	asymptomatic	Wild type	Wild type	Wild type	None
	IV-5	21	F	asymptomatic	Wild type	Wild type	Wild type	None
	IV-6	7	F	asymptomatic	c.3028C>T	p.R1010*	Heterozygous	None
IV-7	4	M	asymptomatic	Wild type	Wild type	Wild type	None	
7	I-1	NA	F	NA	NA	NA	NA	NA
	I-2	NA	M	NA	NA	NA	NA	NA
	III-1	45	F	affected	c.3028C>T	p.R1010*	Heterozygous	MDS, nail dystrophy
8	I-1	NA	F	asymptomatic	NA	NA	NA	None
	I-2	NA	M	asymptomatic	NA	NA	NA	None
	II-1	55	M	affected	c.2992C>T	p.R998*	Heterozygous	MDS (low risk), cirrhosis
	II-2	47	M	affected	c.2992C>T	p.R998*	Heterozygous	MDS (low risk), cirrhosis
	II-3	NA	F	asymptomatic	NA	NA	NA	None
9	I-1	46	F	affected	NA	NA	NA	Liver and lung disease
	I-2	NA	M	asymptomatic	NA	NA	NA	None
	II-1	NA	M	asymptomatic	NA	NA	NA	None
	II-2	NA	F	asymptomatic	NA	NA	NA	None
	III-1	NA	F	asymptomatic	NA	NA	NA	None
	III-2	54	M	affected	c.3012_3028del	p.E1005Kfs*80	Heterozygous	MDS (low risk), interstitial lung disease, cirrhosis, osteoporosis, baldness and psoriasiform skin
	III-3	49	F	asymptomatic	Wild type	Wild type	Wild type	None
	IV-1	32	F	asymptomatic	c.3012_3028del	p.E1005Kfs*80	Heterozygous	None
IV-2	30	F	asymptomatic	Wild type	Wild type	Wild type	None	


DC: dyskeratosis congenita; MDS: myelodysplasia; NA: not available; F: female; M: male.

The families we present clearly illustrate the variable penetrance of heterozygous *RTEL1* variants. This is exemplified by Family 6 (Figure 1C, p.R1010\*) where the index case had DC features, which did not become apparent until age 77 years. His daughter had liver disease at the age of 52 years, and segregation analysis identified four asymptomatic carriers aged below 50 years. This family highlights not only variable penetrance of heterozygous LOF variants, but also suggests a late onset disease predisposition. The same *RTEL1* variant was identified in Family 7 (Figure 1C and Table 2), where it was associated with MDS and nail dystrophy in the 45-year-old index case. Interestingly, this same variant is reported by Ballew *et al.*<sup>6</sup> in a heterozygous state as being the cause of HH in two siblings (aged three and one years) with very short telomeres. In the family in question, the mother also harboured the variant, had short telomeres but was asymptomatic. Indeed, in most of the families where the index case has disease due to biallelic *RTEL1* variants, both here and in previous reports, the heterozygous parents are generally asymptomatic. However, we now note that these individuals may nevertheless be predisposed to developing disease in their later years. This is suggested by Family 2 (Figure 1A, p.G1096W) where there is a history of pulmonary disease in the grandmother in her 70s and for the p.R998\* variant, which has been seen in both severe recessive<sup>3,5,6</sup> and late onset dominant settings (Families 4 and 8 and Cogan *et al.*<sup>7</sup>). Thus, it is important to be careful when counselling families.

We have identified 14 unrelated patients (Table 1 and Online Supplementary Table S2) with nine heterozygous

missense variants we believe to be bystanders due to their occurrence at an allele frequency of more than 1 in 3,000 in the ExAC population. Additionally, 12 unrelated patients with DC (n=5), AA (n=5), MDS (n=1) and AML (n=1) were found to harbour rare heterozygous missense variants. We have classified these as variants of unknown significance (VUS) as they are either not seen in the ExAC population or are present at an allele frequency of less than 1 in 10,000 (Table 1 and Online Supplementary Table S2). The average CADD score for these VUS (average 15.43, range 0.001 – 33), is lower compared to those that we believe to be disease-causing (average 30.13, range 12.9 – 37, Table 1). We also note that of the 15 heterozygous *RTEL1* variants previously reported to be associated with pulmonary disease,<sup>2,7,8</sup> eight of them are missense.

We have measured telomere lengths by monochrome multiplex quantitative PCR<sup>14</sup> in peripheral blood DNA from all patients except for one who had poor DNA quality (Table 1 and Figure 1D). In agreement with previous studies reporting the impact of *RTEL1* variants on telomere length, we observed that patients with biallelic variants and those with heterozygous loss of function variants had significantly shorter telomeres than controls, as determined by the age-adjusted T/S ratio ( $P=0.0005$  and  $P=0.003$  respectively, 1-way ANOVA with Dunn's multiple comparison test). The median age adjusted T/S ratio for the biallelic group is below the 1<sup>st</sup> centile (-0.6 compared with -0.54), and for the LOF group it is below the 10<sup>th</sup> centile (-0.43 compared with -0.34). It is interesting to note that for the VUS variants, there appears to be two subgroups. The lower four points correspond to p.G664V, p.P908R, p.R981W and p.T1377A. Three of



**Figure 1. The segregation, location and impact of *RTEL1* variants.** (A) Families with biallelic *RTEL1* variants and sequencing traces of index cases (homozygous, Families 1, 2, 3 and 5; compound heterozygous, Family 4). The genotyping is described as follows: wild-type (+/+), heterozygous (+/-) or biallelic (-/-). The age at study is given in years. Affected individuals are coloured in black. NA: not available. (B) *RTEL1* protein (NP\_116575.3) schematic showing the location of the biallelic variants in blue, and the heterozygous LOF variants in red. Conserved protein domains include the P-loop NTPase (yellow); the Rad3 domain (green) which includes the DEAD2 domain (red) and the Helicase C-terminal domain (purple); Harmonin N-like domain (blue); PIP-box – the proliferating cell nuclear antigen interacting protein domain (black). (C) Families with heterozygous LOF *RTEL1* variants and sequencing traces of index cases, annotated as in panel A. (D) Age adjusted telomere length values (delta-tel) were measured by subtracting the observed T/S ratio from the expected T/S ratio, using the equation derived from the line of best fit through the plot of T/S ratios from healthy control samples against age. Patients with *TERC* variants are included as a group with known short telomeres. Centiles were calculated from the control delta-tel values as follows: 99<sup>th</sup> centile = 0.95, 90<sup>th</sup> centile = 0.42, 50<sup>th</sup> centile = 0.06, 10<sup>th</sup> centile = -0.34, 1<sup>st</sup> centile = -0.54. The different genotypes are represented as follows, *TERC*-circles (n=44); biallelic-squares (n=5); loss of function (LOF)-triangles (n=6); variants of unknown significance (VUS)-diamonds (n=12); bystanders-inverted triangles (n=13); controls-grey squares (n=202). (E) T-circle amplification using Phi29 polymerase detected by Southern Blot analysis. Samples: p.R70C - patient with sporadic DC carrying this variant of unknown significance (patient 13 in Table 1); p.R998\* - proband of Family 8 carrying this LOF variant (Table 1); positive control - genomic DNA extracted from WI-38 VA-13 cells, known to produce T-circle.

these variants affect key domains within the protein and may impact on the function of RTEL1 (Figure 1B). These are the helicase C domain (G644V) and the harmonin domain (P908R and R981W).

We previously reported a recurrent missense variant p.R981W as a compound heterozygote in three young unrelated probands (under 12-years-old) with HH.<sup>5</sup> Here, we observed the same variant in a heterozygous state in a 24-year-old patient with AA from a consanguineous family (patient 14 in Table 1 and *Online Supplementary Table S2*). In this case, there is no strong evidence that this variant p.R981W is the cause of AA on account of the relatively high frequency of this variant in the ExAC population (6 in 119930 alleles). However, we do note the short telomeres in this patient and the very high CADD score of this variant, indicating the possibility that it acts as a risk factor for disease.

When a patient presents with an *RTEL1* variant, it is difficult to know whether or not it should be considered pathogenic as there are a multitude of rare coding *RTEL1* variants in the population at large. Using the ExAC database, the sum of the number of very rare heterozygous coding alleles (at a frequency of <0.0001) is 1,195 in an average of approximately 56,700 people. This is significantly lower than the number of very rare coding variants that we have identified in our cohort (22 in 429 patients, Fisher's exact test,  $P = 0.003$ ), but on a case-by-case basis, this background poses a problem. In addition to looking at the ExAC database for population frequency, there are several parameters that we have used to assign pathogenic status. The association of the rare variant with the pathology is a given if the patient under review is presenting with one of the *RTEL1* related disease features. Telomere length measurement is now widely used and our experience here is that the heterozygotes, who are often more elderly, may have telomere lengths that are short, but not necessarily very short. We have also looked at T-circles<sup>15</sup> and shown that, in some cases, their presence is clearly increased where there is a loss of function variant compared to a common missense variant (Figure 1E). However, this test is not easy to perform, and a normal range has not been established. The *in silico* prediction tools are helpful and improving, but remain a guide rather than a definitive test. Finally, the segregation of the variant with disease can be decisive. This is more often the case in exclusion rather than inclusion, as we show in Family 5 (Figure 1A and *Online Supplementary Table S4*).

In summary, our study reports on several important observations. Firstly, heterozygous LOF *RTEL1* variants are associated with myelodysplasia and liver disease in adulthood. Secondly, biallelic *RTEL1* variants can present with just bone marrow failure in adulthood. Thirdly, many heterozygous variants, and even some biallelic *RTEL1* variants, are bystanders. Therefore, in order to assign an accurate status to each *RTEL1* variant, detailed clinical and laboratory studies are necessary.

Shirley R. Cardoso,<sup>1\*</sup> Alicia C.M. Ellison,<sup>1</sup>  
Amanda J. Walne,<sup>1</sup> David Cassiman,<sup>2</sup> Manoj Raghavan,<sup>3</sup>  
Bhuvan Kishore,<sup>4</sup> Philip Ancliff,<sup>5</sup> Carmen Rodríguez-Vigil,<sup>6</sup>  
Bieke Dobbels,<sup>2</sup> Ana Rio-Machin,<sup>7</sup> Ahad F.H. Al Seraili,<sup>7</sup>  
Nikolas Pontikos,<sup>8</sup> Hemanth Tummala,<sup>1</sup> Tom Vulliamy<sup>1#</sup>  
and Inderjeet Dokal<sup>1#</sup>

\*Joint senior authors

<sup>1</sup>Centre for Genomics and Child Health, Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, UK; <sup>2</sup>Metabolic Centre and Department of

Gastroenterology-Hepatology, University of Leuven and University Hospitals, Belgium; <sup>3</sup>Centre for Clinical Haematology, University Hospitals Birmingham NHS Foundation Trust, UK; <sup>4</sup>Heart of England NHS Foundation Trust, Birmingham, UK; <sup>5</sup>Camelia Botnar Laboratories, Great Ormond Street Hospital, London, UK; <sup>6</sup>Unidad de Hematología y Oncología Pediátricas, Hospital Universitario Miguel Servet, Zaragoza, Spain; <sup>7</sup>Centre for Haemato-Oncology, Barts Cancer Institute, Queen Mary University of London, UK and <sup>8</sup>UCL Genetics Institute, University College London, UK

*Acknowledgments: the authors would like to thank all the clinicians and patients who have helped us over the years, particularly Dr Kiliz. Financial support was provided by The Brazilian National Council for Scientific and Technological Development, Bloodwise, Children with Cancer and the Medical Research Council, UK.*

Correspondence: s.r.cardoso@qmul.ac.uk  
doi:10.3324/haematol.2017.167056

*Information on authorship, contributions, and financial & other disclosures was provided by the authors and is available with the online version of this article at www.haematologica.org.*

## References

- Ding H, Schertzer M, Wu X, et al. Regulation of murine telomere length by Rtel: an essential gene encoding a helicase-like protein. *Cell*. 2004;117(7):873-886.
- Kannengiesser C, Borie R, Ménard C, et al. Heterozygous RTEL1 mutations are associated with familial pulmonary fibrosis. *Eur Respir J*. 2015;46(2):474-485.
- Walne AJ, Vulliamy T, Kirwan M, Plagnol V, Dokal I. Constitutional mutations in RTEL1 cause severe dyskeratosis congenita. *Am J Hum Genet*. 2013;92(3):448-453.
- Jullien L, Kannengiesser C, Kermasson L, et al. Mutations of the RTEL1 helicase in a Hoyeraal-Hreidarsson syndrome patient highlight the importance of the ARCH domain. *Hum Mutat*. 2016; 37(5):469-472.
- Deng Z, Glusker G, Molczan A, et al. Inherited mutations in the helicase RTEL1 cause telomere dysfunction and Hoyeraal-Hreidarsson syndrome. *Proc Natl Acad Sci USA*. 2013; 110(36):E3408-3416.
- Ballew BJ, Yeager M, Jacobs K, et al. Germline mutations of regulator of telomere elongation helicase 1, RTEL1, in Dyskeratosis congenita. *Hum Genet*. 2013;132(4):473-480.
- Cogan JD, Kropski JA, Zhao M, et al. Rare variants in RTEL1 are associated with familial interstitial pneumonia. *Am J Respir Crit Care Med*. 2015;191(6):646-655.
- Stuart BD, Choi J, Zaidi S, et al. Exome sequencing links mutations in PARN and RTEL1 with familial pulmonary fibrosis and telomere shortening. *Nat Genet*. 2015;47(5):512-517.
- Walne AJ, Collopy L, Cardoso S, et al. Marked overlap of four genetic syndromes with dyskeratosis congenita confounds clinical diagnosis. *Haematologica*. 2016;101(10):1180-1189.
- Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. A general framework for estimating the relative pathogenicity of human genetic variants. *Nat Genet*. 2014;46(3):310-315.
- Le Guen T, Jullien L, Touzot F, et al. Human RTEL1 deficiency causes Hoyeraal-Hreidarsson syndrome with short telomeres and genome instability. *Hum Mol Genet*. 2013;22(16):3239-3249.
- Calado RT, Regal JA, Kleiner DE, et al. A spectrum of severe familial liver disorders associate with telomerase mutations. *PLoS One*. 2009;4(11):e7926.
- Marrone A, Walne A, Tamary H, et al. Telomerase reverse-transcriptase homozygous mutations in autosomal recessive dyskeratosis congenita and Hoyeraal-Hreidarsson syndrome. *Blood*. 2007;110(13):4198-5205.
- Cawthon RM. Telomere length measurement by a novel monochrome multiplex quantitative PCR method. *Nucleic Acids Res*. 2009;37(3):e21.
- Zellinger B, Akimcheva S, Puizina J, et al. Ku suppresses formation of telomeric circles and alternative telomere lengthening in Arabidopsis. *Mol Cell*. 2007;27(1):163-169.