

Pro-inflammatory proteins S100A9 and tumor necrosis factor- α suppress erythropoietin elaboration in myelodysplastic syndromes

Thomas Cluzeau,^{1,2,3} Kathy L. McGraw,¹ Brittany Irvine,¹ Erico Masala,⁴ Lionel Ades,^{3,5} Ashley A. Basiorka,⁶ Jaroslaw Maciejewski,⁷ Patrick Auburger,² Sheng Wei,¹ Pierre Fenaux,^{3,5} Valeria Santini⁴ and Alan List¹

¹H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL, USA; ²Cote d'Azur University, INSERM U1065, Centre Méditerranéen de Médecine Moléculaire, Nice, France; ³Groupe Français des Myélodysplasies, Paris, France; ⁴Hematology Unit, AOU Careggi, Firenze, Italy; ⁵Senior Hematology Unit, Saint Louis Hospital, Paris, France; ⁶H. Lee Moffitt Cancer Center and Research Institute and the Cancer Biology Ph.D. Program, University of South Florida, Tampa, FL, USA and ⁷Cleveland Clinic, Taussig Cancer Institute, Cleveland, OH, USA

©2017 Ferrata Storti Foundation. This is an open-access paper. doi:10.3324/haematol.2016.158857

Received: October 24, 2016.

Accepted: September 28, 2017.

Pre-published: October 5, 2017.


Correspondence: cluzeau.t@chu-nice.fr

Supplemental figures legends


Supplemental figure 1 : Relationship between inflammatory protein and Epo concentrations in patient serum.

(A) Correlations between Epo and S100A9, (B) Correlations between Epo and TNF α ,
(C) Correlations between TNF α and S100A9. All data are represented on scatter plots.


A.



B.



C.



Supplemental Figure 1. Cluzeau et al.