



EUROPEAN
HEMATOLOGY
ASSOCIATION



Ferrata Storti
Foundation

WT1 loss attenuates the TP53-induced DNA damage response in T-cell acute lymphoblastic leukemia

Fulvio Bordin,^{1*} Erich Piovan,^{1,2*} Elena Masiero,² Alberto Ambesi-Impiombato,^{3,4} Sonia Minuzzo,¹ Roberta Bertorelle,² Valeria Sacchetto,² Giorgia Pilotto,¹ Giuseppe Basso,⁵ Paola Zanovello,^{1,2} Alberto Amadori,^{1,2} and Valeria Tosello²

¹Dipartimento di Scienze Chirurgiche, Oncologiche e Gastroenterologiche, Università degli Studi di Padova, Italy; ²U.O.C. Immunologia e Diagnostica Molecolare Oncologica, Istituto Oncologico Veneto IOV - IRCCS, Padova, Italy; ³Institute for Cancer Genetics, Columbia University, New York, NY, USA; ⁴PsychoGenics Inc., Tarrytown, New York, NY, USA and ⁵Dipartimento di Salute della Donna e del Bambino, Università degli Studi di Padova, Italy

*FB and EP contributed equally to this work.

Haematologica 2018
Volume 103(2):266-277

ABSTRACT

Loss-of-function mutations and deletions in Wilms tumor 1 (*WT1*) gene are present in approximately 10% of T-cell acute lymphoblastic leukemia. Clinically, *WT1* mutations are enriched in relapsed series and are associated to inferior relapse-free survival in thymic T-cell acute lymphoblastic leukemia cases. Here we demonstrate that *WT1* plays a critical role in the response to DNA damage in T-cell leukemia. *WT1* loss conferred resistance to DNA damaging agents and attenuated the transcriptional activation of important apoptotic regulators downstream of TP53 in TP53-competent MOLT4 T-leukemia cells but not in TP53-mutant T-cell acute lymphoblastic leukemia cell lines. Notably, *WT1* loss positively affected the expression of the X-linked inhibitor of apoptosis protein, XIAP, and genetic or chemical inhibition with embelin (a XIAP inhibitor) significantly restored sensitivity to γ -radiation in both T-cell acute lymphoblastic leukemia cell lines and patient-derived xenografts. These results reveal an important role for the *WT1* tumor suppressor gene in the response to DNA damage, and support the view that anti-XIAP targeted therapies could have a role in the treatment of *WT1*-mutant T-cell leukemia.

Correspondence:

valeria.tosello@iov.veneto.it

Received: April 12, 2017.

Accepted: November 15, 2017.

Pre-published: November 23, 2017.

doi:10.3324/haematol.2017.170431

Check the online version for the most updated information on this article, online supplements, and information on authorship & disclosures: www.haematologica.org/content/103/2/266

©2018 Ferrata Storti Foundation

Material published in *Haematologica* is covered by copyright. All rights are reserved to the Ferrata Storti Foundation. Use of published material is allowed under the following terms and conditions:

<https://creativecommons.org/licenses/by-nc/4.0/legalcode>.

Copies of published material are allowed for personal or internal use. Sharing published material for non-commercial purposes is subject to the following conditions:

<https://creativecommons.org/licenses/by-nc/4.0/legalcode>, sect. 3. Reproducing and sharing published material for commercial purposes is not allowed without permission in writing from the publisher.



Introduction

The Wilms tumor 1 (*WT1*) tumor suppressor gene plays an important role in embryonic development and tumorigenesis.¹ The *WT1* protein contains a N-terminal transactivation domain and a C-terminus with four zinc-fingers that can act as an activator or a repressor depending on the cell context.^{2,3} The complexity of the protein is augmented by the generation of alternative isoforms that translates into a different capacity of DNA-binding. In mammals, exon 5 and nine nucleotides at the end of exon 9 are subjected to alternative splicing, generating four different predominant splice isoforms. Alternative splicing of exon 5 gives rise to proteins that differ in the presence or absence of a 17-amino-acid insertion (Ex5⁺ and Ex5⁻). A different splicing, which cuts three amino acids [Lysine, Serine, Threonine, (KTS)] at 3' end of exon 9, generates proteins which vary for the KTS insert (KTS⁺ and KTS⁻). These two isoforms (KTS⁺ and KTS⁻) are conserved in all vertebrates and imbalanced expression of these variants are associated to developmental abnormalities. Importantly, *WT1* isoforms lacking the KTS insertion bind to DNA more efficiently, and significantly affect transcription (reviewed by Hastie⁴). Notably, *WT1* has been demonstrated to interact with a variety of proteins, such as TP53, STAT3 and TET2, that play crucial roles in transformation.⁵⁻⁸

Mutations and deletions in the *WT1* gene were first described in inherited and sporadic Wilms tumors, a pediatric malignancy resulting from the transformation of pluripotent embryonic renal precursor cells.^{9,10} Subsequently, *WT1* gene mutations were also found in acute myeloid and bi-phenotypic leukemia subtypes.¹¹

More recently, *WT1* mutations and/or deletions were also reported in approximately 10% of both pediatric and adult T-cell acute lymphoblastic leukemia (T-ALL).¹² Leukemia-associated *WT1* mutations typically consist of heterozygous frameshift-generating deletions and insertions in exon 7 leading to premature stop codons which may ultimately result in truncated proteins lacking the C-terminal DNA-binding domain or in loss-of-function due to nonsense-mediated RNA decay.¹³ *WT1* mutations are particularly prevalent in patients with relapsed T-ALL,¹⁴ and have been associated with inferior relapse-free survival in cases with standard risk thymic T-ALL.¹⁵ Here we describe a previously unrecognized direct mechanistic role of *WT1* loss in the attenuation of DNA damage-induced apoptosis in T-ALL.

Methods

Cell lines and patient-derived xenografts

MOLT4, PF382 and CCRF-HSB2 T-ALL cells and U2OS cells were obtained from the American Type Culture Collection (ATCC). The P12-Ichikawa T-ALL cells were from the German Collection of Microorganisms and Cell Cultures (Leibniz Institute DSMZ). T-ALL cell lines were cultured *in vitro* with complete

RPMI medium supplemented with 10% FCS (Gibco). T-ALL patient-derived xenografts (T-ALL PDX) had been previously established from pediatric T-ALL samples in non-obese/severe combined immunodeficiency mice (NOD/SCID).^{16,17} T-ALL PDX were expanded *in vivo* via intravenous (i.v.) injection into NOD-*scid*IL2R^{null} immunodeficient mice (NSG mice, Jackson Laboratory). For prolonged *in vitro* culture, T-ALL xenografts were maintained in complete RPMI medium supplemented with 20% FCS, cytokines (10 ng/mL IL-7, 20 ng/mL FLT3-L, and 50 ng/mL SCF, all from Peprotech) and 20 nM insulin (Sigma Aldrich). Procedures involving animals and their care conformed with institutional guidelines and were authorized by the animal ethical committee (Italian Ministry of Health).

Statistical analysis

Results were expressed as mean value±Standard Deviation (SD). Unpaired Student *t*-test was used to analyze data. *P*<0.05 was considered statistically significant.

Results

***WT1* alterations confer resistance to DNA damage in T-ALL cells**

Given the association of *WT1* mutations and loss with relapsed T-ALL, we hypothesized that *WT1* inactivation




Figure 1. *WT1* mutations are associated with increased resistance to γ -radiation-induced apoptosis in T-ALL PDX. (A) Cell viability analysis of *WT1* wild-type *TP53* wild-type (wt-*WT1*/wt-*TP53*), *WT1* wild-type *TP53*-mutant (wt-*WT1*/mut-*TP53*), *WT1*-mutant *TP53* wild-type (mut-*WT1*/wt-*TP53*) and *WT1*-mutant *TP53*-mutant (mut-*WT1*/mut-*TP53*) T-ALL PDX samples after 24 hours (h) from 0.5, 1, 2, 4 and 6 Gray of γ -radiation; analysis is shown as fold change (untreated cells fixed at 1). (B) Group column scatter analysis showing the specific apoptosis in wt-*WT1*/wt-*TP53* and mut-*WT1*/wt-*TP53* T-ALL PDX after 24 h from 1 Gray of γ -radiation (**P*<0.05). (C) Representative FACS analysis of apoptosis, using Annexin-V/Sytox Red staining of wt-*WT1*/wt-*TP53* (n=2; sample ns. 8 and 12) and mut-*WT1*/wt-*TP53* (n=2; sample ns. 48 and 51) T-ALL PDX; apoptosis analysis was performed at 24 h from 1 Gray of γ -radiation.

could result in impaired response to DNA damaging agents in this disease. To test this, we investigated the effects of γ -radiation in a panel of T-ALL patient-derived xenografts (T-ALL PDX) including both *WT1* wild-type [sample ns. 8, 9, 10, 11, 12, 15, previously obtained from T-ALL cells at diagnosis; sample 46R, previously obtained from T-ALL cells at relapse (R)] and *WT1*-mutant (sample ns. 48 and 51, and sample ns. 47R and 51R). Consistent with previous reports, all *WT1* mutations in these samples consisted of truncating nonsense or frameshift alterations in exon 7 (Table 1). Of note, only 2 of these specimens (samples 46R, *WT1* wild-type and 47R, *WT1*-mutant) presented inactivating mutations in *TP53* (Table 1). Additional data, such as *NOTCH1*, *FBXW7* mutations and *PTEN* expression, that are frequently altered in T-ALL samples, showed that alterations were homogenously distributed amongst the *WT1* wild-type and *WT1*-mutant

specimens (Online Supplementary Table S1).

Cell viability assays in response to different doses of γ -radiation (0.5, 1, 2, 4 and 6 Gray) divided these T-ALL PDX into sensitive (median lethal dose = LD50<1.5 Gray) and resistant (LD50>1.5 Gray) (Figure 1A). Importantly, the T-ALL PDX resistant to DNA damage included all the *WT1*-mutant *TP53* wild-type xenografts together with *TP53*-mutant samples 46R and 47R. Sample ns. 9 and 16 also clustered in the resistant group which, in contrast, did not present deletions in *TP53* and *WT1* loci, as assessed by Array-based Comparative Genomic Hybridization analysis (*data not shown*). In line with cell viability assays, apoptosis analysis at 24 hour (h) following 1 Gray of γ -radiation showed increased apoptosis in *WT1* wild-type *TP53* wild-type samples compared with *WT1*-mutant *TP53* wild-type tumors ($P<0.05$) (Figure 1B and C). Overall, these results showed an association between




Figure 2. *WT1* loss protects from DNA damage in MOLT4 T-ALL cells. (A) Western blot analysis of *WT1*-knockdown in MOLT4 cells (upper panel); Western blot quantification of *WT1* protein is reported on top of the gel. Values are normalized with respect to the loading control and relative intensity signal calculated with respect to Control (Ctrl) cells (fixed at 1). Cell viability assay and apoptosis analysis in MOLT4 cells infected either with sh-Scramble (Ctrl) or shRNA-*WT1* (*WT1*-KD) after 24 hours (h)-treatment with increasing doses of γ -radiation (left and right panels, respectively). Quantitative data are shown as mean \pm Standard Deviation (SD); assays were performed in triplicates and reproduced at least three times. * $P<0.05$; ** $P<0.005$; *** $P<0.001$. (B) FACS analysis of active proliferation (S-phase, EDU positive) in Ctrl and *WT1*-KD MOLT4 cells through direct measurement of DNA synthesis. The Click-iT[®] EdU Flow Cytometry Assay kit was used in combination with cell cycle analysis performed with Sytox Red staining. Analysis was performed after 12 h from 2 Gray of γ -radiation. Analysis is shown as fold change (untreated cells fixed at 1). Quantitative data are shown as mean \pm SD; assays were performed in triplicates. Two independent experiments were performed. (Right) A representative experiment is shown. Square regions identify cells in active proliferation.

WT1 mutations and resistance to DNA damage, thus suggesting a putative role of WT1 in DNA damage response.

In order to further investigate a potential mechanistic role of WT1 in DNA damage-induced apoptosis, we performed WT1-knockdown in MOLT4 T-ALL cells (Figure

2A, left panel). We used MOLT4 cells that, among several T-ALL cell lines tested, were reported to be wild-type for TP53 locus, even though results were contradictory.¹⁹ Sequencing analysis of our MOLT4 cells showed a heterozygous nonsense mutation in TP53 locus (R306*) that




Figure 3. WT1-knockdown affects the expression of apoptosis-related proteins following γ -radiation and chemotherapy. (A) Whole cell extracts from Control (Ctrl) and WT1-knockdown (WT1-KD) MOLT4 cells after 24 hours (h) from 6 Gray of γ -radiation were hybridized to human apoptosis arrays. Apoptotic proteins that were consistently modulated are labeled in red (left panel). Immunoblot analysis of cleaved-Caspase3 and XIAP in Ctrl or WT1-KD MOLT4 cells; analysis was performed after 24 h from 6 Gray of γ -radiation. β -Actin is shown as loading control (right panel). Western blot quantification of XIAP protein is reported on top of the gel. Values are normalized with respect to the loading Ctrl and the numbers represent the relative intensity signal calculated respect to untreated Ctrl cells (fixed at 1). (B) Cell viability assays in Ctrl or WT1-KD MOLT4 cells after 24 h treatment with increasing doses of etoposide (left). Analysis is shown as fold change (untreated cells fixed at 1). Quantitative data are shown as mean \pm Standard Deviation (SD); assays were performed in triplicates and reproduced at least three times. Apoptosis analysis using AnnexinV/PI staining in Ctrl or WT1-KD MOLT4 cells after 24 h treatment with increasing doses of etoposide (middle panel). Error bars represent SD for triplicate experiments. * P <0.05; ** P <0.005; *** P <0.001. Immunoblot analysis of PARP, cleaved-Caspase 3 and XIAP in Ctrl or WT1-KD MOLT4 cells; analysis was performed after 24 h treatment with 0.5 μ M etoposide (right panel). β -Actin is shown as loading Ctrl. (C) Cell viability assay in Ctrl or WT1-KD MOLT4 cells after 24 h treatment with increasing doses of cytarabine (ARA-C) (0.05, 0.25, 0.5 and 1 μ M), vincristine (0.01, 1, 10 and 100 ng/mL), or methotrexate (0.01, 0.1, 0.3 and 3 μ M). Analysis is shown as fold change (untreated cells fixed at 1). Quantitative data are shown as mean \pm SD; assays were performed in triplicates. * P <0.05; ** P <0.005; *** P <0.001.

was not detectable at the cDNA level, as previously described.²⁰ Analysis of cell viability in *WT1*-knockdown and shRNA scramble control cells treated with increased doses of γ -radiation showed significantly increased survival at 24 h in *WT1*-knockdown cells (Figure 2A, left panel). Similarly, analysis of γ -radiation-induced apoptosis showed significant protection from DNA damage-induced programmed cell death in *WT1*-knockdown T-ALL cells (Figure 2A, right panel). Moreover, analysis of cell cycle and DNA synthesis demonstrated a decreased number of cells in S phase in response to γ -radiation in Control but not in *WT1*-shRNA expressing cells (Figure 2B). These results were further validated using the CRISPR-Cas9 strategy that allows us to directly target *WT1* locus leading to complete *WT1*-knockout (Online Supplementary Figure S1A and B, left panels). Overall, these results implicate *WT1* loss with increased resistance to DNA damage in T-ALL.

WT1 resistance to DNA damage results in increased expression of pro-survival factors

To explore potential mechanisms implicated in the attenuation of DNA-damage-induced apoptosis following *WT1* loss, we analyzed the expression of apoptosis-related proteins in Control and *WT1*-knockdown MOLT4 cells

following 6 Gray of γ -radiation. These analyses revealed increased cleaved-Caspase3 in *WT1* control cells compared with *WT1*-knockdown samples (Figure 3A and Online Supplementary Figure S2A). However, early stages of DNA-damage response upstream of *TP53* seem to be intact in the absence of *WT1*, as *TP53* phosphorylation was detected in both *WT1* Control and *WT1*-knockdown T-ALL lymphoblasts. In addition, we noted increased levels of pro-survival factors survivin and XIAP as well as upregulation of *HMOX2* (*HO-2*), an important factor in protection against oxidative stress, in response to γ -radiation in *WT1*-knockdown cells, as demonstrated by peptide array quantification (Online Supplementary Figure S2A). Western blot analysis using different validated antibodies and independent lysates, documented the strong down-regulation of *XIAP* in MOLT4 cells treated with γ -radiation and the effective and sustained abrogation of this effect upon *WT1*-knockdown (Figure 3A, right panel, and Online Supplementary Figure S2B). This result was further confirmed in *WT1*-knockout MOLT4 cells (Online Supplementary Figure S2C). Analysis of the DNA-damage response in response to etoposide, a topoisomerase II inhibitor, showed similar results. In fact, *WT1*-knockdown cells showed increased cell viability and decreased apop-




Figure 4. WT1 loss does not impair DNA damage recognition in MOLT4 cells. (A) Immunoblot of cleaved-Caspase3, P-TP53 (S15), Ac-TP53 (K382), and total TP53. β -Actin is shown as loading Control (Ctrl). (B) Immunoblot analysis of phosphorylated ataxia telangiectasia mutated (P-ATM; S1981), and P-CHK2 (T68) in Ctrl and *WT1*-KD MOLT4 cells following 1, 3 and 6 hours (h) from 6 Gray of γ -radiation. Tubulin is shown as loading Ctrl. Western blot quantification of P-TP53 (S15), Ac-TP53 (K382), TP53, P-ATM and P-CHK2 proteins is reported on top of the gel. Values are normalized with respect to the loading Ctrl. The numbers represent the relative intensity signal calculated with respect to untreated Ctrl cells (fixed at 1). For P-TP53, P-ATM and P-CHK2 proteins, that were not detected in untreated Ctrl cells, the relative intensity signal is calculated respected to 1 h (Ctrl cells fixed at 1). (C) Flow cytometry analysis of Histone H2AX phosphorylation (S139) in combination with PI incorporation and cell cycle analysis. Ctrl and *WT1*-KD cells were treated with 6 Gray of γ -radiation for 10 and 30 minutes. Three independent experiments were performed. One representative experiment is shown.

tosis compared with Controls when treated with increasing doses of etoposide (Figure 3B, left and middle panels). As in the case of γ -radiation, *WT1* inactivation resulted in decreased cleaved-Caspase3 and impaired XIAP downregulation following etoposide treatment (Figure 3B, right panel). In line with these results, increased survival of *WT1* deficient MOLT4 T-ALL cells was also observed after cytarabine (ARA-C), vincristine and methotrexate treatment (Figure 3C).

WT1 loss promotes cell survival by attenuating TP53 apoptotic response

The TP53 pathway is extremely efficient in detecting DNA lesions in cells. Once induced, TP53 regulates the expression of a wide range of genes, leading to the biological outcomes of repair, growth arrest or apoptosis.²¹ We thus analyzed a possible interaction between WT1 and TP53 pathways in Control or *WT1*-knockdown MOLT4

T-ALL cells treated with γ -radiation. Western blot analysis of phosphorylated TP53 (P-TP53, S15), acetylated TP53 (Ac-TP53, K382) and total TP53 demonstrated induction and activation of TP53 protein in both Control and *WT1*-knockdown MOLT4 cells (Figure 4A). Importantly, at early time points, TP53 activation resulted more strongly induced in Control with respect to *WT1*-knockdown MOLT4 cells. These results were confirmed also using *WT1*-knockout cells (*Online Supplementary Figure S3A*). This effect did not depend on impaired recognition of DNA damage due to *WT1* loss. Indeed, phosphorylation of crucial proteins involved in the initial phases of DNA damage recognition and repair subsequent to DNA double strand breaks such as ATM, CHK2 and histone H2A variant X (P-H2AX, S139; also known as γ H2AX), was comparable between Control and knockdown cells treated with γ -radiation (Figure 4B and C, and *Online Supplementary Figure S3B*).





Figure 5. WT1-knockdown impairs TP53 DNA damage response in MOLT4 cells. (A) Relative expression of specific genes involved in the DNA-damage response. Analysis is performed in both Control (Ctrl) and *WT1*-KD cells after different times [3, 6 and 12 hours (h)] from 6 Gray of γ -radiation. Expression is calculated relative to untreated sh-Scramble MOLT4 cells (Ctrl 0 h) fixed as 1. (B) Western blot analysis of CDKN1A, BBC3, BAX and XIAP in Ctrl and *WT1*-KD MOLT4 cells after 1, 3, 6 and 12 h from γ -radiation. β -Actin is shown as loading control. Western blot quantification of CDKN1A, BBC3, BAX and XIAP proteins is reported on top of the gel. Values are normalized respect to the loading control. The numbers represent the relative intensity signal calculated respect to untreated Ctrl cells (fixed at 1). For CDKN1A and BBC3 proteins, that were not detected in untreated Ctrl cells, the relative intensity signal is calculated respect to 3 h and 1 h, respectively (Ctrl cells fixed at 1).

Following these results, we analyzed the expression of genes involved in DNA repair, apoptosis and cell cycle progression in control and *WT1*-knockdown MOLT4 cells under basal conditions and at 3, 6 and 12 h after γ -radiation. These analyses revealed that *WT1*-knockdown

affected the expression of prominent genes involved in apoptosis, including *BAX*, *BBC3*, *FAS* and *GADD45A* ($P < 0.001$) (Figure 5A). Similarly, upregulation of *CDKN1A*, a key TP53 target gene involved in cell cycle regulation, was also impaired in *WT1*-knockdown cells




Figure 6. *WT1*-knockdown does not affect survival in *TP53*-mutated T-cell acute lymphoblastic leukemia (T-ALL) cells. (A) Western blot analysis of *WT1*-knockdown in P12-Ichikawa and PF382 cells (top panels); cell viability analysis of Control (Ctrl) and *WT1*-knockdown P12-Ichikawa and PF382 cells after 24 hours (h) treatment with increasing doses of γ -radiation (middle panels). Analysis is shown as fold change (untreated cells fixed at 1). Western blot quantification of *WT1* protein is reported on top of the gel. Values are normalized respect to the loading. The numbers represent the relative intensity signal calculated with respect to Ctrl cells (fixed at 1). Immunoblot analysis of PARP and XIAP in Ctrl and *WT1*-knockdown P12-Ichikawa and PF382 cells (lower panels); analysis was performed after 24 h from 6 Gray of γ -radiation. β -Actin is shown as a loading control. Western blot quantification of XIAP protein is reported on top of the gel. Values are normalized respect to the loading control. The numbers represent the relative intensity signal calculated respect to untreated Ctrl cells (fixed at 1). (B) Cell viability analysis in Ctrl or *WT1*-KD PF382 cells after 1, 3, 6, 12 and 24 h after 6 Gray of γ -radiation (left panel). Analysis is shown as fold change. Immunoblot of PARP, P-TP53 (S15), Ac-TP53 (K382), and total TP53. β -Actin is shown as loading control (right panel). Western blot quantification of P-TP53 (S15), Ac-TP53 (K382), and TP53, P-ATM and P-CHK2 proteins is reported on top of the gel. Values are normalized respect to the loading control. The numbers represent the relative intensity signal calculated respect to untreated Ctrl cells (fixed at 1). (C) Relative expression of specific genes involved in the DNA damage response. Analysis was performed in both Ctrl and *WT1*-KD after different times (3, 6 and 12 h) from 6 Gray of γ -radiation. Expression is calculated relative to untreated Ctrl PF382 cells (Ctrl 0 h) fixed as 1.

treated with γ -radiation compared with *WT1* scramble shRNA Control ($P < 0.001$). Among these, the BBC3 pro-apoptotic factor showed markedly impaired protein upregulation in *WT1*-knockdown irradiated cells (Figure 5B). We thus hypothesized that *WT1* directly regulates BBC3 expression co-operating with TP53 in the DNA damage response. To test this possibility, we analyzed the effects of overexpression of the *WT1* isoforms on the activity of a *WT1*-regulatory element located in intron 1-2 of the BBC3 gene in a luciferase reporter assay. In these experiments, expression of both *WT1*-(KTS-) isoforms induced an approximate 9-fold increase of BBC3 reporter activity (Online Supplementary Figure S4). Consistent with data in literature (reviewed by Hastie⁴), overexpression of both *WT1*-(KTS+) isoforms determined a weak transcriptional response (approx. 2-fold induction). Luciferase activity was seen to be abolished following transfection of mutant *WT1* (E384*) isoforms, the expression of which was found to be barely detectable by Western blot analysis. Overall, these data suggest that, in response to DNA damage, *WT1* loss de-regulates prominent TP53 targets affecting cell survival.

Viability of *TP53*-mutant T-ALL cell lines is not significantly affected by *WT1* loss when exposed to γ -radiation

In order to further validate the resistance to DNA damage conferred by *WT1* loss in MOLT4 cells, we extended our analysis to additional T-ALL cell lines with known *TP53* status (Online Supplementary Table S2). To this end, we infected CCRF-HSB2, which resulted wild-type for *TP53* with a specific hairpin for *WT1* (*WT1*-KD CCRF-HSB2). These cells showed decreased levels of *WT1* as

compared to control cells (Online Supplementary Figure S5A). Importantly, treatment of CCRF-HSB2 cells with increasing doses of γ -radiation and etoposide, showed that loss of *WT1* conferred increased resistance to apoptosis, even if this was to a lesser extent than MOLT4 cells ($P < 0.005$) (Online Supplementary Figure S5A and B). Notably, Western blot of XIAP protein showed higher levels upon *WT1* loss at 1 μ M etoposide (Online Supplementary Figure S5B, right panel). These experiments were also carried out on *TP53*-mutated T-ALL cell lines, PF382 and P12-Ichikawa, that presented deleterious mutations in the *TP53* gene (Online Supplementary Table S2). Treatment of *TP53*-mutated P12-Ichikawa cells with increasing doses of γ -radiation revealed that *WT1*-knockdown cells were not protected from apoptosis compared to control cells (Figure 6A, top left panel). Molecular analysis showed that cleaved-PARP and XIAP were not significantly differentially expressed between Control and *WT1*-knockdown cells upon DNA damage (Figure 6A, bottom left). Similar results were obtained in *TP53*-mutated PF382 cells (Figure 6A, right panel). As expected from a *TP53*-mutated cell line, PF382 cells displayed *TP53* stabilization in the absence of DNA damage and failed to induce the transcription of crucial apoptotic genes involved in *TP53* response, such as *BAX*, *BBC3*, *FAS*, and *GADD45A*, following γ -radiation treatment (Figure 6B and C). Consistently, loss of *WT1* in PF382 cells did not alter this profile (Figure 6C). *CDKN1A* resulted similarly up-regulated following γ -radiation in Control and *WT1*-knockdown PF382 cells (Figure 6C). In conclusion, differently to MOLT4 and CCRF-HSB2 cells that displayed functional *TP53* activation and *WT1* loss increased cell viability following DNA damage, *TP53*-mutated T-ALL cell

Table 1. *WT1* and *TP53* genetic status in T-cell acute lymphoblastic leukemia PDX. HGVS-nomenclature was used for the description of sequence variants.¹⁸

T-ALL samples	WT1	TP53
8	wt	wt
9	wt	wt
10	wt	wt
11	wt	wt
12	wt	wt
15	wt	wt
16	wt	wt
46R	wt	c.743G>A; p.R248Q
47R	Mut c.1142C>A; p. S381*	Mut c.708_709insC; p.M237fs c.638G>A; p.R213Q.
48	Mut c.1109_1110insGGGCCCGG; p.V371fs	wt
51	Mut c.1102_1104delinsCGGCCACTCCCCGGGGGTCC; p.V368fs	wt
51R	Mut c.1102_1104delinsCGGCCACTCCCCGGGGGTCC; p.V368fs	wt

wt: wild-type; Mut: mutant; R: *.

lines, PF382 and P12 Ichikawa, were not significantly affected by *WT1* loss when exposed to γ -radiation, suggesting a possible cross-talk between WT1 and functional TP53.

Increased resistance of *WT1*-deficient T-ALL cells to DNA damage can be rescued by pre-treatment with XIAP inhibitors

Molecular characterization of DNA damage response in *WT1* deficient MOLT4 cells showed an impaired TP53 response characterized by de-regulation of several proteins involved at different steps of the apoptotic cascade downstream of TP53 activation. To further validate these

results, we molecularly characterized the effects of γ -radiation on *WT1* wild-type *TP53* wild-type T-ALL PDX samples (n=2; sample ns. 8 and 12) and on *WT1*-mutated *TP53* wild-type samples (n=2; sample ns. 48 and 51). All the tumors showed TP53 protein stabilization upon DNA damage (Figure 7A). Importantly, *WT1* wild-type samples showed higher levels of cleaved-Caspase3 compared to *WT1*-mutant tumors. Following γ -radiation, BAX, survivin and HO-2 proteins were not significantly regulated either in *WT1* wild-type nor in *WT1*-mutated T-ALL xenograft samples, while BBC3 was not differentially regulated between *WT1* wild-type and *WT1*-mutated T-ALL, with *WT1*-mutated T-ALL PDX sample n. 51 undergoing a




Figure 7. XIAP is a critical player in *WT1* resistance to DNA damage in T-cell acute lymphoblastic leukemia (T-ALL) cells. (A) Western blot analysis of TP53, cleaved-Caspase3, BBC3, BAX, survivin, HO-2 and XIAP in wt-*WT1* (n=2; sample ns. 8 and 12) and mut-*WT1* (n=2; sample ns. 48 and 51) T-ALL PDX following 3 hours (h) from 6 Gray of γ -radiation. Histogram plots representing Western blot quantification of TP53, cleaved-Caspase3, BBC3, BAX and XIAP proteins is reported on the right. Values are normalized respect to the loading control (Ctrl). The plots show the relative intensity signal calculated with respect to untreated Control cells (fixed at 1). (B) Apoptosis assay in *WT1*-mutated PDX sample n. 48 pre-treated for 24 h with 15 μ M embelin and subjected to γ -radiation or etoposide treatment (0.5 Gray and 0.1 μ M, respectively) in the presence or absence of embelin. (C) Apoptosis assay in MOLT4 cells pre-treated for 24 h with 7.5 μ M or 10 μ M embelin and subjected to γ -radiation or etoposide treatment (2 Gray and 0.2 μ M, respectively). Analysis is shown at 48 h following γ -radiation and presented as percentage of apoptotic cells. **P*<0.05; ***P*<0.005; ****P*<0.001.

strong induction of BBC3 (Figure 7A and *Supplementary Figure S6*). On the other hand, XIAP protein resulted grossly down-regulated following γ -radiation in *WT1* wild-type but not in *WT1*-mutant T-ALL PDX, in line with results obtained in MOLT4 and CCRF-HSB2. XIAP is one of the best characterized members of the inhibitor of apoptosis (IAP) family proteins and is considered to be a key regulator of apoptosis. We thus investigated whether inhibition of XIAP could contribute to rescue the effects of *WT1* loss in T-ALL cells upon DNA damage. T-ALL PDX samples were treated with increasing doses of embelin, a natural compound that binds to XIAP and promotes apoptosis.²² Amongst 5 T-ALL PDX samples analyzed, 2 resulted resistant at 30 μ M embelin (n. 12 and n. 15) whereas T-ALL PDX sample ns. 8, 48 and 51 showed decreased viability and increased apoptosis starting from 15 μ M of embelin (*Online Supplementary Figure S7*). We thus pre-treated *WT1*-mutated PDX sample n. 48 with 15 μ M of embelin, followed by treatment with γ -radiation and etoposide at 0.5 Gray and 0.1 μ M etoposide, respectively. Apoptosis analysis showed a significant rescue of resistance to both γ -radiation and etoposide treatments ($P < 0.001$) (Figure 7B). Importantly, a complete rescue was achieved by γ -radiation and etoposide treatments in MOLT4 cells at 10 μ M embelin. Indeed, apoptosis analysis revealed that pre-treatment with 7.5 μ M and especially 10 μ M embelin before DNA damaging agents (γ -radiation and etoposide), resulted in a significant increase in apoptosis with respect to that induced by γ -radiation or etoposide alone in *WT1*-knockdown MOLT4 cells ($P < 0.001$) (Figure 7C). Notably, XIAP-knockdown in MOLT4 cells deficient for *WT1* (*WT1*-KD) efficiently rescued resistance to apoptosis in the presence of increasing doses of etoposide ($P < 0.001$ at 0.25 μ M etoposide) (*Online Supplementary Figure S8*). Overall, these results identified XIAP as a critical player in regulating DNA damage resistance induced by *WT1* loss, opening new therapeutic opportunities in *WT1*-mutant patients.

Discussion

The introduction of intensive combination chemotherapy protocols in T-ALL treatment has led to remarkable improvements in survival.²³ Unfortunately, primary resistance to chemotherapy or acquired resistance are strongly associated with poor outcome and the molecular mechanisms that allow leukemia cells to resist chemotherapy are not yet completely understood. A recent study has demonstrated that refractory T-ALL clones can develop mutations in drug-resistance genes under chemotherapy pressure.²⁴ Moreover, resistant clones may originate from minor sub-clones present in patients at diagnosis through a dramatic selection during chemotherapy.^{25,26} A better understanding of the genetic lesions that may contribute to chemo-resistance is imperative to improve treatment protocols and for the identification of more effective anti-leukemic drugs. In particular, it is necessary to better characterize genetic alterations associated to genes with poorly characterized function.

The *WT1* gene is mutated in approximately 10% of T-ALL and AML cases, and these alterations are mainly heterozygous frameshift mutations that cluster in exon 7 and are predicted to lead to a truncated protein which lacks the zinc finger domain and the property to bind DNA.^{11,12,27}

Monoallelic or subclonal deletions have also been observed in some cases in association with mutations.^{12,28,29} Even if *WT1* alterations clearly suggest a role as a tumor suppressor mediated at least in part by de-regulation of its transcriptional activity, its role in T-ALL is still poorly understood. Most studies have focused their attention on acute myeloid leukemia (AML). Recently, *WT1* inactivating mutations have been found to inversely correlate with *TET2/IDH1/IDH2* mutations; indeed, *WT1*-mutant AML patients have reduced 5-hydroxymethyl-cytosine levels, similar to *TET2/IDH1/IDH2* mutant AML, suggesting a critical role for *WT1* as an epigenetic regulator.⁷ Moreover, always in the context of AML, *WT1* mutations were found to be associated with an increased risk of relapse and an inferior outcome, even if contradictory results have been reported,³⁰⁻³⁶ probably due to the difference between the AML subgroups analyzed. Interestingly, a case report of AML demonstrated that a minor sub-clone, characterized by a mutation in the *WT1* gene, emerges as a dominant clone after repetitive chemotherapy, including stem cell transplant.³⁷ In T-ALL, studies on the role of *WT1* mutations are still limited. In pediatric and adult T-ALL, the presence of *WT1* mutations in the entire cohort were not predictive of poor clinical outcome.^{12,27} Interestingly, one study observed that within the standard risk group of thymic T-ALL, the small group of *WT1*-mutant patients had inferior relapse-free survival compared to wild-type patients.¹⁵ Notably, a very recent study in a group of high-risk T-ALL composed of paired diagnosis and relapse samples reported that *WT1* mutations resulted highly enriched with respect to unselected T-ALL cases.¹⁴ Moreover, *WT1* was shown to be among the frequently altered genes in the Early T Progenitor ALL (ETP) subgroup.³⁸ ETP ALL were originally identified as a high-risk subtype of leukemia that lacks several T-cell markers and is characterized by aberrant expression of myeloid and stem cell markers.³⁹

Our study highlights an important role for *WT1* as a critical regulator of the *TP53*-dependent apoptotic response following DNA damage, and indicates an association between *WT1* loss-of-function and resistance both in T-ALL PDX and MOLT4 T-ALL cells. In fact, in the presence of DNA-damaging stimuli, such as γ -radiation and chemotherapy, loss of *WT1* in MOLT4 cells conferred increased survival associated with an impaired induction of *TP53* apoptotic response. Amongst the de-regulated genes, some of which have been previously described as *WT1* targets, such as *BAX* and *GADD45A*, we identified *BBC3* as a new important *WT1* target. *BBC3*, which encodes for the pro-apoptotic factor Puma, is a potent antagonist of anti-apoptotic BCL2-related proteins contributing to the release and activation of the pro-apoptotic factors BAX and BAK. Although *BBC3* was found to be a critical regulator of apoptosis in primary thymocytes in response to DNA damage and glucocorticoid treatment⁴⁰ and a crucial factor in MOLT4 cells following *WT1* loss, one mut-*WT1* T-ALL xenograft in our series (n. 51) showed strong induction of *BBC3* much like the wt-*WT1* PDX samples, suggesting that other pro-apoptotic factors may be important in this sample.

An analysis of the *WT1*-knockout mouse first pointed out the involvement of *WT1* in regulating apoptosis, which demonstrated that lack of *WT1* expression determined severe abnormalities of renal development and that the metanephric blastema of *WT1*-null embryos was char-

acterized by massive apoptosis compared to wild-type embryos.⁴¹ A number of subsequent studies have demonstrated that WT1 can directly or indirectly regulate several BCL-2 family members including pro-apoptotic factors, such as BAK and BAX, and the anti-apoptotic proteins BCL2 and BCL2A1 with different outcomes depending on the cell context (reviewed by Toska and Roberts² and by Yang *et al.*³). Contradictory results regarding WT1 regulation may be due in part to the complex structure of the WT1 protein and its multiple isoforms. Another level of complexity is due to co-operativity between WT1 and its interacting proteins. WT1 was found to physically interact with TP53, and this interaction was demonstrated to modulate the ability of WT1 and TP53 to trans-activate their respective targets.^{5,42,43} In this scenario, it is plausible that the cellular context may substantially determine WT1 effects on specific targets. In our study, in fact, WT1 significantly directly or indirectly affects the transcription of important mediators of TP53-apoptotic response only in the presence of a functional TP53. MOLT4 cells carry a heterozygous nonsense mutation in *TP53* locus (p.R306*) that was not detectable at the cDNA level as previously described by others;²⁰ concerning this trait, unlike the majority of T-ALL cell lines, MOLT4 cells resemble primary T-ALL samples at diagnosis as they rarely display deleterious mutations in the *TP53* gene.⁴⁴ Indeed, treatment of MOLT4 cells with DNA-damaging stimuli determines TP53 stabilization and strong induction of TP53-specific target genes. This response is significantly down-regulated when MOLT4 cells are engineered to lose *WT1*, as in knockdown or knockout experiments. Our data suggest that *WT1* loss can influence both stabilization and transcriptional activity of TP53. On the other hand, we found that *TP53*-mutated T-ALL cell lines, such as PF382 and P12-Ichikawa, were not significantly affected by *WT1* loss in response to DNA damage, suggesting a possible link between TP53 and WT1, that remains a subject for future studies. The effects on TP53 response mediated by *WT1* loss in MOLT4 cells also resulted in deregulation of downstream effectors such as XIAP. In fact, we found elevated levels of XIAP following γ -radiation and etoposide treatment in *WT1*-deficient MOLT4 cells. Importantly, pre-treatment of *WT1*-deficient MOLT4 cells

with embelin, a natural compound that specifically inhibits XIAP^{22,45} and direct inactivation of XIAP using specific shRNAs, rescued the resistance of *WT1*-deficient MOLT4 towards γ -radiation and etoposide. Notably, in T-ALL PDX samples, as for MOLT4 cells, XIAP stability was maintained following DNA damage in *WT1*-mutant samples but not in *WT1* wild-type xenografts, confirming a prominent role of XIAP downstream of WT1 resistance in T-ALL. In addition, also *WT1*-mutant T-ALL xenografts were seen to be sensitized to γ -radiation and etoposide when the treatments were performed in the presence of the XIAP inhibitor embelin.

XIAP acts by inhibiting caspase activation at an important point where intrinsic and extrinsic apoptotic pathways converge, making it an attractive therapeutic target. Pre-clinical studies concerning the use of embelin and small molecule XIAP inhibitors have demonstrated antitumor effects in xenograft and mouse models.⁴⁶⁻⁴⁹ Importantly, no specific toxicity was reported following XIAP inhibition in these studies.⁴⁶ In conclusion, our study unveils an important role for the *WT1* tumor suppressor gene in dampening TP53 DNA damage response and suggest that the combination of XIAP inhibitors with conventional chemotherapy or radiotherapy in T-ALL patients carrying *WT1* mutations might be beneficial.

Acknowledgments

We are grateful to W. Pear (Perelman School of Medicine, University of Pennsylvania, Philadelphia) for providing MigR1 empty vector; E. Peta for performing some molecular analyses; S. Indraccolo for providing patient-derived xenografts (PDX); V. Agnusdei and M. Pimazza for maintaining PDX samples in immunodeficient mice; Cinzia Candiotto for TP53 mutational analysis in T-ALL cell lines; R. Grancara for computational assistance.

Funding

This work was supported by the Italian Association for Cancer Research (AIRC) grant to VT (MFAG #13053), to AA (IG#14032), to PZ (IG #14256) and to GB (IG#19186); Ministero dell'Istruzione, dell'Università e della Ricerca (MIUR) Ex 60% to E.P.; Progetto di Ricerca di Ateneo (PRAT; #152403) to EP; Istituto Oncologico Veneto 5x1000 fund.

References

1. Miller-Hodges E, Hohenstein P. WT1 in disease: shifting the epithelial-mesenchymal balance. *J Pathol.* 2012;226(2):229-240.
2. Toska E, Roberts SG. Mechanisms of transcriptional regulation by WT1 (Wilms' tumor 1). *Biochem J.* 2014;461(1):15-32.
3. Yang L, Han Y, Suarez Saiz F, Minden MD. A tumor suppressor and oncogene: the WT1 story. *Leukemia.* 2007;21(5):868-876.
4. Hastie ND. Life, sex, and WT1 isoforms--three amino acids can make all the difference. *Cell.* 2001;106(4):391-394.
5. Maheswaran S, Park S, Bernard A, et al. Physical and functional interaction between WT1 and p53 proteins. *Proc Natl Acad Sci USA.* 1993;90(11):5100-5104.
6. Rong Y, Cheng L, Ning H, et al. Wilms' tumor 1 and signal transducers and activators of transcription 3 synergistically promote cell proliferation: a possible mechanism in sporadic Wilms' tumor. *Cancer Res.* 2006;66(16):8049-8057.
7. Rampal R, Alkalin A, Madzo J, et al. DNA hydroxymethylation profiling reveals that WT1 mutations result in loss of TET2 function in acute myeloid leukemia. *Cell Rep.* 2014;9(5):1841-1855.
8. Wang Y, Xiao M, Chen X, et al. WT1 recruits TET2 to regulate its target gene expression and suppress leukemia cell proliferation. *Mol Cell.* 2015;57(4):662-673.
9. Call KM, Glaser T, Ito CY, et al. Isolation and characterization of a zinc finger polypeptide gene at the human chromosome 11 Wilms' tumor locus. *Cell.* 1990;60(3):509-520.
10. Gessler M, Poustka A, Cavenee W, Neve RL, Orkin SH, Bruns GA. Homozygous deletion in Wilms tumours of a zinc-finger gene identified by chromosome jumping. *Nature.* 1990;343(6260):774-778.
11. King-Underwood L, Pritchard-Jones K. Wilms' tumor (WT1) gene mutations occur mainly in acute myeloid leukemia and may confer drug resistance. *Blood.* 1998;91(8):2961-2968.
12. Tosello V, Mansour MR, Barnes K, et al. WT1 mutations in T-ALL. *Blood.* 2009;114(5):1038-1045.
13. Abbas S, Erpelinck-Verschueren CA, Goudswaard CS, Lowenberg B, Valk PJ. Mutant Wilms' tumor 1 (WT1) mRNA with premature termination codons in acute myeloid leukemia (AML) is sensitive to nonsense-mediated RNA decay (NMD). *Leukemia.* 2010;24(3):660-663.
14. Oshima K, Khiabani H, da Silva-Almeida AC, et al. Mutational landscape, clonal evolution patterns, and role of RAS mutations in relapsed acute lymphoblastic leukemia. *Proc Natl Acad Sci USA.* 2016; 113(40):11306-11311.
15. Heesch S, Goekbuget N, Stroux A, et al. Prognostic implications of mutations and

- expression of the Wilms tumor 1 (WT1) gene in adult acute T-lymphoblastic leukemia. *Haematologica*. 2010;95(6):942-949.
16. Agnusdei V, Minuzzo S, Frasson C, et al. Therapeutic antibody targeting of Notch1 in T-acute lymphoblastic leukemia xenografts. *Leukemia*. 2014;28(2):278-288.
 17. Pinazza M, Borga C, Agnusdei V, et al. An immediate transcriptional signature associated with response to the histone deacetylase inhibitor Givinostat in T acute lymphoblastic leukemia xenografts. *Cell Death Dis*. 2016;6:e2047.
 18. den Dunnen JT, Dalgleish R, Maglott DR, et al. HGVS Recommendations for the Description of Sequence Variants: 2016 Update. *Hum Mutat*. 2016;37(6):564-569.
 19. Leroy B, Girard L, Hollestelle A, Minna JD, Gazdar AF, Soussi T. Analysis of TP53 mutation status in human cancer cell lines: a reassessment. *Hum Mutat*. 2014;35(6):756-765.
 20. Ikediobi ON, Davies H, Bignell G, et al. Mutation analysis of 24 known cancer genes in the NCI-60 cell line set. *Mol Cancer Ther*. 2006;5(11):2606-2612.
 21. Riley T, Sontag E, Chen P, Levine A. Transcriptional control of human p53-regulated genes. *Nat Rev Mol Cell Biol*. 2008;9(5):402-412.
 22. Nikolovska-Coleska Z, Xu L, Hu Z, et al. Discovery of embelin as a cell-permeable, small-molecular weight inhibitor of XIAP through structure-based computational screening of a traditional herbal medicine three-dimensional structure database. *J Med Chem*. 2004;47(10):2430-2440.
 23. Pui CH. Recent advances in acute lymphoblastic leukemia. *Oncology*. 2011;25(4):341, 346-347.
 24. Tzoneva G, Perez-Garcia A, Carpenter Z, et al. Activating mutations in the NT5C2 nucleotidase gene drive chemotherapy resistance in relapsed ALL. *Nat Med*. 2013;19(3):368-371.
 25. Mullighan CG, Phillips LA, Su X, et al. Genomic analysis of the clonal origins of relapsed acute lymphoblastic leukemia. *Science*. 2008;322(5906):1377-1380.
 26. Clappier E, Gerby B, Sigaux F, et al. Clonal selection in xenografted human T cell acute lymphoblastic leukemia recapitulates gain of malignancy at relapse. *J Exp Med*. 2011;208(4):653-661.
 27. Renneville A, Kaltenbach S, Clappier E, et al. Wilms tumor 1 (WT1) gene mutations in pediatric T-cell malignancies. *Leukemia*. 2010;24(2):476-480.
 28. Van Vlierberghe P, Homminga I, Zuurbier L, et al. Cooperative genetic defects in TLX3 rearranged pediatric T-ALL. *Leukemia*. 2008;22(4):762-770.
 29. Zhang J, Ding L, Holmfeldt L, et al. The genetic basis of early T-cell precursor acute lymphoblastic leukaemia. *Nature*. 2012;481(7380):157-163.
 30. Bachas C, Schuurhuis GJ, Reinhardt D, et al. Clinical relevance of molecular aberrations in paediatric acute myeloid leukaemia at first relapse. *Br J Haematol*. 2014;166(6):902-910.
 31. Becker H, Marcucci G, Maharry K, et al. Mutations of the Wilms tumor 1 gene (WT1) in older patients with primary cytogenetically normal acute myeloid leukemia: a Cancer and Leukemia Group B study. *Blood*. 2010;116(5):788-792.
 32. Hollink IH, van den Heuvel-Eibrink MM, Zimmermann M, et al. Clinical relevance of Wilms tumor 1 gene mutations in childhood acute myeloid leukemia. *Blood*. 2009;113(23):5951-5960.
 33. Virappane P, Gale R, Hills R, et al. Mutation of the Wilms' tumor 1 gene is a poor prognostic factor associated with chemotherapy resistance in normal karyotype acute myeloid leukemia: the United Kingdom Medical Research Council Adult Leukaemia Working Party. *J Clin Oncol*. 2008;26(33):5429-5435.
 34. Paschka P, Marcucci G, Ruppert AS, et al. Wilms' tumor 1 gene mutations independently predict poor outcome in adults with cytogenetically normal acute myeloid leukemia: a cancer and leukemia group B study. *J Clin Oncol*. 2008;26(28):4595-4602.
 35. Krauth MT, Alpermann T, Bacher U, et al. WT1 mutations are secondary events in AML, show varying frequencies and impact on prognosis between genetic subgroups. *Leukemia*. 2015;29(3):660-667.
 36. Ho PA, Zeng R, Alonzo TA, et al. Prevalence and prognostic implications of WT1 mutations in pediatric acute myeloid leukemia (AML): a report from the Children's Oncology Group. *Blood*. 2010;116(5):702-710.
 37. Nyvold CG, Stentoft J, Braendstrup K, et al. Wilms' tumor 1 mutation accumulated during therapy in acute myeloid leukemia: Biological and clinical implications. *Leukemia*. 2006;20(11):2051-2054.
 38. Liu Y, Easton J, Shao Y, et al. The genomic landscape of pediatric and young adult T-lineage acute lymphoblastic leukemia. *Nat Genet*. 2017;49(8):1211-1218.
 39. Coustan-Smith E, Mullighan CG, Onciu M, et al. Early T-cell precursor leukaemia: a subtype of very high-risk acute lymphoblastic leukaemia. *Lancet Oncol*. 2009;10(2):147-156.
 40. Jeffers JR, Parganas E, Lee Y, et al. Puma is an essential mediator of p53-dependent and -independent apoptotic pathways. *Cancer Cell*. 2003;4(4):321-328.
 41. Kreidberg JA, Sariola H, Loring JM, et al. WT-1 is required for early kidney development. *Cell*. 1993;74(4):679-691.
 42. Idelman G, Glaser T, Roberts CT Jr, Werner H. WT1-p53 interactions in insulin-like growth factor-I receptor gene regulation. *J Biol Chem*. 2003;278(5):3474-3482.
 43. Menke AL, Clarke AR, Leitch A, et al. Genetic interactions between the Wilms' tumor 1 gene and the p53 gene. *Cancer Res*. 2002;62(22):6615-6620.
 44. Hof J, Krentz S, van Schewick C, et al. Mutations and deletions of the TP53 gene predict nonresponse to treatment and poor outcome in first relapse of childhood acute lymphoblastic leukemia. *J Clin Oncol*. 2011;29(23):3185-3193.
 45. Chen J, Nikolovska-Coleska Z, Wang G, Qiu S, Wang S. Design, synthesis, and characterization of new embelin derivatives as potent inhibitors of X-linked inhibitor of apoptosis protein. *Bioorg Med Chem Lett*. 2006;16(22):5805-5808.
 46. Vogler M, Walczak H, Stadel D, et al. Small molecule XIAP inhibitors enhance TRAIL-induced apoptosis and antitumor activity in preclinical models of pancreatic carcinoma. *Cancer Res*. 2009;69(6):2425-2434.
 47. Dai Y, Desano J, Qu Y, et al. Natural IAP inhibitor Embelin enhances therapeutic efficacy of ionizing radiation in prostate cancer. *Am J Cancer Res*. 2011;1(2):128-143.
 48. Lunardi A, Ala U, Epping MT, et al. A clinical approach identifies mechanisms and potential therapies for androgen deprivation resistance in prostate cancer. *Nat Genet*. 2013;45(7):747-755.
 49. Yang T, Lan J, Huang Q, et al. Embelin sensitizes acute myeloid leukemia cells to TRAIL through XIAP inhibition and NF-kappaB inactivation. *Cell Biochem Biophys*. 2015;71(1):291-297.