

**Unsatisfactory efficacy in randomized study of reduced-dose CPX-351 for medically less fit adults with newly diagnosed acute myeloid leukemia or other high-grade myeloid neoplasm**

Roland B. Walter,<sup>1,2,3</sup> Megan Othus,<sup>4</sup> Kaysey F. Orlowski,<sup>1</sup> Emily N. McDaniel,<sup>1</sup> Bart L. Scott,<sup>1,5</sup> Pamela S. Becker,<sup>1,2</sup> Mary-Elizabeth M. Percival,<sup>1,2</sup> Paul C. Hendrie,<sup>2</sup> Bruno C. Medeiros,<sup>6</sup> Michael T. Chiarella,<sup>7</sup> Arthur C. Louie<sup>7</sup> and Elihu H. Estey<sup>1,2</sup>

<sup>1</sup>Clinical Research Division, Fred Hutchinson Cancer Research Center, Seattle, WA; <sup>2</sup>Department of Medicine, Division of Hematology, University of Washington, Seattle, WA; <sup>3</sup>Department of Epidemiology, University of Washington, Seattle, WA; <sup>4</sup>Public Health Sciences Division, Fred Hutchinson Cancer Research Center, Seattle, WA; <sup>5</sup>Department of Medicine, Division of Medical Oncology, University of Washington, Seattle, WA and <sup>6</sup>Stanford University, CA; <sup>7</sup>Jazz Pharmaceuticals, Ewing, NJ, USA

Correspondence: [rwalter@fredhutch.org](mailto:rwalter@fredhutch.org)  
doi:10.3324/haematol.2017.182642

**SUPPLEMENTAL TABLE. Cycle #1 treatment-emergent adverse events with CPX-351 at 32 units/m<sup>2</sup> or 64 units/m<sup>2</sup> per dose**

<b>Parameter</b>	<b>32 units/m<sup>2</sup> n (% cycles)</b>	<b>64 units/m<sup>2</sup> n (% cycles)</b>
Fever, infection		
Bacteremia		
Catheter-related infection	2 (5.3%)	1 (10%)
Cellulitis		1 (10%)
Lung infection	3 (7.9%)	3 (30%)
Neutropenic fever	7 (18.4%)	5 (50%)
Sepsis	2 (5.3%)	1 (10%)
Cardiac		
Atrial tachycardia		1 (10%)
Cardiac arrest	1 (2.6%)	
Tachycardia		1 (10%)
Gastrointestinal		
Mucositis	2 (5.3%)	2 (20%)
General		
Multi-system organ failure		1 (10%)
Immune System		
Anaphylaxis		
Investigations		
Acute kidney injury	4 (10.5%)	2 (20%)
Alanine aminotransferase increase	1 (2.6%)	
Aspartate aminotransferase increase	1 (2.6%)	
Metabolism and nutritional		
Dehydration	1 (2.6%)	
Tumor lysis	1 (2.6%)	1 (10%)
Nervous system disorders		
Intracranial hemorrhage	1 (2.6%)	
Respiratory		
ARDS		1 (10%)
Atelectasis		1 (10%)
Bronchopulmonary hemorrhage	1 (2.6%)	
Hypoxia	3 (7.9%)	2 (20%)
Respiratory failure		1 (10%)
Other		
Deep vein thrombosis	1 (2.6%)	1 (10%)
Leg pain		1 (10%)
Urinary retention		1 (10%)

Table summarizing treatment-emergent grade 3-5 non-hematologic adverse effects that were experienced during treatment cycle #1 and were considered as definitively, probably, or possibly related to study treatment by the investigator.