

Haematologica
HAEMATOL/2017/177808
Version 3

TNF receptor signaling is a driver of chronic lymphocytic leukemia that can be therapeutically targeted by the flavonoid wogonin

Claudia Dürr, Bola S. Hanna, Angela Schulz, Fabienne Lucas, Manuela Zucknick, Axel Benner, Andrew Clear, Sibylle Ohl, Selcen Öztürk, Thorsten Zenz, Stephan Stilgenbauer, Min Li-Weber, Peter H. Krammer, John G. Gribben, Peter Lichter, and Martina Seiffert

Disclosures: This study was supported by the Helmholtz Virtual Institute Understanding and overcoming resistance to apoptosis and therapy in leukemia, by the Else Kroener-Fresenius Foundation, by the German José Carreras Foundation, Grant no. R14/23, and the ERA-NET Transcan-2 Network FIRE-CLL. StSt was supported by the SFB1074 project B1. The authors declare no competing financial interests. This study does not include any patents or copyrights.

Contributions: C.D. designed the study, acquired, analyzed and interpreted data, prepared figures, and wrote the manuscript. B.H., A.S., and F.M. acquired, analyzed and interpreted data. A.C., S.O., and S.Oe. acquired and analyzed data. M.Z. and A.B. analyzed the bead array data and performed statistical analyses. T.Z. and S.S. provided clinical samples and data, advised the study and reviewed the manuscript. M.L.-W. and P.K. advised the wogonin treatment studies and reviewed the manuscript. J.G. provided tissue arrays, advised the study and reviewed the manuscript. P.L. supervised the study and reviewed the manuscript. M.S. designed and supervised the study, analyzed and interpreted data, and wrote the manuscript.